















AN  
ELEMENTARY TREATISE  
ON  
ASTRONOMY.

IN FOUR PARTS.

CONTAINING

A SYSTEMATIC AND COMPREHENSIVE EXPOSITION OF THE  
THEORY, AND THE MORE IMPORTANT PRACTICAL  
PROBLEMS; WITH SOLAR, LUNAR, AND  
OTHER ASTRONOMICAL TABLES.

*Designed for Use as a Text-Book in Colleges and Academies.*

BY WILLIAM A. NORTON,

LATE PROFESSOR OF NATURAL PHILOSOPHY AND ASTRONOMY  
IN THE UNIVERSITY OF THE CITY OF NEW YORK.

PUBLISHED BY

WILEY & PUTNAM, New York; THOMAS, COWPERTHWAIT  
& CO., PHILADELPHIA; C. C. LITTLE & CO., BOSTON.

---

1839.

Entered according to an Act of Congress, in the year 1839, by  
WILLIAM A. NORTON,  
in the Clerk's Office of the District Court of the Southern District of New York.

## PREFACE.

---

THE object in writing the present treatise, has been to provide a suitable text book for the use of the students of Colleges and the higher Academies, and at the same time to furnish the practical astronomer with rules or formulæ, and accurate tables for performing the more important astronomical calculations.

The work is divided into four Parts. The first three Parts contain the theory: the First Part treating of the determination of the places and motions of the heavenly bodies; the Second, of the phenomena resulting from the motions of these bodies, and of their appearances, dimensions, and physical constitution; and the Third, of the theory of Universal Gravitation. The Fourth Part consists of practical problems, which are solved with the aid of the tables appended to the work. An Appendix is added, containing a large collection of useful trigonometrical formulæ, and such investigations of astronomical formulæ as, from their length, could not, consistently with the plan of the work, be admitted into the text, and which it was still deemed advisable to retain for the benefit of the few who might wish to pursue them.

The chief peculiarities of the present treatise are,

1. The adoption of the Copernican System as an hypothesis at the outset, leaving it to be established by the agreement between the conclusions to which it leads and the results of observation.
2. A connected exposition of the principles and methods of astronomical observation, embracing the doctrine of the sphere, the construction and use of the principal astronomical instruments, and the theory of the corrections for refraction, parallax, aberration, precession, and nutation.
3. The exhibition of the methods of determining the motions and places of the different classes of the heavenly bodies in one connection.
4. The explanation of the principles of the construction of astronomical tables.
5. The addition of a chapter on the measurement of time, embracing the explanation of the different kinds of time, the processes by which one is converted into another, the methods of determining the time from astronomical observations with the transit instrument and sextant, and the calendar.
6. The contemplation of the phenomena of the aspect and apparent motions of the heavenly bodies as *consequences* of their motions in space, and the deduction of the various circumstances of these phenomena from the theory of the orbital motions previously established.
7. A comprehensive view of the theory of Universal Gravitation, followed out into its various consequences.
8. An exposition of the operations of the disturbing forces in producing the perturbations of the motions of the Solar System.
9. The solution of practical problems by means of loga-

rithmic formulæ, instead of rules. 10. The addition of lunar, solar, and other astronomical tables of peculiar accuracy and improved arrangement.

It may further be mentioned, that many of the investigations have been materially simplified, and that the aim has been to introduce into all of them as much simplicity and uniformity of method as possible. Particular attention has also been paid to the diagrams, it being of great importance that they should convey correct notions to the mind of the student.

The problems in the Fourth Part are principally for making calculations relative to the Sun, Moon, and Fixed Stars. The tables of the Sun and Moon, used in finding the places of these bodies, have, for the most part, been abridged and computed from the tables of Delambre, as corrected by Bessel, and those of Burckhardt; and the tables of epochs have all been reduced to the meridian of Greenwich. These tables will give the places and motions of the Sun and Moon within a fraction of a second of the tables from which they were derived. But as this degree of accuracy will not generally be required, rules are also given in the Fourth Part for obtaining approximate results. The entire set of tables has been stereotyped, and great pains has been taken, by repeated revisions and verifications, to render them accurate.

The principal astronomical works which have been consulted in writing the present treatise, are those of *Vince, Gregory, Woodhouse, Delambre, Biot, Laplace, Herschel*, and *Gummere*; also *Francœur's Uranogra-*



*phy*, *Francœur's Practical Astronomy*, *Encyclopædia Metropolitana*, Article *Astronomy*, and *Bailey's Tables and Formulæ*. Free use has been made of the methods of investigation and demonstration pursued in these treatises, such modifications being introduced, in those which have been adopted, as the plan of the work required.

A list of the Errata that have been detected, which, it is believed, contains all that are important, may be found immediately after the Table of Contents. It is recommended to the student to make some note of these at the places where they occur in the text, before taking up the subject in course.

WILLIAM A. NORTON.

# TABLE OF CONTENTS.

---

## P A R T I.

### ON THE DETERMINATION OF THE PLACES AND MOTIONS OF THE HEAVENLY BODIES.

#### CHAPTER I.

	Page
Introductory Remarks—General Phenomena of the Heavens . . . . .	1

#### CHAPTER II.

On the Celestial and Terrestrial Spheres . . . . .	8
--	---

#### CHAPTER III.

On the Construction and Use of the Principal Astronomical Instruments . . . . .	19
<i>Transit Instrument</i> . . . . .	22
<i>Astronomical Clock</i> . . . . .	27
<i>Astronomical Quadrant</i> . . . . .	28
<i>Altitude and Azimuth Instrument</i> . . . . .	31
<i>Equatorial</i> . . . . .	ib.
<i>Sextant</i> . . . . .	32
<i>Micrometer—Errors of Instrumental Admeasurement</i> . . . . .	34

#### CHAPTER IV.

Theory of Corrections—Refraction—Parallax—Aberration—Precession—Nutation . . . . .	34
<i>Refraction</i> . . . . .	35
<i>Parallax</i> . . . . .	41
<i>Aberration</i> . . . . .	47
<i>Precession and Nutation</i> . . . . .	52
<i>Remarks on the Corrections.—Verification of the Hypothesis that the Diurnal Motion of the Stars is Uniform and Circular</i> . . . . .	59

## CHAPTER V.

	Page
Of the Earth ;—its Figure and Dimensions ;—Latitude and Longitude of a Place . . . . .	61
<i>Determination of the Latitude and Longitude of a Place . . . . .</i>	64

## CHAPTER VI.

Of the Places of the Fixed Stars . . . . .	67
--	----

## CHAPTER VII.

Of the Apparent Motion of the Sun in the Heavens . . . . .	72
--	----

## CHAPTER VIII.

Of the Motions of the Sun, Moon, and Planets, in their Orbits . . . . .	77
<i>Kepler's Laws . . . . .</i>	ib.
<i>Definitions of Terms . . . . .</i>	81
<i>Elements of the Orbit of a Planet . . . . .</i>	82
<i>Methods of Determining the Elements of the Sun's Apparent Orbit, or of the Earth's Real Orbit . . . . .</i>	83
<i>Methods of Determining the Elements of the Moon's Orbit . . . . .</i>	88
<i>Methods of Determining the Elements of a Planet's Orbit . . . . .</i>	91
<i>Mean Elements and their Variations . . . . .</i>	98

## CHAPTER IX.

On the Determination of the Place of a Planet, or of the Sun, or Moon, for a Given Time, by the Elliptical Theory ; and of the Verification of Kepler's Laws . . . . .	100
<i>Place of a Planet, or of the Sun or Moon in its Orbit . . . . .</i>	ib.
<i>Heliocentric Place of a Planet . . . . .</i>	102
<i>Geocentric Place of a Planet . . . . .</i>	ib.
<i>Places of the Sun and Moon . . . . .</i>	104
<i>Verification of Kepler's Laws . . . . .</i>	105

## CHAPTER X.

On the Inequalities of the Motions of the Planets and of the Moon ; and of the Construction of Tables for finding the Places of these Bodies . . . . .	106
<i>Construction of Tables . . . . .</i>	112

## CHAPTER XI.

Of the Motions of the Comets . . . . .	117
--	-----

## CHAPTER XII.

Of the Motions of the Satellites . . . . .	121
--	-----

## CHAPTER XIII.

	Page
On the Measurement of Time . . . . .	124
<i>Different Kinds of Time</i> . . . . .	ib.
<i>Conversion of one Species of Time into another</i> . . . . .	126
<i>Determination of the Time and Regulation of Clocks by Astronomical</i>	
<i>Observations</i> . . . . .	128
<i>Of the Calendar</i> . . . . .	132

---

## PART II.

ON THE PHENOMENA RESULTING FROM THE MOTIONS  
OF THE HEAVENLY BODIES, AND ON THEIR APPEAR-  
ANCES, DIMENSIONS, AND PHYSICAL CONSTITUTION.

## CHAPTER XIV.

	Page
Of the Sun and the Phenomena attending its Apparent Motions . . . . .	137
<i>Inequality of Days</i> . . . . .	ib.
<i>Twilight</i> . . . . .	141
<i>The Seasons</i> . . . . .	142
<i>Appearance, Dimensions, and Physical Constitution of the Sun</i> . . . . .	145

## CHAPTER XV.

Of the Moon and its Phenomena . . . . .	149
<i>Phases of the Moon</i> . . . . .	ib.
<i>Moon's Rising, Setting, and Passage over the Meridian</i> . . . . .	152
<i>Rotation and Librations of the Moon</i> . . . . .	155
<i>Dimensions and Physical Constitution of the Moon</i> . . . . .	156

## CHAPTER XVI.

Eclipses of the Sun and Moon—Occultations of the Fixed Stars . . . . .	158
<i>Eclipses of the Moon</i> . . . . .	ib.
<i>Eclipses of the Sun</i> . . . . .	168
<i>Occultations</i> . . . . .	186

## CHAPTER XVII.

Of the Planets and the Phenomena occasioned by their Motions in Space . . . . .	187
<i>Apparent Motions of the Planets with respect to the Sun</i> . . . . .	ib.

	Page
<i>Stations and Retrogradations of the Planets</i> . . . . .	191
<i>Phases of the Inferior Planets</i> . . . . .	193
<i>Transits of the Inferior Planets</i> . . . . .	194
<i>Appearances, Dimensions, Rotation, and Physical Constitution of the Planets</i> . . . . .	195

## CHAPTER XVIII.

Of Comets.—Their Appearance, Magnitude, and Physical Constitution . . . . .	202
---	-----

## CHAPTER XIX.

Of the Fixed Stars.—Their Number and Distribution over the Heavens— Annual Parallax and Distance—Variable Stars—Double Stars—Clusters of Stars, and Nebulæ . . . . .	204
<i>Annual Parallax and Distance of the Stars</i> . . . . .	205
<i>Variable Stars</i> . . . . .	206
<i>Double Stars</i> . . . . .	208
<i>Clusters of Stars—Nebulæ</i> . . . . .	209

## PART III.

## OF THE THEORY OF UNIVERSAL GRAVITATION.

## CHAPTER XX.

Of the Principle of Universal Gravitation . . . . .	Page 211
---	-------------

## CHAPTER XXI.

Theory of the Elliptic Motion of the Planets . . . . .	213
--	-----

## CHAPTER XXII.

Theory of the Perturbations of the Elliptic Motion of the Planets and of the Moon . . . . .	218
--	-----

## CHAPTER XXIII.

Of the Relative Masses and Densities of the Sun, Moon, and Planets; and of the Relative Intensity of the Gravity at their surface . . . . .	235
--	-----

# TABLE OF CONTENTS.

xi

## CHAPTER XXIV.

	Page
Of the Figure and Rotation of the Earth ; and of the Precession of the Equinoxes and Nutation . . . . .	236

## CHAPTER XXV.

Of the Tides . . . . .	240
------------------------	-----

# PART IV.

## ASTRONOMICAL PROBLEMS.

	Page
EXPLANATION OF THE TABLES . . . . .	247
PROB. I. To work, by logistical logarithms, a proportion the terms of which are degrees and minutes, or minutes and seconds of an arc ; or hours and minutes, or minutes and seconds of time . . . . .	254
PROB. II. To take from a table the quantity corresponding to a given value of the argument, or to given values of the arguments of the table . . . . .	255
PROB. III. To convert Degrees, Minutes, and Seconds of the Equator into Time . . . . .	262
PROB. IV. To convert Time into Degrees, Minutes, and Seconds . . . . .	262
PROB. V. The Longitudes of two Places, and the Time at one of them being given, to find the corresponding time at the other . . . . .	262
PROB. VI. The Apparent Time being given, to find the corresponding Mean Time ; or the Mean Time being given, to find the Apparent . . . . .	264
PROB. VII. To correct the Observed Altitude of a Heavenly Body for Refraction . . . . .	267
PROB. VIII. The Apparent Altitude of a Heavenly Body being given, to find its True Altitude . . . . .	269
PROB. IX. To find the Sun's Longitude, Semi-diameter, and Hourly Motion, for a given Time, from the Tables . . . . .	271
PROB. X. To find the Apparent Obliquity of the Ecliptic, for a given Time, from the Tables . . . . .	274
PROB. XI. Given the Sun's Longitude and the Obliquity of the Ecliptic, to find his Right Ascension and Declination . . . . .	275
PROB. XII. Given the Sun's Right Ascension and the Obliquity of the Ecliptic, to find his Longitude and Declination . . . . .	276

	Page
PROB. XIII. The Sun's Longitude and the Obliquity of the Ecliptic being given, to find the Angle of Position . . . . .	277
PROB. XIV. To find from the Tables, the Moon's Longitude, Latitude, Equatorial Parallax, Semi-diameter, and Hourly Motions in Longitude and Latitude, for a given Time . . . . .	277
PROB. XV. The Moon's Equatorial Parallax, and the Latitude of a Place being given, to find the Reduced Parallax and Latitude . . . . .	288
PROB. XVI. To find the Longitude and Altitude of the Nonagesimal Degree of the Ecliptic, for a given Time and Place . . . . .	289
PROB. XVII. To find the Apparent Longitude and Latitude, as affected by Parallax, and the Augmented Semi-diameter of the Moon ; the Moon's True Longitude, Latitude, Horizontal Semi diameter, and Equatorial Parallax, and the Longitude and Altitude of the Nonagesimal Degree of the Ecliptic, being given . . . . .	292
PROB. XVIII. To find the Mean Right Ascension and Declination, or Longitude and Latitude of a Star, for a given Time, from the Tables . . . . .	297
PROB. XIX. To find the Aberration of a Star in Right Ascension and Declination, for a given Day . . . . .	299
PROB. XX. To find the Nutation of a Star in Right Ascension and Declination, for a given Day . . . . .	300
PROB. XXI. To find the Apparent Right Ascension and Declination of a Star for a given Day . . . . .	301
PROB. XXII. To find the Aberration of a Star in Longitude and Latitude, for a given Day . . . . .	302
PROB. XXIII. To find the Apparent Longitude and Latitude of a Star, for a given Day . . . . .	303
PROB. XXIV. To compute the Longitude and Latitude of a Heavenly Body from its Right Ascension and Declination, the Obliquity of the Ecliptic being given . . . . .	304
PROB. XXV. To compute the Right Ascension and Declination of a Heavenly Body from its Longitude and Latitude, the Obliquity of the Ecliptic being given . . . . .	305
PROB. XXVI. The Longitude and Declination of a Body being given, and also the Obliquity of the Ecliptic, to find the Angle of Position . . . . .	307
PROB. XXVII. To find from the Tables the Time of New or Full Moon, for a given Year and Month . . . . .	308
PROB. XXVIII. To determine the number of Eclipses of the Sun and Moon that may be expected to occur in any given Year, and the Times nearly at which they will take place . . . . .	311
PROB. XXIX. To calculate an Eclipse of the Moon . . . . .	315
PROB. XXX. To calculate an Eclipse of the Sun, for a given Place . . . . .	320



## APPENDIX.

	Page
TRIGONOMETRICAL FORMULÆ . . . . .	341
I. <i>Relative to a Single arc or angle a</i> . . . . .	ib.
II. <i>Relative to Two Arcs a and b, of which a is supposed to be the greater</i> . . . . .	342
III. <i>Trigonometrical Series</i> . . . . .	343
IV. <i>Differences of Trigonometrical Lines</i> . . . . .	ib.
V. <i>Resolution of Right Angled Spherical Triangles</i> . . . . .	344
VI. <i>Resolution of Oblique Angled Spherical Triangles</i> . . . . .	346
INVESTIGATION OF ASTRONOMICAL FORMULÆ . . . . .	349
<i>Formulæ for the Parallax in Right Ascension and Declination, and in Longitude and Latitude</i> . . . . .	ib.
<i>Formulæ for the Aberration in Longitude and Latitude, and in Right Ascension and Declination</i> . . . . .	357
<i>Formulæ for the Nutation in Right Ascension and Declination</i> . . . . .	362
<i>Solution of Kepler's Problem, by which a Body's Place is found in an Elliptical Orbit</i> . . . . .	367
NOTE TO PROBLEM XIV . . . . .	371
<i>Rules for finding the Moon's Longitude, Latitude, Hourly Motions, Equatorial Parallax, and Semi-diameter, for a given time, from the Nautical Almanac</i> . . . . .	ib.



## E R R A T A.

Page 31, line 12. *For S Q, read T Q.*

" 44. *For  $\text{tang } \frac{1}{2} (z + p)$ , read  $\text{tang } (\frac{1}{2} z + p)$ .*

" ib. *For  $z$ , read  $\frac{1}{2} z$ .*

" ib. *For  $o s'$ , read  $o s$ .*

" 58, " 14. *For  $s a'$ , read  $\sin s a'$ .*

" 63, " 4. *For Art. 132, read Art. 142.*

" 69, " 22. *For Table LXI, read Table XC.*

" 73, " 10. *For declination, read right ascension.*

" 84, " 31. *For Art. 179, read Art. 178.*

" ib., " 36. *For Art. 174, read Art. 184.*

" 87, " 28. *For motion of the apogee and perigee in longitude, read sidereal motion of apogee and perigee.*

" 103. *For  $\frac{1 - \theta}{1 + \theta}$ , read  $\frac{1 - \text{tang } \theta}{1 + \text{tang } \theta}$ .*

" 105, " 13. *For 246, read 247.*

" 116, " 6. *For Art. 265, read Art. 267.*

" 119, " 20. *For 1836, read 1826.*

" 131, " 4. *For 4 20, read 3 50. This will change slightly the result of the Problem.*

" 145. *For  $\text{tang } \frac{1}{2} A E B$ , read  $\sin \frac{1}{2} A E B$ .*

" ib. *For  $\text{tang } \frac{1}{2} \delta$ , read  $\sin \frac{1}{2} \delta$ .*

" 155, " 3. *For  $m - S$ , read  $m - s$ .*

" 156, " last. *For  $\frac{61' 24''}{33' 31''}$ , read  $\frac{33' 31''}{122' 48''}$ .*

" 159. *For  $\text{tang } E C a$ , read  $\sin E C a$ .*

" ib. *For  $\text{tang } (\delta - p)$ , read  $\sin (\delta - p)$ .*

" ib., " 26. *For tangent, read sine.*

" 160, " 25. *For maximum, read minimum, and for minimum, read maximum.*

- Page 165, line 13. *For C' f, read M f, and for C' f', read M f'.*  
 " 167. *For C R, read C R'.*  
 " 178, " 16. *For Prob. XXVII, read Prob. XVII.*  
 " 183, equ. 121. *For 3600<sup>s</sup>, read — 3600<sup>s</sup>.*  
 " ib., last equ. *For 1800<sup>s</sup>, read — 1800<sup>s</sup>.*  
 " 185, line 17. *For M m, read M n.*  
 " 200, " 25. *For 1848, read 1847.*  
 " 205, " 33. *For diametrically to, read diametrically opposite to.*  
 " 215, " 6. *For conclave, read concave.*  
 " 220, last equ. *For cos  $\gamma$ , read cos  $\beta$ .*  
 " 225, line 20. *For apogee to the perigee, read perigee to the apogee.*  
 " 228, " 19. *For (equa. 137), read (equa. 139).*  
 " 229, " 14. *For Art. 563, read Art. 564.*  
 " 230, " 13 & 19. *For S, read N.*  
 " ib., " 16. *For the same of, read the same side of.*  
 " 238, " 6. *For  $\frac{1}{2} \frac{1}{2} \frac{1}{6}$ , read  $\frac{1}{2} \frac{1}{3} \frac{1}{6}$ .*  
 " 252, " 21. *For LVII, read LXVII.*  
 " 254, " 23. *For logistical of, read logistical logarithm of.*  
 " 265, " 7. *For XVII, read XVIII.*  
 " 271. *For last Problem, read Problem VI.*  
 " 279, " 5. *For XXVII, read LIV.*  
 " 281, " 6. *For — 2' 37".2, read 237".2.*  
 " 293, " 1. *For R', read R.*  
 " 309, " 17. *For approximate of, read approximate time of.*  
 " 319. *For  $\lambda$ , read  $\lambda'$ .*  
 " 348, case 2. *For A, c, b, read A, C, b, and for C, read c.*  
 " 350, 2d equ. *For s p s', read z p s'.*  
 " 354. *For tang (s — 90°), read tang  $\frac{1}{2}$  (s — 90°).*

# PART I.

## ON THE DETERMINATION OF THE PLACES AND MOTIONS OF THE HEAVENLY BODIES.

---

### CHAPTER I.

#### INTRODUCTORY REMARKS. — GENERAL PHENOMENA OF THE HEAVENS.

1. ASTRONOMY is a mixed mathematical science, which treats of the motions, positions, distances, appearances, magnitudes, and physical constitution of the heavenly bodies. That part of the science, which has for its object the determination of these several particulars from observation, is called *Plane Astronomy*; and that, in which the physical causes of the motions and constitution of the heavenly bodies are investigated, is denominated *Physical Astronomy*.

2. To be able to form correct notions of the phenomena of the heavens, it is necessary to know the form of the earth. We learn from the following circumstances, that the earth is a body of a globular form, insulated in space. 1st. When a vessel is receding from the land, an observer, stationed upon the coast, first loses sight of the hull, then of the lower parts of the sails, and lastly, of the top-sails. This is the case, whatever is the direction of the course of the vessel, and at whatever part of the earth it is observed. 2d. At sea, the visible horizon, or the line bounding the visible portion of the earth's surface, is every where a circle, of a greater or less extent, according to the altitude of the point of observation, and is, on all sides, equally depressed. 3d. Navigators have sailed around the earth, and,

by steering their course continually in one direction, arrived, at length, at the place from which they departed. These facts prove the surface of the sea to be convex, and the surface of the land conforms very nearly to that of the sea ; for, the elevations of the highest mountains above the level of the ocean bear an exceedingly small proportion to the dimensions of the whole earth.

3. If, on a clear night, we observe the heavens, we shall find that the stars, while they retain the same situations with respect to each other, undergo a continual change of position with respect to the earth. Some will be seen to ascend from a quarter called the *East*, being replaced by others that come into view, or *rise* ; others, to descend towards the opposite quarter, the *West*, and to go out of view, or *set*. And, if our observations be continued throughout the night, with the east on our left and the west on our right, the stars which rise in the east will be seen to move in parallel circles entirely across the visible heavens, and finally to set in the west. Each star will ascend in the heavens during the first half of its course, and descend during the remaining half. The greatest heights of the several stars will be different, but they will all be attained towards that part of the heavens which lies directly in front, called the *South*. If we now turn our backs to the south, and direct our attention to the opposite quarter, the *North*, new phenomena will present themselves. Some stars will appear, as before, ascending, reaching their greatest heights, and descending ; but other stars will be seen, farther to the north, that never set, and which appear to revolve in circles, from east to west, about a certain star, that seems to remain stationary. This seemingly stationary star is called the *Polar Star* ; and those stars that revolve about it, and never set, are called *Circumpolar Stars*. It should be remarked, however, that the polar star, when accurately observed by means of instruments, is found not to be strictly stationary, but to describe a small circle about a point at a little distance from it, as a fixed centre. This point is called the *North Pole*. It is, in reality, about the north pole, as thus defined, and not the polar star, that the apparent revolutions of the stars at the north are performed. At the corresponding hours of the following night, the aspect of the heavens will be the same, from which it

appears that the stars return to the same position once in about 24 hours. It would seem, then, that the stars all appear to move, from east to west, exactly as if attached to the concave surface of a hollow sphere, which rotates in this direction about an axis passing through the station of the observer and the north pole of the heavens, in a space of time nearly equal to 24 hours. This motion, common to all the heavenly bodies, is called their *Diurnal Motion*.

4. It is ascertained by certain accurate methods of observation and computation, hereafter to be exhibited, that the diurnal motion of the stars is strictly *uniform and circular*.

5. A circle cut out of the heavens by a plane passing through the axis of rotation, has a *north and south* direction; and a circle cut out by a plane perpendicular to the axis, has an *east and west* direction.

6. The greater number of the stars preserve constantly the same relative position with respect to each other; and they are therefore called *Fixed Stars*. There are, however, a few stars, called *Planets*, which are perpetually changing their places in the heavens. The number of the planets is *ten*. Each has a distinctive name, as follows: Mercury, Venus, Mars, Jupiter, Saturn, Uranus, Ceres, Pallas, Juno, and Vesta. Mercury, Venus, Mars, Jupiter and Saturn, are visible to the naked eye, and have been known from the most ancient times. The other five, namely, Uranus, Ceres, Pallas, Juno, and Vesta, cannot be seen without the assistance of the telescope, and were discovered by modern observers.\*

7. The planets are distinguishable from each other, either by a difference of aspect, or by a difference of apparent motion with respect to the sun. Venus and Jupiter are the two most brilliant planets: they are quite similar in appearance, but their apparent motions, with respect to the sun, are very different. Venus never recedes beyond  $40^{\circ}$  or  $50^{\circ}$  from the sun, while Jupiter is

---

\* The planet Uranus was discovered in 1781 by Dr. Herschel, who gave it the name of the *Georgium Sidus*. By the European astronomers it was called *Herschel*. It is now generally known by the name given in the text. Ceres, Pallas, Juno, and Vesta, have been discovered since 1800; the first by Piazzi, the second and fourth by Olbers, and the third by Harding.



seen at every variety of angular distance from him. Mars is known by the ruddy color of his light. Saturn has a pale, dull aspect.

8. The apparent motion of the planets is generally directed towards the east ; occasionally, however, they are seen moving towards the west. As their easterly prevails over their westerly motion, they all, in process of time, accomplish a revolution around the earth. The periods of revolution are different for each planet.

9. The Sun and Moon are also continually changing their places among the fixed stars.

10. From repeated examinations of the situation of the moon among the stars, it is found, that she has with respect to them a progressive circular motion, from *west* to *east*, and completes a revolution around the earth in about 27 days.

11. The motion of the sun is also constantly progressive, and directed from *west* to *east*. This will appear, on observing for a number of successive evenings, the stars which first become visible in that part of the heavens where the sun sets. It will be found, that those stars, which in the first instance were observed to set just after the sun, soon cease to be visible, and are replaced by others that were seen immediately to the east of them ; and that these, in their turn, give place to others situated still farther to the east. The sun, then, is continually approaching the stars that lie on the eastern side of him. The period of time, in which he accomplishes a revolution in the heavens, is about 365 days.

It is to be observed that the sun does not advance *directly* towards the east. He has also some motion from south to north, and north to south. It is a matter of common observation, that the sun is moving towards the north from winter to summer, and towards the south from summer to winter.

12. When the place of the sun in the heavens is accurately found from day to day by certain methods of observation, hereafter to be explained, it appears that his path is an exact circle, inclined about  $23^{\circ}$  to a circle running due east and west. (Art. 5.)

13. The motions of the sun, moon, and planets, are for the most part confined to a certain zone, of about  $18^{\circ}$  in breadth,

extending around the heavens from west to east, which has received the name of the *Zodiac*.

14. There is yet another class of bodies, called *Comets*, that have a motion among the fixed stars. They appear only occasionally in the heavens, and continue visible only for a few weeks, or months. They shine with a diffusive, nebulous light, and are commonly accompanied by a faint divergent stream of similar light, called a *tail*.

15. The motions of the comets are not restricted to the zodiac. These bodies are seen in all parts of the heavens, and moving in every variety of direction.

16. By inspecting the planets with telescopes, it has been discovered that some of them are constantly attended by a greater or less number of small stars, whose positions are continually varying. These attendant stars are called *Satellites*. The planets which have satellites are Jupiter, Saturn, and Uranus. The satellites are sometimes called *Secondary Planets*; the planets upon which they attend being denominated *Primary Planets*.

17. The sun and moon, the planets, (including the earth,) together with their satellites, and the comets, compose the *Solar System*.

18. From the consideration of the apparent motions and other phenomena of the Solar System, several theories have been formed in relation to the arrangement and actual motions in space of the bodies that compose it. The theory, or *system*, now universally received, is (in its most prominent features) that which was taught by Copernicus in the sixteenth century, and which is known by the name of the *Copernican System*. It is as follows:

19. The sun occupies a fixed centre, about which the planets (including the earth) revolve from west to east,\* in planes that are but slightly inclined to each other, and in the following order: Mercury, Venus, the Earth, Mars, Vesta, Juno, Ceres, Pallas, Jupiter, Saturn, and Uranus. The earth rotates from west to east about an axis, inclined to the plane of its orbit, under an

---

\* A motion in space from *West* to *East*, is a motion from *right* to *left*, to a person situated within the orbit described, and on the north side of its plane.

angle of about  $66\frac{1}{2}^{\circ}$ , and which remains continually parallel to itself as the earth revolves around the sun. The moon revolves from west to east around the earth as a centre ; and, in like manner, the satellites circulate from west to east around their primaries. Without the Solar System, and at immense distances from it, are the fixed stars. (See Plate 1, which is a diagram of the Solar System in projection.)

20. We shall here, at the outset, adopt this system as an *hypothesis*, and shall rely upon the simple and complete explanations it affords of the celestial phenomena, as they come to be investigated, together with the evidence furnished by Physical Astronomy, to produce entire conviction of its truth in the mind of the student.

21. The following are the characters or symbols employed by astronomers for denoting the several planets, and the sun and moon :—

The Sun, . . . . . ☉	Ceres, . . . . . ♀
Mercury, . . . . . ☿	Pallas, . . . . . ♀
Venus, . . . . . ♀	Jupiter, . . . . . ♃
The Earth, . . . . . ⊕	Saturn, . . . . . ♄
Mars, . . . . . ♂	Uranus, . . . . . ♅
Vesta, . . . . . ♁	The Moon, . . . . . ☾
Juno, . . . . . ♃	

22. The angular distance between any two fixed stars is found to be the same, from whatever point on the earth's surface it is measured. It follows, therefore, that the diameter of the earth is insensible, when compared with the distance of the fixed stars ; and that, with respect to the region of space which separates us from these bodies, the whole earth is a mere point. Moreover, the angular distance between any two fixed stars is the same, at whatever period of the year it is measured. Whence, if the earth revolves around the sun, its entire orbit must be insensible, in comparison with the distance of the stars.

23. On the hypothesis of the earth's rotation, the diurnal motion of the heavens is a mere illusion, occasioned by the rotation of the earth. To explain this, suppose the axis of the earth prolonged on till it intersects the heavens, considered as concentric with the earth. Conceive a great circle to be traced through the two points of intersection and the point directly

over head, and let the position of the stars be referred to this circle. It will be readily perceived, that the relative motion of this circle and the stars will be the same, whether the circle rotates with the earth from west to east, or, the earth being stationary, the whole heavens rotate about the same axis and at the same rate in the opposite direction. Now, as the motion of the earth is perfectly equable, we are insensible of it, and, therefore, attribute the changes in the situations of the stars, with respect to the earth, to an actual motion of these bodies. It follows, then, that we must conceive the heavens to rotate as above mentioned, since, as we have seen, such a motion would give rise to the same changes of situation as the supposed rotation of the earth. It was stated (Art. 3) that the sphere of the heavens appears to rotate about a line passing through the north pole and the station of the observer; but, as the radius of the earth is insensible in comparison with the distance of the stars, an axis passing through the centre of the earth will, in appearance, pass through the station of the observer, wherever it may be upon the earth's surface.

24. We in like manner infer, that the observed motion of the sun in the heavens is only an apparent motion, occasioned by the orbital motion of the earth. Let  $E E'$  (Fig. 1) represent two positions of the earth in its orbit  $E E' E''$  about the sun  $S$ . When the earth is at  $E$ , the observer will refer the sun to that part of the heavens marked  $s$ ; but when the earth is arrived at  $E'$ , he will refer it to the part marked  $s'$ ; and being in the mean time insensible of his own motion, the sun will appear to him to have described in the heavens the arc  $s s'$ , just the same as if it had actually passed over the arc  $S S'$  in space, and the earth had, during that time, remained quiescent at  $E$ . The motion of the sun from  $s$  towards  $s'$  will be from west to east, since the motion of the earth from  $E$  towards  $E'$  is in this direction. Moreover, as the axis of the earth is inclined to the plane of its orbit under an angle of  $66\frac{1}{2}^\circ$ , (Art. 19,) the plane of the sun's apparent path, which is the same as that of the earth's orbit, will be inclined  $23\frac{1}{2}^\circ$  to a circle perpendicular to the earth's axis, or to a circle directed due east and west.

## CHAPTER II.

## ON THE CELESTIAL AND TERRESTRIAL SPHERES.

25. IN determining from observation the apparent positions and motions of the heavenly bodies, and in general, in all investigations that have relation to their apparent positions and motions, Astronomers conceive all these bodies, whatever may be their actual distance from the earth, to be referred to a spherical surface of an indefinitely great radius, having the station of the observer, or what comes to the very same thing, the centre of the earth, for its centre. This imaginary spherical surface is called the *Sphere of the Heavens*, or, the *Celestial Sphere*. It is important to observe, that by reason of its great dimensions, if two lines be drawn through any two points of the earth, and parallel to each other, they will, when prolonged on, meet it sensibly in the same point; and that, if two parallel planes be passed through any two points of the earth, they will intersect it sensibly in the same great circle.

26. For the purposes of observation and computation, certain imaginary points, lines, and circles, appertaining to the celestial sphere, are employed, which we shall now proceed to explain.

1. The *Vertical Line*, at any place on the earth's surface, is the line of descent of a falling body, or the position assumed by a plumb line, when the plummet is freely suspended and at rest.

Every plane that passes through the vertical line is called a *Vertical Plane*. Every plane that is perpendicular to the vertical line, is called a *Horizontal Plane*.

2. The *Sensible Horizon* of a place on the earth's surface, is the circle in which a horizontal plane, drawn through the place, cuts the celestial sphere. As its plane is tangent to the earth, it separates the visible from the invisible portion of the heavens.

3. The *Rational Horizon* is a circle parallel to the former, the plane of which passes through the centre of the earth. The zone of the heavens comprehended between the sensible and



rational horizon is imperceptible, or the two circles appear as one and the same, at the distance of the earth.

4. The *Zenith* of a place is the point in which the vertical prolonged upwards pierces the celestial sphere. The point in which the vertical, when produced downwards, intersects the celestial sphere, is called the *Nadir*.

The zenith and nadir are the poles of the horizon.

5. The *Axis of the Heavens* is an imaginary right line, passing through the north pole (Art. 3) and the centre of the earth. It is the line about which the apparent rotation of the heavens is performed. It is, also, on the hypothesis of the earth's rotation, the axis of rotation of the earth prolonged on to the heavens.

6. The *South Pole* of the heavens is the point in which the axis of the heavens meets the southern part of the celestial sphere.

To illustrate the preceding definitions, let the circle N E S Q (Fig. 2.) represent the earth, and O Z the *vertical* of a point O on its surface. Then, H O R will be the plane of the *sensible horizon*, H' C R' the plane of the *rational horizon*, O Z the direction of the *zenith*, and O C that of the *nadir*. And, if O P represent the direction in which an observer at O will see the north pole, C P, parallel to O P, will be the *axis of the heavens*.

Now, neglecting the size of the earth, or conceiving the observer to be stationed at its centre, let C (Fig. 3) be the place of observation, C Z the vertical corresponding to O Z in Fig. 2, and P the north pole; then will Z be the *zenith* and N the *nadir*; the great circle H A R a, the poles of which are Z, N, the *horizon*; P C P' the axis of the heavens; and P' the *south pole*.

7. *Vertical Circles* are great circles passing through the zenith and nadir. They cut the horizon at right angles, and their planes are vertical. Thus, Z S M represents a vertical circle passing through the star S, called the *Vertical Circle of the Star*.

8. The *Meridian* of a place is that vertical circle which contains the north and south poles of the heavens. The plane of the meridian is called the *Meridian Plane*.

Thus, P Z R P' is the meridian of the station C. The half H Z R, above the horizon, is termed the *Superior Meridian*, and the other half R N H, below the horizon, is termed the *Inferior*

*Meridian.* The two points, as H and R, in which the meridian cuts the horizon, are called the *North* and *South Points* of the horizon; and the line of intersection, as H C R, of the meridian plane with the plane of the horizon, is called the *Meridian Line*, or the *North and South Line*.

9. The *Prime Vertical* is the vertical circle which crosses the meridian at right angles. It cuts the horizon in two points, as *e, w*, called the *East* and *West Points of the Horizon*.

10. The *Altitude* of any heavenly body is the arc of a vertical circle, intercepted between the centre of the body and the horizon, or the angle at the centre of the sphere, measured by this arc. Thus, S M is the altitude of the star S.

11. The *Zenith Distance* of a heavenly body is the arc of a vertical circle, intercepted between its centre and the zenith; or the distance of the centre of the body from the zenith, as measured by the arc of a great circle. Thus, Z S is the zenith distance of the star S.

It is obvious that the zenith distance and altitude of a body are *compliments* of each other, and, therefore, when either one is known, that the other may be found.

12. The *Azimuth* of a heavenly body is the arc of the horizon, intercepted between the meridian and the vertical circle, passing through the centre of the body; or the angle comprehended between the meridian plane and the vertical plane containing the centre of the body. It is reckoned either from the north or from the south point, and each way from the meridian. H M represents the azimuth of the star S.

*The Azimuth and Altitude, or azimuth and zenith distance of a heavenly body, ascertain its position with respect to the horizon and meridian, and, therefore, its place in the visible hemisphere.* Thus, the azimuth H M determines the position of the vertical circle Z S M of the star S, with respect to the meridian Z P H, and the altitude M S, or the zenith distance Z S, the position of the star in this circle.

13. The *Amplitude* of a heavenly body at its rising, is the arc of the horizon intercepted between the point where the body rises and the east point. Its amplitude at setting is the arc of the horizon, intercepted between the point where the body sets and the west point. It is reckoned towards the north, or towards



the south, according as the point of rising or setting is north or south of the east or west point. Thus, if  $aBSA$  represents the circle described by the star  $S$  in its diurnal motion,  $ea$  will be its amplitude at rising, and  $wa$  its amplitude at setting.

14. The *Celestial Equator*, or the *Equinoctial*, is a vertical of the celestial sphere, the plane of which is perpendicular to the axis of the heavens. The north and south poles of the heavens are therefore its geometrical poles. The celestial equator is represented in Fig. 3, by  $EwQe$ . This circle is also frequently called the *Equator*, simply.

15. *Parallels of Declination* are small circles, parallel to the celestial equator.  $aBSA$  represents the parallel of declination of the star  $S$ .

The parallels of declination passing through the stars, are the circles described by the stars, in their apparent diurnal motion. These, by way of abbreviation, we shall call *Diurnal Circles*.

16. *Celestial Meridians*, *Hour Circles*, and *Declination Circles*, are different names given to all great circles, which pass through the poles of the heavens, cutting the equator at right angles.  $PS P'$  is a celestial meridian. The angles comprehended between the hour circles and the meridian, reckoning from the meridian towards the west, are called *Hour Angles* or *Horary Angles*.

17. The *Ecliptic* is that great circle of the heavens which the sun appears to describe in the course of the year.

18. The *Obliquity of the Ecliptic* is the angle under which the ecliptic is inclined to the equator.

19. The *Equinoctial Points* are the two points in which the ecliptic intersects the equator. That one of these points, which the sun passes in the spring, is called the *Vernal Equinox*, and the other, which is passed in the autumn, is called the *Autumnal Equinox*. These terms are also applied to the *epochs* when the sun is at the one or the other of these points.

20. The *Solstitial Points* are the two points of the ecliptic  $90^\circ$  distant from the vernal and autumnal equinox. The one that lies to the north of the equator is called the *Summer Solstice*, and the other the *Winter Solstice*. The epochs of the

sun's arrival at these points are also designated by the same terms.

21. The *Equinoctial Colure* is the celestial meridian passing through the equinoctial points; and the *Solstitial Colure*, is the celestial meridian passing through the solstitial points.

22. The *Polar Circles* are parallels of declination at a distance from the poles equal to the obliquity of the ecliptic. The one about the north pole is called the *Arctic Circle*; the other, about the south pole, is called the *Antarctic Circle*.

The polar circles contain the poles of the ecliptic.

23. The *Tropics* are parallels of declination at a distance from the equator equal to the obliquity of the ecliptic. That which is on the north side of the equator, is called the *Tropic of Cancer*, and the other, the *Tropic of Capricorn*.

The tropics touch the ecliptic at the solstitial points.

Let C (Fig. 4) represent the centre of the earth and sphere, C P the axis of the heavens, E V Q A the equator, W V L A the ecliptic, and K, K', its poles. Then will V be the *vernal* and A the *autumnal equinox*; W the *winter*, and T the *summer solstice*; P V P' A the *equinoctial colure*; P K W K' T the *solstitial colure*; the angle T C Q, or its measure the arc T Q, the *obliquity of the ecliptic*; K m U, K' m' U', the *polar circles*; and T n Z, W' n' Z', the *tropics*.

24. The *Zodiac* (Art. 13) extends about  $9^\circ$  on each side of the ecliptic.

25. The ecliptic and zodiac are divided into twelve equal parts, called *Signs*. Each sign contains  $30^\circ$ . The division commences at the vernal equinox. Setting out from this point, and following around from west to east, the *Signs of the Zodiac*, with the respective characters by which they are designated, are as follows: Aries ♈, Taurus ♉, Gemini ♊, Cancer ♋, Leo ♌, Virgo ♍, Libra ♎, Scorpio ♏, Sagittarius ♐, Capricornus ♑, Aquarius ♒, Pisces ♓.

A motion in the heavens in the order of the signs, or from west to east, is called a *direct* motion, and a motion contrary to the order of the signs, or from east to west, is called a *retrograde* motion.

26. The *Right Ascension* of a heavenly body is the arc of the equator, intercepted between the vernal equinox and the declina-

tion circle which passes through the centre of the body, as reckoned from the vernal equinox towards the east. It measures the inclination of the declination circle of the body to the equinoctial colure. Thus, P S R being the declination circle of the star S, and V the place of the vernal equinox, V R is the right ascension of the star.

27. The *Declination* of a heavenly body is the arc of a circle of declination, intercepted between the centre of the body and the equator. It therefore expresses the distance of the body from the equator. Thus, R S is the declination of the star S.

Declination is *North*, or *South*, according as the body is north, or south of the equator.

*The right ascension and declination of a heavenly body are two co-ordinates, which, taken together, fix its position in the sphere of the heavens:* for, they make known its situation with respect to two circles, the equinoctial colure, and the equator. Thus, V R and R S ascertain the position of the star S with respect to the circles P V, and V Q A E.

28. The *Polar Distance* of a heavenly body is the arc of a declination circle, intercepted between the centre of the body and the elevated pole. The polar distance is the complement of the declination, and, therefore, when either is known, the other may be found.

29. *Circles of Latitude* are great circles of the celestial sphere, which pass through the poles of the ecliptic, and therefore cut this circle at right angles. Thus, K S L represents a part of the circle of latitude of the star S.

30. The *Longitude* of a heavenly body is the arc of the ecliptic, intercepted between the vernal equinox and the circle of latitude, which passes through the centre of the body, as reckoned from the vernal equinox towards the east, or in the order of the signs. It measures the inclination of the circle of latitude of the body to the circle of latitude passing through the vernal equinox. Thus, V L is the longitude of the star S.

31. The *Latitude* of a heavenly body is the arc of a circle of latitude, intercepted between the centre of the body and the ecliptic. It therefore expresses the distance of the body from the ecliptic. Thus, L S is the latitude of the star S.

Latitude is *North*, or *South*, according as the body is north, or south of the ecliptic.

*The longitude and latitude of a heavenly body are another set of co-ordinates, which serve to fix its position in the heavens.* They ascertain its situation with respect to the circle of latitude passing through the vernal equinox and the ecliptic. Thus, V L and L S fix the position of the star S, making known its situation with respect to the circles K V and V T A W.

32. The *Angle of Position* of a star is the angle included at the star between the circles of latitude and declination passing through it. P S K is the angle of position of the star S.

33. The *Astronomical Latitude*, or the *Latitude* of a place, is the arc of the meridian intercepted between the zenith and the celestial equator. It is *North*, or *South*, according as the zenith is north, or south of the equator. Z E (Fig. 5) represents the latitude of the station o.

27. The earth's surface, considered as spherical (which accurate admeasurement, upon principles that will be explained in the sequel, proves it to be, very nearly,) is called the *Terrestrial Sphere*. The following geometrical constructions appertain to the terrestrial sphere, as it is employed for the purposes of astronomy. It will be observed that they correspond to those of the celestial sphere above described, and are used for similar objects.

1. The *North* and *South Poles of the Earth* are the two points in which the axis of the heavens intersects the terrestrial sphere. They are also the extremities of the earth's axis of rotation.

2. The *Terrestrial Equator* is the great circle, in which a plane passing through the centre of the earth, and perpendicular to the axis of the heavens and earth, cuts the terrestrial sphere. The *terrestrial* and the *celestial equator*, then, lie in the same plane. The poles of the earth are the geometrical poles of the terrestrial equator. The two hemispheres into which the terrestrial equator divides the earth, are called, respectively, the *Northern Hemisphere* and the *Southern Hemisphere*.

3. *Terrestrial Meridians* are great circles of the terrestrial



sphere, passing through the north and south poles of the earth, and cutting the equator at right angles. Every plane that passes through the axis of the heavens, cuts the celestial sphere in a *celestial meridian*, and the terrestrial sphere in a *terrestrial meridian*.

Let  $PP'$  (Fig. 5,) represent the axis of the heavens,  $O$  the centre of the earth, and  $p$  and  $p'$  its poles. Then,  $erq$  will represent the *terrestrial equator* ( $ERQ$  representing the celestial equator,) and  $pep'$  and  $psp'$  *terrestrial meridians* ( $PEP'$  and  $PS P'$  representing celestial meridians.)

4. The *Reduced Latitude* of a place on the earth's surface is the arc of the terrestrial meridian, intercepted between the place and the equator, or the angle at the centre of the earth measured by this arc. Thus,  $oe$ , or the angle  $OOe$ , is the reduced latitude of the place  $o$ . Latitude is *North*, or *South*, according as the place is north, or south of the equator. The reduced latitude differs somewhat from the astronomical latitude, by reason of the slight deviation of the earth from a spherical form. Their difference is called the *Reduction of Latitude*.

5. *Parallels of Latitude* are small circles of the terrestrial sphere parallel to the equator. Every point of a parallel of latitude has the same latitude.

The parallels of latitude which correspond in situation with the polar circles and tropics in the heavens, have received the same appellations as these circles. (See Defs. 22, 23, p. 12.)

6 The *Longitude* of a place on the earth's surface, is the inclination of its meridian to that of some particular station, fixed upon as a circle to reckon from, and called the *First Meridian*. It is measured by the arc of the equator, intercepted between the first meridian and the meridian passing through the place, and is called *East*, or *West*, according as the latter meridian is to the east, or to the west of the first meridian. Thus, if  $pqp'$  be supposed to represent the first meridian, the angle  $spq$ , or the arc  $ql$ , will be the longitude of the place  $s$ .

Different nations have, for the most part, adopted different first meridians. The English use the meridian which passes through the observatory at Greenwich, near London; and the French, the meridian of the observatory of Paris. In the United States, as we have no public observatory, the longitude

is, for astronomical purposes, reckoned from the meridian of Greenwich, or Paris, (generally the former.)

*The longitude and latitude of a place designate its situation on the earth's surface.* They are precisely analogous to the right ascension and declination of a star in the heavens.

28. The diagram (see Fig. 3) which we made use of in Art. 26, in illustrating our description of the circles of the celestial sphere, represents the aspect of this sphere at a place at which the north pole of the heavens is somewhere between the zenith and horizon. Such is the position of the north pole at all places situated between the equator and the north pole of the earth. For, let O (Fig. 6) represent a place on the earth's surface, H O R the horizon, O Z the vertical, H Z R the meridian, and Z E the latitude. Q O E will then represent the equinoctial, and P, P', 90° distant from E and on the meridian, the poles. Now, we have,

$$H P = Z H - Z P = 90^\circ - Z P; \quad Z E = P E - Z P = 90^\circ - Z P. \quad \text{Whence } H P = Z E.$$

Thus, *the altitude of the pole is every-where equal to the latitude of the place.* It follows, therefore, that in proceeding from the equator to the north pole, the altitude of the north pole of the heavens will gradually increase from 0° to 90°.

If the spectator is in the southern hemisphere, the elevated pole, as it is always on the opposite side of the zenith, from the equator, will be the south pole. At corresponding situations of the spectator, it will obviously have the same altitude as the north pole in the northern hemisphere.

29. Let us now inquire into the principal circumstances of the diurnal motion of the stars, as it is seen by a spectator situated somewhere between the equator and the north pole. And, in the first place, it is a simple corollary from the proposition just established, that the parallel of declination to the north, whose *polar distance is equal to the latitude of the place*, will lie entirely above the horizon, and just touch it at the north point. This circle is called the *circle of perpetual apparition*; the line a H (Fig. 7) represents its projection on the meridian plane. The stars comprehended between it and the north pole will *never set*. As the depression of the south pole is equal to the altitude of the north pole, the parallel of declination o R,

at a distance from the south pole equal to the latitude of the place, will lie entirely below the horizon, and just touch it at the south point. The parallel thus situated, is called the *circle of perpetual occultation*. The stars comprehended between it and the south pole will *never rise*.

The celestial equator (which passes through the east and west points) will intersect the meridian at a point E, whose zenith distance, Z E, is equal to the latitude of the place (Def. 33, Art. 26,) and consequently, whose *altitude*, R E, is equal to the *co-latitude of the place*. Therefore, in the situation of the observer above supposed, the equator Q O E, passing to the south of the zenith, will, together with the diurnal circles  $n r$ ,  $s t$ , &c., which are all parallel to it, be *obliquely* inclined to the horizon, making with it an angle equal to the co-latitude of the place. As the centres  $c$ ,  $c'$ , &c., of the diurnal circles lie on the axis of the heavens, which is inclined to the horizon, all diurnal circles situated between the two circles of perpetual apparition and occultation,  $a H$  and  $o R$ , with the exception of the equator, will be divided unequally by the horizon. The greater parts of the circles  $n r$ ,  $n' r'$ , &c., to the north of the equator, will be above the horizon; and the greater parts of the circles  $s t$ ,  $s' t'$ , &c., to the south of the equator, will be below the horizon. Therefore, while the stars situated in the equator will remain an equal length of time above and below the horizon, those to the north of the equator will remain a longer time above the horizon than below it; and those to the south of the equator, on the contrary, a longer time below the horizon than above it. It is also obvious, from the manner in which the horizon cuts the different diurnal circles, that the disparity between the intervals of time that a star remains above and below the horizon, will be the greater the more distant it is from the equator. Again, the stars will all *culminate*, or attain to their greatest altitude, in the meridian: for, since the meridian crosses the diurnal circles at right angles, they will have the least zenith distance when in this circle. Moreover, as the meridian bisects the portions of the diurnal circles which lie above the horizon, the stars will all employ the same length of time in passing from the eastern horizon to the meridian, as in passing from the meridian to the western horizon. The circumpolar



stars will pass the meridian *twice* in 24 hours ; once *above*, and once *below* the pole. These meridian passages are called respectively *Upper* and *Lower Culminations*, or *Inferior* and *Superior Transits*.

It is evident from what is stated in Art. 28, that the circumstances of the diurnal motion will be the same at any place in the southern hemisphere, as at the place which has the same latitude in the northern.

The celestial sphere in the position relative to the horizon which we have now been considering, which obtains at all places situated between the equator and either pole, is called an *Oblique Sphere*, because all bodies rise and set obliquely to the horizon.

30. When the spectator is situated on the equator, both the celestial poles will be in his horizon, (Art. 28,) and, therefore, the celestial equator and the diurnal circles in general will be perpendicular to the horizon. This situation of the sphere is called a *Right Sphere*, for the reason that all bodies rise and set at right angles with the horizon. It is represented in Fig. 8. As the diurnal circles are bisected by the horizon, the stars will all remain the same length of time above, and below the horizon.

31. If the observer be at either of the poles, the elevated pole of the heavens will be in his zenith, (Art. 28,) and, consequently, the celestial equator will be in his horizon. The stars will move in circles parallel to the horizon, and the whole hemisphere, on the side of the elevated pole, will be continually visible, while the other hemisphere will be continually invisible. This is called a *Parallel Sphere*. It is represented in Fig. 9.

## CHAPTER III.

ON THE CONSTRUCTION AND USE OF THE PRINCIPAL  
ASTRONOMICAL INSTRUMENTS.

32. Astronomical Instruments are, for the most part, used for the admeasurement of arcs of the celestial sphere, or of angles corresponding to such arcs at the earth's surface. They consist, essentially, of a telescope turning upon a horizontal axis, and of a vertical graduated limb, (or, in some cases, of both a vertical and a horizontal graduated limb,) to indicate the angle passed over by the telescope. At the common focus of the object-glass and eye-glass of the telescope is a diaphragm, or circular plate, attached to which are two very fine wires, or threads, crossing each other at right angles in its centre. The place of this diaphragm may be altered by adjusting screws; it is by this means brought into such a position, that the cross of the wires will lie on the axis of the telescope, (that is, the line joining the centre of the object-glass and eye-glass.) The line joining the centre of the object-glass and the cross of the wires, is technically termed the *Line of Collimation*. Bringing the cross of the wires upon the axis of the telescope, is called *Adjusting the Line of Collimation*. A star is known to be on the line of collimation when it is bisected by the cross-wires.

The telescope either turns around the centre of the graduated limb, or, which is more common, the limb and telescope are firmly attached to each other, and turn together. In the first arrangement, a small steel plate firmly connected with the telescope, slides along the limb. Upon this plate a small mark is drawn, which is called the *Index*. The required angle is *read off*, by noting the angle upon the limb which is pointed out by the index; the zero on the limb being generally, in practice, the point from which the angle is reckoned. When the telescope and graduated limb are firmly connected, the limb slides past the index, which is now stationary. The limbs of even the largest instruments are not divided into smaller parts than about

5', but, by means of certain subsidiary contrivances, the angle may, with some instruments, be read off to within a fraction of a second.

33. The principal contrivances for increasing the accuracy of the reading off of angles, are the *Vernier*, the *Micrometer Screw*, and the *Microscope Micrometer*. The Vernier is only the index plate, so graduated that a certain number of its divisions occupy the same space as a number one less on the limb. Fig. 10 represents a vernier, and a portion of the limb of the instrument, 15 divisions on the vernier corresponding to 14 on the limb. If we suppose the smallest divisions of the limb to be 15', and call  $x$  the number of minutes in one division of the vernier, then,

$$15x = 14 \times 15', \text{ and } x = 14'.$$

Thus, the *difference* between a division on the vernier and one on the limb, will be 1'. Accordingly, if the index, which is the first mark on the vernier, should be little past the mark  $40^\circ$  on the limb, and the second mark of the vernier should coincide with the next point of division, marked  $40^\circ 15'$ , the angle would be  $40^\circ 1'$ . If the third mark on the vernier were coincident with the next division of the limb, the angle would be  $40^\circ 2'$ . If the fourth with the next division to this,  $40^\circ 3'$ ; and so on.

By making the divisions on the vernier more numerous, the angle can be read off with greater precision; but a better expedient is provided in the *Micrometer Screw*. This piece of mechanism is represented in Fig. 10. The part E can be fastened to the limb of the instrument by means of a screw. F G is a screw, with a milled head at F, working in a collar fixed in the under part of E, and in a nut fixed in the under part of the telescope T t. When the part E is fixed or *clamped*, and the screw is turned around by its milled head at F, it must communicate a direct motion to the nut, and, consequently, to the telescope and vernier in the direction of F G. Attached to the screw, or to the small cylinder on which it is formed, is an index D, moveable together with the screw, and on a thin graduated immoveable plate, the profile only of which is shown in the figure. Suppose now that the screw is of such fineness, that while, together with the index D, it makes a complete revolution, the vernier moves through an arc of 1'. Then,

if the plate be divided into 60 equal parts, a motion of the index over one of these parts would answer to a motion of  $1''$  on the limb. This being understood, to show the use of the micrometer screw, suppose that no two marks on the vernier and limb are coincident: bring the two nearest into coincidence by turning the screw, and the number of divisions passed over by the index D will be the seconds to be added to, or subtracted from, the angle read off with the vernier. In observing the coincidence of the divisions of the limb and vernier, the eye is assisted by a microscope.

34. The Microscope Micrometer is a compound microscope firmly fixed opposite to the limb, and furnished with cross-wires in the focus of the eye-glass, or conjugate focus of the object-glass, moveable by a fine threaded screw. The observer looks through it at the limb. By turning its screw, the cross-wire is brought into exact coincidence with the nearest of the divisions of the limb, and, as with the micrometer screw, the distance through which the screw is turned makes known the distance from this division of the fixed centre of the microscope, which corresponds to the index of a fixed vernier plate.

35. It is obvious, that, other things being the same, instruments are accurate in proportion to the power of the telescope, and the size of the limb. The large instruments now in use in astronomical observatories, are relied upon as furnishing angles to within  $1''$  of the truth.

36. *Time* is an essential element in astronomical observation. Three different kinds of time are employed by astronomers: *Sidereal*, *Apparent* or *True Solar*, and *Mean Solar Time*.

37. *Sidereal Time* is time as measured by the diurnal motion of the stars, or, more properly, of the vernal equinox. A *Sidereal Day* is the interval between two successive meridian transits of a star, or, (as it is now most generally considered,) the interval between two successive transits of the vernal equinox. It commences at the instant when the vernal equinox is on the superior meridian, and is divided into 24 *Sidereal Hours*.

38. *Apparent*, or *True Solar Time*, is deduced from observations upon the sun. An *Apparent Solar Day* is the interval



between two successive meridian passages of the sun's centre ; commencing when the sun is on the superior meridian. It appears from observation, that it is a little longer than a sidereal day, and that its length is variable during the year. It is divided into 24 *Apparent Solar Hours*.

39. *Mean Solar Time* is measured by the diurnal motion of an imaginary sun, called the *Mean Sun*, conceived to move uniformly from west to east in the equator, with the real sun's mean motion in the ecliptic, and to have at all times a right ascension equal to the sun's mean longitude. A *Mean Solar Day* commences when the mean sun is on the superior meridian, and is divided into 24 *Mean Solar Hours*.

Since the mean sun moves uniformly and directly towards the east, the length of the mean solar day must be invariable.

40. The *Astronomical Day* commences at noon, and is divided into 24 hours ; but the *Calendar Day* commences at midnight, and is divided into two portions of 12 hours each.

41. Astronomical observations are, for the most part, made in the plane of the meridian. But some of minor importance are made out of this plane. The chief instruments employed for meridian observations, are the *Astronomical Quadrant*, and the *Transit Instrument*, in connection with the *Astronomical Clock*. These are the capital instruments of an observatory, inasmuch as they serve (as will soon be explained) for the determination of the places of the heavenly bodies, which are the fundamental data of astronomical science. The principal instruments used for observation at different azimuths, are the *Altitude and Azimuth Instrument*, the *Equatorial*, and the *Sextant*.

#### *Transit Instrument.*

42. The *Transit Instrument* is a meridional instrument, employed in conjunction with a clock or chronometer for observing the passage of celestial objects across the meridian, either for the purpose of determining their difference of right ascension, or of obtaining the correct time. Fig. 11 represents a fixed transit instrument. A D is a telescope, fixed, as it is represented in the figure, to an horizontal axis formed of two cones. The two small ends of these cones are ground into two perfectly equal cylinders ; which cylindrical ends are called *Pivots*.

These pivots rest on two angular bearings, in form like the upper part of a Y, and denominated Y's. The Y's are placed in two dove-tailed brass grooves fastened in two stone pillars E and W, so erected as to be perfectly steady. One of the grooves is horizontal, the other vertical, so that, by means of screws, one end of the axis may be pushed a little forwards or backwards, and the other end may be either slightly depressed or elevated: which two small movements are necessary, as it will be soon explained, for two adjustments of the telescope.

Let E be called the eastern pillar, W the western. On the eastern end of the axis is fixed (so that it revolves with the axis) an index  $n$ , the upper part of which, when the telescope revolves, nearly slides along the graduated face of a circle, attached, as it is shown in the figure, to the eastern pillar. The use of this part of the apparatus is to adjust the telescope to the altitude or zenith distance of a star the transit of which is to be observed. Thus, suppose the index  $n$  to be at  $o$ , in the upper part of the circle, when the telescope is horizontal: then, by elevating the telescope, the index is moved downwards. Suppose the position to be that represented in the figure, then the number of degrees between  $o$  and the index is the altitude.

The wire plate placed in the focus of the transit telescope has attached to it five vertical wires, together with one horizontal wire. In order to be seen at night, these wires require to be illuminated by artificial light. Their illumination is effected by making one of the cones hollow, and admitting the light of a lamp placed in the pillar opposite the orifice; which light is directed to the wires by a reflector placed diagonally in the telescope. The reflector, having a large hole in its centre, does not interfere with the rays passing down the telescope from the object.

43. We will now explain the principal adjustments of the transit. Upon setting the instrument up, it should be so placed, that the telescope, when turned down to the horizon, should point north and south, as near as can possibly be ascertained. This being done, then—

1. *To adjust the line of collimation.*

This adjustment consists in bringing the central vertical wire, within the telescope, to intersect the optical axis, which is sup-

posed to be fixed by the maker of the instrument perpendicularly to the axis of rotation. There is no occasion with this instrument, to have the horizontal wire intersect the optical axis with exactness. Direct the telescope to some small, distant, well defined object, (the more distant the better,) and bisect it with the middle of the central vertical wire; then lift the telescope out of its angular bearings, or Y's, and replace it, with the axis reversed. Point the telescope again to the same object, and if it be still bisected, the collimation adjustment is correct; if not, move the wires one half the angle of deviation, by turning the small screws that hold the wire plate, near the eye end of the telescope, and the adjustment will be accomplished; but, as half the deviation may not be correctly estimated in moving the wires, it becomes necessary to verify the adjustment, by moving the telescope the other half, which is done by turning the screw that gives the small azimuth motion to the Y before spoken of, and consequently, to the pivot of the axis which it carries. Having thus again bisected the object, reverse the axis as before, and if half the error was correctly estimated, the object will be bisected upon the telescope being directed to it. If it should not be bisected, the operation of reversing and correcting half the error must be gone through again, and, until after successive approximations the object is found to be bisected in both positions of the axis; the adjustment will then be perfect.

It is desirable that the central wire should be truly vertical, as we should then have the power of observing the transit of a star on any part of it, as well as the centre. It may be ascertained whether it is so, by elevating and depressing the telescope, when directed to a distant object; if it is bisected by every part of the wire, the wire is vertical; if it is not bisected, the wire should be adjusted, by turning the inner tube carrying the wire plate until the above test of its verticality be obtained.

2. *To set the axis of rotation of the telescope horizontal.*

44. This adjustment may be effected by means of a spirit-level, attached to two upright arms bent at their upper extremities, by which it is hung on the two pivots of the axis. At one end of the level is a vertical adjusting screw, by which that end may be elevated or depressed. Put the telescope in its place, and



observe to which end of the level the bubble runs, which will always be the more elevated end ; bring it back to the middle by the Y screw for vertical motion, and take off the level, and hang it on again with the ends reversed. Then, if the bubble is again found in the middle, the level is already parallel to the axis, and the axis horizontal ; but if not, adjust one half the error by the adjusting screw of the level, and the other half by the Y screw ; and let the operation of reversing, and adjusting by halves, be repeated until the bubble will remain stationary in either position of the level, in which case, both the level and axis will be horizontal.

3. *To adjust the line of collimation to the plane of the meridian.*

45. We have said, that upon setting the instrument up, the telescope is to be brought into the meridian plane, as near as can be ascertained. One mode of establishing it, is to direct the telescope to the pole star, and by repeated observations find the position corresponding to its greatest or least altitude. At the present time, we may, instead, compute by means of existing tables founded on observation, the time of the meridian transit of the pole star, and at that computed time bisect the star by the middle vertical wire. Afterwards, the line of collimation may be placed still more exactly in the plane of the meridian, in the following manner : Note the times of two successive superior transits of the pole star across the central vertical wire, and the time of the intervening inferior transit. If the line of collimation were exactly in the plane of the meridian, as the diurnal circles are bisected by this plane, the interval between the superior and next inferior transit, would be precisely equal to the interval between the inferior and next superior transit. Accordingly, if these intervals are not in fact equal, find by repeated trials, the position of the telescope and vertical wire for which they are equal, and the line of collimation will then be in the plane of the meridian. The method of regulating the clock, required in making this adjustment, will be explained when we come to treat of the astronomical clock.

46. When the transit telescope has once been placed accurately in the meridian plane, in order to avoid the repetition of troublesome verifications of its position, a meridian mark should

be set up, and permanently established, at a distance from the instrument ; its place being determined by means of the middle or meridional wire. At Greenwich, two such marks, one to the north and another to the south, are used ; they are vertical stripes of white paint upon a black ground, on buildings about two miles distant from the observatory. The position of the telescope is verified, by sighting at the meridian mark, when it is once established.

47. The times of the transits of the heavenly bodies are ascertained as follows : In the case of a star, the moments of its crossing each of the five vertical wires are noted ; as the wires are equally distant from each other, the mean of these times (or their sum divided by 5) will be the time of the star's crossing the middle wire, or of its meridian transit. The utility of having five wires, instead of the central one only, will be readily understood, from the consideration that a mean result of several observations is deserving of more confidence than a single one ; since the chances are, that an error which may have been made at one wire, will be compensated by an opposite error at another. If the body observed has a disc of perceptible magnitude, as in the cases of the sun, moon, and planets, the times of the passage of both the western and eastern limb across each of the five wires are noted, and the mean of the whole taken, which will be the instant of the meridian transit of the centre of the body.

The time of the transit of a body, may, in this manner, be ascertained within a few tenths of a second.

48. In the interval between the transits of any two stars, the arc of the equator which expresses their difference of right ascension, will pass across the meridian, the rate of the motion being that of  $15^{\circ}$  to a sidereal hour : hence the difference of the times of transit of two stars, as observed with a sidereal clock, when converted into degrees, by allowing  $15^{\circ}$  to the hour, will be the difference between the right ascensions of the two stars. We may then, in this manner, by means of a transit instrument and sidereal clock, find the differences between the right ascension of any one star and the right ascensions of all the others. This being done, as soon as the position of the vernal equinox with respect to the same star becomes known, (and

we shall show how to find it in the sequel,) the *absolute* right ascensions of all the stars will also become known. In the actually existing state of astronomical science, the right ascensions of all the stars are more or less accurately known, and a right ascension sought is now obtained *directly*, by noting the time of the transit of the body, with a sidereal clock regulated so as to indicate 0h. 0m. 0s., when the vernal equinox is on the meridian, and converting it into degrees.

*Astronomical Clock.*

49. The astronomical clock is very similar to the common clock. It has a compensation pendulum; that is, a pendulum so constructed, that its length is unaffected by changes of temperature. The hours on the face are marked from 1 to 24.

50. Astronomers make use of sidereal time (as already stated) in determining the right ascensions of the heavenly bodies, but for all other purposes, they generally use mean solar time.

51. *To regulate a sidereal clock.* When a clock is used for determining differences of right ascension, (Art. 48,) it is adjusted to sidereal time, if it goes equally and marks out 24 hours in a sidereal day; it being altogether immaterial at what time it indicates 0h. 0m. 0s. To ascertain the *daily rate* of going of a clock which is to be adjusted to sidereal time for the purpose just mentioned, note by the clock the times of two successive meridian transits of the same star. The difference between the interval of the transits and 24 hours, will be the daily *gain* or *loss* (as the case may be) of the clock with respect to a perfectly accurate sidereal clock.\* If the gain or loss, when found after this manner, proves to be the same each day, then the mean rate of going is the same each day.

Next, to be able to discover the rate from hour to hour during the day, it is necessary to have obtained beforehand, at various times, and under various states of the circumstances likely to influence the rate of going of the clock, the differences between the times of the transits of a number of different stars, (correct-

---

\* It is not necessary, in order to obtain the daily rate of a sidereal clock, that the transit instrument should be adjusted to the plane of the meridian. It is only requisite that it should be kept fixed in some one vertical plane.

ing proportionally for the daily rate,) and to take the mean of the several differences found for each pair of stars for the exact difference of their transits. When this has been done, the rate of the clock may be found at all hours during the day, by noting by the clock *the differences between the times of the transits* of these stars, and comparing these with the exact differences already found. *At the present time*, the right ascensions of the stars being known, we have only, in order to ascertain the rate from hour to hour, to compare the intervals of time given by the clock between the transits of different stars taken in the order of their right ascension, with their *differences of right ascension*.

52. The sidereal clocks now in use, are made to indicate 0h. 0m. 0s. when the vernal equinox is on the superior meridian. For the regulation of such clocks, it is necessary to know not only their *rate*, but also their *error*. This is found by noting the time of the transit of a star, and comparing it with its right ascension expressed in time. If the two are equal, the clock is right, otherwise their difference will be its error.

If the error of the rate of a clock be considerable, it should be diminished by altering the length of the pendulum; otherwise, it may be allowed for. The stars best adapted to the regulation of clocks, are those in the vicinity of the equator; for, as their motion is more rapid than that of the stars more distant from the equator, there is less liability to error in noting their transits.

53. A mean solar clock is usually regulated by observations upon the sun. The method of regulating it cannot be adequately explained until we have treated of the apparent motions of the sun. It will here suffice to state, that with the instruments we have now described the sun's motions can be ascertained; and that, therefore, as a knowledge of these is all that is necessary, in order that we may be able to obtain the mean solar time at any instant, it is *possible* to express all intervals of time in *mean solar time*.

#### *Astronomical Quadrant.*

54. An Astronomical Quadrant is an instrument designed for the measurement of the meridian zenith distances or altitudes of the heavenly bodies, of which the essential parts are a grad-

uated quadrantal limb, a telescope turning upon an axis which passes through the centre of the limb, and a plumb line or level, to ascertain the vertical or horizontal line. (See Fig. 12.) Astronomical quadrants may be either portable or fixed. Portable quadrants are mounted upon an upright stem resting upon a tripod, and can be turned around in azimuth. Fixed quadrants have their axis firmly fastened in a wall, in a horizontal position, and perpendicularly to the plane of the meridian. They are hence called *Mural Quadrants*. The large mural quadrants of the Greenwich Observatory are of 8 feet radius.

55. The same adjustments are necessary for the quadrant as for the transit instrument; and in addition, the horizontal wire must be brought to intersect the axis, and the vertical, or horizontal point on the limb, must be found. The methods of effecting the adjustments are also the same with the fixed quadrant, except in the case of the collimation adjustment. This cannot be effected without the intervention of a portable quadrant or similar instrument. The horizontal wire of the telescope of a portable quadrant may be brought to intersect the optical axis, by directing the telescope to some star near the zenith, bisecting it with the horizontal wire, then turning the instrument  $180^\circ$  in azimuth, and moving the wire over half its angular distance from the star in the new position of the telescope, and repeating the process until the star is bisected in both positions of the instrument. This adjustment may be avoided altogether, by taking the half difference of the zenith distances of the star in the two positions of the instrument for the constant index error. The *horizontal point* (technically so called) is the place of the index answering to the horizontal position of the telescope. It may be found by means of a level or plumb line.

56. In place of mural quadrants, *Mural Circles* are often used. For greater accuracy, the angle is read off at six different points of the limb, which is an entire circle, by means of six stationary microscope micrometers, (Art. 34,) and the mean of the different readings taken for the angle required.

57. There is another modification of the quadrant, called the *Zenith Sector*, which is used to measure the meridian zenith distances of stars that cross the meridian within a few degrees



of the zenith. The limb extends only about  $10^\circ$  on each side of the lowest point. The zenith sector in the observatory at Greenwich has a radius of 12 feet.

58. The meridian altitude of a star is obtained by bringing the telescope into such a position, that the star will be bisected by the horizontal wire as it passes through the field of view, and observing the angle upon the limb. That of the sun, moon, or any planet, may be ascertained by measuring the altitudes of the upper and lower limbs, and taking their half sum for the altitude of the centre: or, if the apparent semidiameter be known, by adding this to the altitude of the lower limb, or subtracting it from the altitude of the upper limb.

59. The meridian altitude or zenith distance of a heavenly body having been measured with an astronomical quadrant, or other similar instrument, at a place the latitude of which is known, its declination may easily be found. For, let  $s, s', s''$  (Fig. 6,) represent the points of meridian passage of three different stars; one to the north of the zenith (Z,) one between the zenith and the equator (E,) and a third to the south of the equator, and we shall have—

$$\begin{aligned} E s = Z E + Z s, \quad E s' = Z E - Z s', \quad E s'' = Z s'' - Z E = \\ -(Z E - Z s''); \end{aligned}$$

or, in general,

Declination = latitude + zenith distance . . . (1);  
the latitude being taken always positive, the zenith distance being also taken positive when of the same name with the latitude, but negative when of a contrary name; and the declination being north, if it comes out positive, and south, if it comes out negative.

The latitude, which is here supposed to be known, may be found by measuring the meridian altitudes of a circumpolar star at its inferior and superior transits, and taking their half sum. For, as the pole lies midway between the points at which the transits take place, its altitude will be the arithmetical mean, or the half sum of the altitudes of these points, and the altitude of the pole is equal to the latitude of the place (Art. 28.)

60. When the right ascension and declination of a heavenly body have been obtained from observation, with a transit instrument and quadrant, (Arts. 48, 59,) its longitude and latitude may

be computed. For, Let S (Fig. 4) represent the place of the body, V R Q E the equator, V L T W the ecliptic, and P, K, the poles of the equator and ecliptic. In the triangle P K S we shall know, P S the compliment of S R the declination, and the angle K P S = E R - E V + V R =  $90^\circ$  + right ascension ; and, if we suppose the obliquity of the ecliptic to be known, we shall know P K. We may therefore compute K S, and the angle P K S. But K S is the compliment of S L, which is the latitude of the body S ; and P K S =  $180^\circ$  - E K S =  $180^\circ$  - (W V + V L) =  $180^\circ$  - ( $90^\circ$  + longitude) =  $90^\circ$  - longitude.

The obliquity of the ecliptic, which we have here supposed to be known, is, in practice, easily found ; for, it is equal to S Q the sun's greatest declination.

#### *Altitude and Azimuth Instrument.*

61. The Altitude and Azimuth Instrument consists, essentially, of a telescope with two graduated limbs, the one horizontal, and the other vertical. The telescope turns about the centre of the vertical limb, or turns with the limb about its centre ; and the vertical limb turns, with the telescope, about the vertical axis of the horizontal limb.

If the telescope be brought into the meridian plane, and afterwards directed upon a star out of this plane, the arc of the horizontal limb passed over by the index will be the *azimuth* of the star. The vertical limb will serve to measure its *altitude*.

62. The *Meridian Line* at a place may easily be determined with the altitude and azimuth instrument, by a method called the *Method of Equal Altitudes*. Let, O (Fig. 13) represent the place of observation, N P Z the meridian, and S, S' two positions of the same star, at which the altitude is the same. Now, the triangles Z P S and Z P S' have the side Z P common, Z S = Z S', and (allowing that the stars move in *circles*) P S = P S'. Hence, they are equal, and consequently the angle P Z S = P Z S' ; that is, *equal altitudes* of a star correspond to *equal azimuths*. Therefore, by bisecting the arc of the horizontal limb, comprehended between two positions of the vertical limb for which the observed altitude of a star is the same, we shall obtain the meridian line.

#### *Equatorial.*

63. The Equatorial is very similar, in its construction, to the



altitude and azimuth instrument. It is so called, from the circumstance of one of the limbs being placed in a position parallel to the plane of the equator. The axis of this limb is then parallel to the axis of the heavens; and the limb, to the centre of which the telescope is attached, is parallel in every one of its positions to the plane of some one celestial meridian. This instrument is particularly useful in the measurement of apparent diameters, and in all observations that require the telescope to be directed upon a body for a considerable period of time; as, by giving the limb to which the telescope is attached a slow motion from east to west, the body may be followed in its diurnal motion, and kept continually within the field of view.

#### *Sextant.*

64. The Sextant serves for the direct admeasurement of the angular distance between any two objects. Its essential parts are a graduated limb B C (Fig. 14), comprising about 60 degrees of the entire circle, which is attached to a triangular frame B A C; two mirrors, of which one (A), called the *Index Glass*, is moveable in connection with an index G about A, the centre of the limb, and the other (D), called the *Horizon-Glass*, is permanently fixed parallel to the radius A C drawn to the zero point of the limb, and is only half-silvered, (the upper half being transparent;) and an immoveable telescope at E, directed towards the horizon-glass. The principle of the construction and use of the sextant may be understood from what follows. A ray of light S A from a celestial object S, which impinges against the index-glass, is reflected off at an equal angle, and striking the horizon-glass (D), is again reflected to E, where the eye likewise receives through the transparent part of that glass, a direct ray from another point or object S'. Now, if A S' be drawn, directed to the object S', S A S', the angular distance between the two objects S and S' is equal to double the angle C A G, measured upon the limb of the instrument (A C being parallel to the horizon-glass.) For, when the index-glass is parallel to the horizon-glass and the angle on the limb is zero, A D, the course of the first reflected ray, will make equal angles with the two glasses, and therefore the angle S A D will become the angle S' A D (= A D E); and the observer, looking through the telescope, will see the same object S' both by direct and reflected

light. Now, if the index-glass be moved from this position through any angle  $C A G$ , the angle made by the reflected ray  $A D$  with this glass, will be diminished by an amount equal to this angle; for, we have  $D A G = D A C - C A G$ . Therefore, the angle made by the incident ray with the index-glass, as it is always equal to that made by the reflected ray, will be diminished by this amount. Consequently, the incident ray will, on the whole, that is, by the diminution of its inclination to the mirror by the angle  $C A G$ , and by the motion of the mirror through the same angle, be displaced towards the right an angle  $S A S'$ , equal to  $2 G A C$ . Thus, the angular distance  $S A S'$  of two objects  $S, S'$ , seen in contact, the one ( $S'$ ) directly, and the other ( $S$ ) by reflection from the two mirrors, is equal to twice the angle  $C A G$  that the index-glass is moved from the position ( $A C$ ) of parallelism to the horizon-glass.

Hence the limb is divided into 120 equal parts, which are called *degrees*; and to obtain the angular distance between two points, it is only necessary to sight directly at one of them, and then move the index until the reflected image of the other is brought into contact with it; the angle read off on the limb will be the angle sought.

65. To obtain the angular distance between two bodies which have a sensible diameter, bring the *nearest* limbs into contact, and to the angle read off on the limb *add* the sum of the apparent semi-diameters of the two bodies, or bring the *farthest* limbs into contact, and *subtract* this sum.

66. The sextant is also employed to take the altitude of a heavenly body. A horizontal reflector, called an *Artificial Horizon*, is placed in front of the observer: the angle between the body and its reflected image is then measured, as if this image were a real object; the half of which will be the altitude of the body.

A shallow vessel of mercury forms a very good artificial horizon.

67. In obtaining the altitude of a body, *at sea*, its altitude above the visible horizon is measured, by bringing the lower limb into contact with the horizon. To this angle is added the apparent semi-diameter of the body, and from the result is sub-

tracted the depression of the visible horizon below the horizontal line, called the *Dip of the Horizon*.

*Micrometer.—Errors of Instrumental Admeasurement.*

68. The *Apparent Diameter* of a heavenly body may be measured with great precision by means of a piece of apparatus attached to telescopes, called a *Micrometer*, which is designed for the admeasurement of small angles.

69. Whatever precautions may be taken, the results of instrumental admeasurement will never be wholly free from errors. Errors that arise from inaccuracy in the workmanship or adjustment of the instrument may be detected and allowed for. But, errors of *observation* are obviously undiscoverable. Since however the chances are, that an error committed at one observation will be compensated by an opposite error at another, it is to be expected that a more accurate result will be obtained, if a great number of observations, under varied circumstances, be made, instead of one, and the *mean* of the whole taken for the element sought. And accordingly, it is the uniform practice of astronomical observers to *multiply* observations as much as is practicable.

---

## CHAPTER IV.

THEORY OF CORRECTIONS.—REFRACTION.—PARALLAX.—  
ABERRATION.—PRECESSION.—NUTATION.

70. Angles measured at the earth's surface with astronomical instruments, answer to the *Apparent Place* of a heavenly body, and are termed *Apparent* elements. In astronomical language, the *True Place* of a heavenly body is its real place in the heavens, as it would be seen from the centre of the earth. Angles which relate to the true place, are denominated *True* elements. The apparent co-ordinates of a star are reduced to the true, by the application of certain corrections, called *Refraction*, *Parallax*, and *Aberration*.

71. Refraction, and aberration, are corrections for errors committed in the estimation of a star's place, while parallax serves to transfer the co-ordinates from the earth's surface to its centre. The object of the reduction of observations from the surface to the centre of the earth, is to render observations made at different places on the earth's surface directly comparable with each other. Observers occupying different stations upon the earth, refer the same body (unless it be a fixed star) to different points of the celestial sphere. Their observations cannot, therefore, be compared together, unless they be reduced to the same point, and the centre of the earth is the most convenient point of reference that can be chosen.

72. The co-ordinate planes or circles, to which the place of a star is referred (p. 13), are not strictly stationary, but, on the contrary, have a continual slow motion with respect to the stars. Hence, the true co-ordinates of a star's place, which have been found for any one epoch, will not answer, without correction, for any other epoch. The reduction from one epoch to another, is effected by applying two corrections, called *Precession* and *Nutation*.

### *Refraction.*

73. We learn from the principles of Pneumatics, as well as by experiments with the barometer, that the atmosphere gradually decreases in density from the earth's surface upwards. We learn also from the same sources, that it may be conceived to be made up of an infinite number of strata, of decreasing density, concentric with the earth's surface. From the known pressure and density of the atmosphere at the surface of the earth, it is computed, that by the laws of the equilibrium of fluids, if its density were throughout the same as immediately in contact with the earth, its altitude would be about 5 miles. Certain facts, hereafter to be mentioned, show that its actual altitude is not far from 50 miles. Now, it is an established principle of Optics, that light in passing from a vacuum into a transparent medium, or from a rarer into a denser medium, is bent, or *refracted*, towards the perpendicular to the surface at the point of incidence. It follows, therefore, that the light which comes from a star, in passing into the earth's atmosphere, or in passing from one stratum of atmosphere into another, is

refracted towards the radius drawn from the centre of the earth to the point of incidence.

74. Let  $M m n N$ ,  $N n o O$ ,  $O o q Q$  (Fig. 15) represent successive strata of the atmosphere. Any ray  $S p$  will then, instead of pursuing a straight course  $S p x$ , follow the broken line  $p a b c$ , being bent downwards at the points  $p$ ,  $a$ ,  $b$ ,  $c$ , &c., where it enters the different strata. But, since the number of strata is infinite, and the density increases by infinitely small degrees, the deflections  $a p x$ ,  $b a y$ , &c., as well as the lengths of the lines  $p a$ ,  $a b$ , &c., are infinitely small; and therefore,  $p a b c$  the path of the ray, is a broken line of an infinite number of parts, or a curved line concave towards the earth's surface, as it is represented in Fig. 16. Moreover, it lies in the vertical plane containing the original direction of the ray; for, this plane is perpendicular to all the strata of the atmosphere, and therefore the ray will continue in it in passing from one to the other.

75. The line  $O S'$  (Fig. 16), drawn tangent to  $p a b$  the curvilinear path of the light, at its lowest point, will represent the direction in which the light enters the eye, and therefore the apparent line of direction of the star. If, then,  $O S$  be the true direction of the star, the angle  $S O S'$  will be the displacement of the star produced by *Atmospherical Refraction*. This angle is called the *Astronomical Refraction*, or simply the *Refraction*.

Since  $p a b$  is concave towards the earth,  $O S'$  will lie above  $O S$ ; consequently, *refraction makes the apparent altitude of a star greater than its true altitude, and the apparent zenith distance of a star less than its true zenith distance.* (We here speak of the true altitude and true zenith distance, as estimated from the station of the observer upon the earth's surface.) Thus, to obtain the *true altitude* from the apparent, we must *subtract* the refraction; and to obtain the *true zenith distance* from the apparent, we must *add* the refraction. As refraction takes effect wholly in a vertical plane (Art. 74), it *does not alter the azimuth of a star.*

76. The amount of the refraction varies with the apparent zenith distance. In the zenith it is zero, since the light passes perpendicularly through all the strata of the atmosphere; and it is the greater, the greater is the zenith distance; for, the



greater the angle  $Z O a$  (Fig. 17), the greater will be the angle of refraction  $O a C$ , and consequently the greater the refraction.

*To find the amount of the refraction for a given zenith distance or altitude.*

77. Let us first show a method of resolving this problem by the general theory of refraction. According to this theory, the amount of the refraction, except so far as the convexity of the strata of the atmosphere may have an effect, depends wholly upon the absolute density of the air immediately in contact with the earth, and not at all upon the law of variation of the density of the different strata; that is, the actual refraction is the same that would take place, if the light passed from a vacuum immediately into a stratum of air of the density which obtains at the earth's surface. Let us suppose then, that the whole atmosphere is brought to the same density as that portion of it which is in contact with the earth, and let  $b a h$  (Fig. 17) represent its surface, also let  $O$  represent the station of the observer upon the earth's surface, and  $S a$  a ray incident upon the atmosphere at  $a$ . Denote the angle of refraction  $O a C$  by  $p$ , and the refraction  $O a x$  by  $r$ . The angle of incidence,

$$Z' a S = Z' a S' + S' a S = O a C + O a x = p + r.$$

Now if we represent the index of refraction of the atmosphere by  $m$ , we have by the laws of refraction,

$$\sin Z' a S = m \sin O a C, \text{ or } \sin (p + r) = m \sin p;$$

developing (App. For. 15,)

$$\sin p \cos r + \cos p \sin r = m \sin p;$$

or dividing by  $\sin p$ ,

$$\cos r + \cot p \sin r = m.$$

But as  $r$  is small, we may take  $\cos r = 1$ , and  $\sin r = r \sin 1''$ . (App. 47.)

Whence,  $1 + \cot p r \sin 1'' = m$ , or  $r = \frac{m-1}{\sin 1''} \times \frac{1}{\cot p} = A \text{ tang}$

$p$ ; putting  $A = \frac{m-1}{\sin 1''}$ . Let  $Z C a = C$ ; and  $Z O a = Z$ .  $O$

$a C = Z O a - Z C a$ , or,  $p = Z - C$ . Substituting, we have,

$$r = A \text{ tang } (Z - C) \dots (2).$$

When the zenith distance is not great,  $C$  is very small with respect to  $Z$ . If we neglect it, we have,

$$r = A \text{ tang } Z \dots (3);$$



which is the expression for the refraction, answering to the supposition that the surface of the earth is a plane, and that the light is transmitted through a stratum of uniformly dense air, parallel to its surface. We perceive, therefore, that the *refraction, except in the vicinity of the horizon, varies nearly as the tangent of the apparent zenith distance.*

78. It has been ascertained by experiment, that  $m$  the index of refraction, (the barometer being = 29.6 inches, and the thermometer =  $50^{\circ}$ ) = 1.0002803. Substituting in equation (3), after having restored the value of  $A$ , and reducing, there results,

$$r = 57''.8 \tan Z \dots \dots (4).$$

79. Dr. Bradley determined the constant  $A$  of formula (3) by means of astronomical observations, and made it  $57''.0$ , which gives the formula,

$$r = 57''.0 \tan Z \dots \dots (5).$$

80. Farther investigations conducted Dr. Bradley to a more accurate formula for the refraction, which is,

$$r = 57''.0 \tan (Z - 3r) \dots \dots (6) :$$

as in the preceding formula,  $57''$  is here the refraction at  $45^{\circ}$  apparent zenith distance. The value of  $r$  is obtained by successive approximations. In the first place,  $r$  is considered as zero, or  $3r$  is neglected in the second member; the resulting value found for the refraction, is then substituted in the place of  $r$  in the second member, and a second more exact value computed.

81. Other astronomers, by a combination of observation with theory, have obtained still more accurate formulæ. But the investigation, as well as the explanation of these, must be omitted in an elementary treatise like the present.

82. When the latitude or co-latitude of a place, and the polar distance of a star which passes the meridian near the zenith, have been determined, the refraction may be found for all altitudes from observation simply, without the aid of theory. For, let  $P$  (Fig. 18) be the elevated pole,  $Z$  the zenith,  $PZ$  the meridian,  $HOR$  the horizon,  $S$  the true place of a star, and  $S'$  its apparent place. Suppose the apparent zenith distance  $ZS'$  to have been measured. Now, in the triangle  $ZPS$ ,  $ZP$  the co-latitude, and  $PS$  the polar distance, are known by hypothesis, and the angle  $P$  is the sidereal time which has elapsed since the star's last meridian transit, (or, if the star be to the east of the

meridian, the difference between this interval and 24 sidereal hours,) converted into degrees by allowing  $15^\circ$  to the hour. Therefore we may compute the true zenith distance  $ZS$ , and subtracting from it the apparent zenith distance  $ZS'$ , we shall have the refraction. For the solution of this problem, the polar distance may be found by taking the complement of the declination computed from an observed meridian zenith distance (Art. 59); and, since the upper and lower transits of a circumpolar star take place at equal distances from the pole, the co-latitude may be found by taking the half sum of the greatest and least zenith distances of the pole star. But it is obvious, that neither of these quantities can be accurately determined, unless the measured zenith distances be corrected for refraction. When, however, the zenith distances in question differ considerably from  $90^\circ$ , the corresponding refractions may be at first ascertained with considerable accuracy by means of equation (4). When more correct formulæ have been obtained by this or any other process, the latitude and polar distance, and therefore the refraction answering to the measured zenith distance, will become more accurately known.

83. The various formulæ of refraction having been tested by numerous observations, it is found that they are all (though in different degrees) liable to material errors, when the zenith distance exceeds  $80^\circ$ , or thereabouts. At greater zenith distances than this, the refraction is *irregular*, or is frequently different in amount when the circumstances upon which it is supposed to depend are the same.

84. The refractive power of the air varies with its density, and hence the refraction must vary with the height of the barometer and thermometer.

85. The refractions which have place when the barometer stands at 29.6 inches (or, according to some astronomers, 30 inches), and the thermometer at  $50^\circ$ , are called *mean refractions*.

The refractions corresponding to any other height of the barometer or thermometer, are obtained by seeking the requisite *corrections* to be applied to the mean refractions, on the hypothesis that the refraction is directly proportional to the density of the atmosphere.

86. To save astronomical observers and computers the trouble of calculating the refraction whenever it is needed, the mean refractions corresponding to various zenith distances or altitudes are computed from the formulæ, as also the corrections for the barometer and thermometer, and inserted in a table. Table VIII is Dr. Young's table of mean refractions, and Table IX his table of corrections. The refraction answering to any zenith distance not inserted in the table can be found by simple proportion. (Sec Prob. VII)\*.

*Other effects of atmospherical refraction.*

87. Atmospherical refraction makes the apparent distance of any two heavenly bodies less than the true; for, it elevates them in vertical circles which continually approach each other from the horizon till they meet in the zenith.

88. Refraction also makes the discs of the sun and moon appear of an elliptical form when near the horizon. As it increases with an increase of zenith distance, the lower limb of the sun or moon is more refracted than the upper, and thus the vertical diameter is shortened, while the horizontal diameter remains the same, or very nearly so. This effect is most observable near the horizon, for the reason that the increase of the refraction is there the most rapid. The difference between the vertical and horizontal diameters may amount to 1-8 part of the whole diameter.

89. In the apparent horizon the refraction is about 34'. It follows, therefore, that when a star appears to be in the horizon, it is actually 34' below it. Refraction, then, retards the setting and accelerates the rising of the heavenly bodies.

Having this effect upon the rising and setting of the sun, it must increase the length of the day.

90. The apparent diameter of the sun is about 32'; as this is little less than the refraction in the horizon, it follows, that when the sun appears to touch the horizon, it is actually entirely below it. The same is true of the moon, as its apparent diameter is nearly the same with that of the sun.

---

\* The tables referred to in the text may be found near the end of the book. The problems referred to are in Part IV.

*Parallax.*

91. The correction for atmospherical refraction having been applied, the zenith distance of a body is reduced from the surface of the earth to its centre, by means of a correction called *Parallax*.

92. Parallax is, in its most general sense, the angle made by the lines of direction, or the arc of the celestial sphere comprised between the places of an object, as viewed from two different stations. It may also be defined to be the angle subtended at an object by a line joining two different places of observation. Let S (Fig. 19) represent a celestial object, and A, B two places from which it is viewed. At A it will be referred to the point  $s$  of the celestial sphere, and at B to the point  $s'$ ; the angle B S A, or the arc  $s s'$ , is the parallax. The arc  $s s'$  is taken as the measure of the angle B S A, on the principle that the celestial sphere is a sphere of an indefinitely great radius, so that the point S is not sensibly removed from its centre.

93. The term parallax is, however, generally used in Astronomy in a limited sense only, namely, to denote the angle included between the lines of direction of a heavenly body, as seen from a point on the earth's surface and from its centre; or the angle subtended at a heavenly body by a radius of the earth. If C (Fig. 20) is the centre of the earth, O a point on its surface, and S a heavenly body, O S C is the parallax of the body.

94. The parallax of a heavenly body above the horizon, is called *Parallax in Altitude*.

95. The parallax of a body at the time its apparent altitude is zero, or when it is in the plane of the horizon, is called the *Horizontal Parallax*. Thus, if the body S (Fig. 20) be supposed to cross the plane of the horizon at S', O S' C will be its horizontal parallax. O S C is a parallax in altitude of this body.

96. It is to be observed, that the definition just given of the horizontal parallax, answers to the supposition that the earth is of a spherical form. In point of fact, the earth (as will be shown in the sequel) is a spheroid, and accordingly the vertical and the radius at any point of its surface are inclined to each other, as represented in Fig. 21, where O C is the radius, and O C' the vertical. The points Z and  $z$  in which the vertical and radius pierce the celestial sphere are called, respectively, the *Apparent*

*Zenith*, and the *True Zenith*. In perfect strictness, the horizontal parallax is the parallax at the time  $z$  O S', the apparent distance from the true zenith, is  $90^\circ$ . No material error, however, will be committed, in supposing the earth to be spherical, except when the question relates to the parallax of the *moon*.

97. Let the apparent zenith distance Z O S = Z (Fig. 20) the true zenith distance Z C S =  $z$ , and the parallax O S C =  $p$ . Since the angle Z O S is the exterior angle of the triangle O S C, we have,

Z O S = Z C S + O S C, and hence also, Z C S = Z O S — O S C ;  
or,

$$Z = z + p, \text{ and } z = Z - p \dots (7).$$

Thus, to obtain the *true* zenith distance from the apparent, we have to *subtract* the parallax, and to obtain the *apparent* zenith distance from the true, to *add* the parallax.

Parallax, then, takes effect wholly in a vertical plane, like the refraction, but in the inverse manner; depressing the star, while the refraction elevates it. Thus, the refraction is added to Z, but the parallax is subtracted from it.

*To find an expression for the parallax in altitude.*

98. 1. *In terms of the apparent zenith distance.* In the triangle S O C (Fig. 20), the angle O S C = parallax in altitude =  $p$ , O C = radius of the earth = R, C S = distance of the body S = D, and C O S =  $180^\circ - \text{Z O S} = 180^\circ - \text{apparent zenith distance} = 180^\circ - Z$ ; and we have by Trigonometry, the proportion,

$$\sin \text{O S C} : \sin \text{C O S} :: \text{C O} : \text{C S};$$

whence,

$$\sin p : \sin (180^\circ - Z) :: R : D;$$

and

$$D \sin p = R \sin Z;$$

or,

$$\sin p = \frac{R}{D} \sin Z \dots (8).$$

This equation shows that the parallax  $p$  depends for any given zenith distance Z upon the distance of the body, and is less in proportion as this distance is greater: also, that for any given distance of the body it increases with an increase in the zenith



distance. When  $Z = 90^\circ$ ,  $p$  has its maximum value, and then = horizontal parallax =  $H$ ;

thus,

$$\sin H = \frac{R}{D} \dots \dots (9),$$

substituting,

$$\sin p = \sin H \sin Z \dots (10).$$

This last equation may be somewhat simplified. The distances of the heavenly bodies are so great, that  $p$  and  $H$  are always very small angles; even for the moon, which is much the nearest, the value of  $H$  does not at any time exceed  $62'$ . We may therefore, without material error, replace  $\sin p$  and  $\sin H$  by  $p$  and  $H$ . This being done, there results,

$$p = H \sin Z \dots (11).$$

Wherefore, *the parallax in altitude equals the product of the horizontal parallax by the sine of the apparent zenith distance.*

If we take notice of the deviation of the earth's form from that of a sphere,  $Z$ , in equation (10), will represent the apparent distance from the true zenith, (Art. 96,) and  $H$  the horizontal parallax as it is defined in Art. 96.

99. 2. *In terms of the true zenith distance.* In the actual state of Astronomy, the true co-ordinates of the places of the heavenly bodies are generally known, or may be obtained by computation, from the results of observations already made, and from these there is often occasion to deduce the apparent co-ordinates. For this purpose there is required an expression for the parallax in altitude in terms of the true zenith distance.

If we make  $Z = z + p$  (Art. 97) in equation (10), we shall have,

$$\sin p = \sin H \sin (z + p), \text{ or } \sin H = \frac{\sin p}{\sin (z + p)};$$

whence,

$$1 + \sin H = 1 + \frac{\sin p}{\sin (z + p)} = \frac{\sin (z + p) + \sin p}{\sin (z + p)};$$

and

$$1 - \sin H = 1 - \frac{\sin p}{\sin (z + p)} = \frac{\sin (z + p) - \sin p}{\sin (z + p)};$$

dividing,

$$\frac{1 + \sin H}{1 - \sin H} = \frac{\sin(z + p) + \sin p}{\sin(z + p) - \sin p};$$

or,

$$\text{tang}^2 (45^\circ + \frac{1}{2} H) = \frac{\text{tang} \frac{1}{2} (z + p)}{\text{tang} \frac{1}{2} z} \quad (\text{see App. For. 36, 29});$$

whence,

$$\text{tang} \frac{1}{2} (z + p) = \text{tang} \frac{1}{2} z \text{ tang}^2 (45^\circ + \frac{1}{2} H) \dots (12).$$

This equation makes known  $z + p$ , from which we may obtain  $p$  by subtracting  $z$ .

In order to be able to compute the parallax in altitude by means of formula (11) or (12), it is necessary to know  $H$  the horizontal parallax.

*To find the horizontal parallax.*

100. Let  $O, O'$  (Fig. 21) represent two stations upon the same terrestrial meridian  $O E O'$ , and at a remote distance from each other,  $Z, Z'$  their apparent zeniths, and  $z, z'$  their true zeniths,  $Q C E$  the equator, and  $S$  the body (supposed to be in the meridian,) the parallax of which is to be found. Let the angle  $O S O' = A$ ,  $z O S = Z$ ,  $z' O' S = Z'$ ; also let  $C O = R$ ,  $C O' = R'$ ,  $C S = D$ , the parallax in altitude  $O S C = p$ , and the parallax in altitude  $O' S C = p'$ . Now, by equation (8), replacing the sine of the parallax by the parallax itself, (Art. 98,)

$$p = \frac{R}{D} \sin Z, \text{ and } p' = \frac{R'}{D} \sin Z';$$

whence,

$$A = p + p' = \frac{R}{D} \sin Z + \frac{R'}{D} \sin Z' = \frac{R \sin Z + R' \sin Z'}{D};$$

but, (equa. 9)

$$H = \frac{R}{D}, \text{ or } D = \frac{R}{H}.$$

Substituting this value of  $D$ , and deducing the value of  $H$ , we have,

$$H = \frac{R \times A}{R \sin Z + R' \sin Z'} \dots (13).$$

It remains now to find an expression for  $A$  in terms of measurable quantities. Let  $O s$  and  $O' s'$  (Fig. 21,) be the directions at  $O$  and  $O'$  of a fixed star which crosses the meridian nearly at the same time with the body. Owing to the immense

distance of the star, these lines will be sensibly parallel to each other (Art. 22). Let the angle  $\text{S O } s$  the difference between the meridian zenith distances of the body and star, as observed at  $\text{O}$ , be represented by  $d$ , and let the same difference  $\text{S O' } s$  for the station  $\text{O'}$ , be represented by  $d'$ . Now,

$$\text{O S O'} = \text{O L O'} - \text{S O' } s = \text{S O } s - \text{S O' } s, \text{ or } A = d - d'.$$

If the body be seen on different sides of the star by the two observers, we shall have,

$$A = d + d'.$$

Substituting in equation (13), there results,

$$H = \frac{R (d \pm d')}{R \sin Z + R' \sin Z'} \dots (14).$$

If we regard the earth as a sphere,  $R = R'$ , and dividing by  $R$ , we have,

$$H = \frac{d \pm d'}{\sin Z + \sin Z'} \dots (15).$$

101. To find the parallax by means of these formulæ, each of the two observers must measure the meridian zenith distance of the body, and also of a star which crosses the meridian nearly at the same time with the body, and correct them for refraction. The difference of the two will be, respectively, the values of  $d$  and  $d'$ ; and the corrected zenith distances of the body will be the values of  $Z$  and  $Z'$ , if formula (15) be used; if formula (14) be used, the measured zenith distances of the body must still be corrected for the reduction of latitude, (p. 15, def. 4.)

It is not necessary that the two stations should be on precisely the same meridian; for, if the meridian zenith distance of the body be observed from day to day, its daily variation will become known; then, knowing also the difference of longitude of the two places, a simple proportion will give the change of zenith distance during the interval of time employed by the body in moving from the meridian of the most easterly to that of the most westerly station. This result, applied to the zenith distance observed at one of the stations, will reduce it to what it would have been, if the observation had been made in the same latitude on the meridian of the other station.

102. The Horizontal Parallax of a heavenly body may be found by the foregoing process, to within  $1''$  or  $2''$  of the truth. No greater degree of accuracy is necessary in the case of the

moon. But there are certain uses made of the horizontal parallax of a body, that will be noticed hereafter, which require that the parallax of the sun, and of the planets, should be known with much greater precision. The more accurate methods employed to determine the parallaxes of these bodies, will be explained (in principle at least) in subsequent parts of the work.

103. In consequence of the spheroidal form of the earth, the horizontal parallax of a body is somewhat different at different places. Let  $H$  and  $H'$  denote the horizontal parallaxes of the same body, and  $R$  and  $R'$  the radii of the earth at two different places. Then, by equation (9),

$$H = \frac{R}{D}, \text{ and } H' = \frac{R'}{D};$$

whence,

$$H : H' :: \frac{R}{D} : \frac{R'}{D} :: R : R'.$$

Thus the parallax at the equator, called the *Equatorial Parallax*, is the greatest, and the parallax at the pole the least. The difference between the parallaxes of the moon at the equator and at the pole may amount to about  $12''$ . For the other heavenly bodies the difference is too small to be taken into account.

104. When the horizontal parallax has been found for any one distance and time, from observation, the horizontal parallax for any other distance and time may be approximately computed, by means of the principle that the parallax of a body is directly proportional to its apparent diameter. The truth of this principle appears from the fact, that both the parallax (Art. 98) and the apparent diameter are inversely proportional to the same quantity, viz: the distance of the body from the centre of the earth.

In the present condition of astronomical science, when the horizontal parallax of either one of the heavenly bodies is required for any particular time, it may be obtained by computation, or from tables. It may also be taken out of the *Nautical Almanac*.\*

---

\* The *Nautical Almanac* is a collection of data to be used in nautical and astronomical calculations, published annually in England, and republished in New York. It may generally be obtained two or three years previous to the year for which it is calculated.

105. The equatorial horizontal parallax of the moon varies from  $53' 48''$  to  $61' 24''$ , according to the distance of the moon from the earth. The equatorial parallax of the moon answering to the mean distance, is  $57' 1''$ .

The horizontal parallax of the sun varies slightly from a change of distance. At the mean distance it is  $8''.6$ .

The horizontal parallaxes of the planets are comprised within the limits  $31''$ , and  $0''.4$ .

The fixed stars have no parallax.\*

*Parallax in right ascension and declination, and in longitude and latitude.*

106. Since parallax displaces a body in its vertical circle, which is generally oblique to the equator and ecliptic, it will alter its right ascension and declination, as well as its longitude and latitude. The difference between the true and apparent right ascension is called the *parallax in right ascension*; the like differences for the other co-ordinates are called, respectively, *parallax in declination*, *parallax in longitude*, and *parallax in latitude*. Formulæ by means of which these corrections may be found, when the right ascension and declination or the longitude and latitude are given, are investigated in the Appendix.

*Aberration.*

107. The celebrated English Astronomer, Dr. Bradley, commenced in the year 1725 a series of accurate observations upon the fixed stars, which were continued for several years, with the view of ascertaining whether the apparent places of the fixed stars were subject to any direct alteration, in consequence of the supposed change of the earth's position in space. The observations showed that there had been in reality, during the period of observation, small changes in the apparent places of each of the stars observed, which, when greatest, amounted to about  $40''$ ; but they were not such as should have resulted from the supposed changes of the earth's position in space. These phenomena Dr. Bradley undertook to examine and reduce to a general law. After repeated trials, he at last succeeded in discovering their true explanation. His theory is, that they are

---

\* The practical method of correcting for parallax is detailed and exemplified in Problem VIII.



different effects of one general cause, a progressive motion of light in conjunction with an orbital motion of the earth.

108. Let us conceive the observer to be stationed at the earth's centre ; and let  $A C B$  (Fig. 22) be a portion of the earth's orbit, so small that it may be considered a right-line,  $C S$  the absolute direction of a fixed star as seen from the point  $C$ ,  $A C$  the distance through which the earth moves in some small portion of time, and  $a C$  the distance through which a particle of light moves in the same time. Then, a particle of light, which, coming from the star in the direction  $S C$ , is at  $a$  at the same time that the earth is at  $A$ , will arrive at  $E$  at the same time that the earth does. Suppose, that when the earth is at  $A$ ,  $A a$  is the position of the axis of a telescope, and, that continuing parallel to itself, it takes up by virtue of the earth's motion, the successive positions  $A' a'$ ,  $A'' a''$  . . . . .  $C S'$  : a particle of light which follows the line  $S C$  in space will descend along this axis : for  $a a'$  is to  $A A'$  and  $a a''$  is to  $A A''$ , as  $a C$  is to  $A C$ , that is, as the velocity of light is to the velocity of the earth ; consequently, when the earth is at  $A'$ , the particle of light is on the axis at  $a'$ , and when the earth is at  $A''$  the particle of light is on the axis at  $a''$ , and so on for all the other positions of the axis, until the earth arrives at  $C$ . The apparent direction of the star  $S$ , as far, at least, as it depends upon the cause under consideration, will therefore be  $C S'$ .

The angle  $S C S'$ , which expresses the change in the apparent place of a star  $S$ , produced by the motion of light combined with the motion of the spectator, is called the *Aberration* of the star ; and the phenomenon of the change of the apparent course of the light coming from a star, thus produced, is called *Aberration of Light*, or simply *Aberration*.

109. If through the point  $a$  (Fig. 23) a line  $a s'$  be drawn parallel to  $A C$ , and terminating in  $C S'$ , the figure  $A a s' C$  will be a parallelogram, and therefore  $a s'$  will be equal to  $A C$ . Hence it appears, that if on  $C S$  the line of direction of a star  $S$ , a line  $C a$  be laid off, representing the velocity of light, and through  $a$  a line  $a s'$  be drawn, having the same direction as the earth's motion, and equal to its velocity, the line joining  $s'$  and  $C$  will be the apparent line of direction of the star, the point  $S'$  its apparent place in the heavens, and the angle  $a C s'$  its aber-

ration. We conclude, therefore, that by virtue of aberration, a star is seen in advance of its true place in the plane passing through the line of direction of the star and the line of the earth's motion.

110. The aberration is the same when a star is viewed with the naked eye, as when it is seen through a telescope. For, let  $a C$  the velocity of the light, be decomposed into two velocities, of which one  $A C$  is equal and parallel to the velocity of the earth; the other will be represented by  $s' C$ . Now, since the velocity  $A C$  is equal and parallel to the velocity of the earth, it will produce no change in the relative position of a particle of light and the eye, and therefore the relative motion of the light and the eye will be the same that it would be if the earth were stationary, and the light had only the velocity  $s' C$ ; accordingly, the light entering the eye just as it would do if it actually came in the direction  $s' C$ , and the eye were at rest,  $C s'$  will be the apparent direction of the star from which it proceeds.

111. If we regard the observer as situated upon the earth's surface, instead of being at its centre, the aberration resulting from the earth's motion of revolution will be still the same: for, all points of the earth advance at the same rate and in the same direction with the centre. The motion of rotation will produce an aberration proper to itself, but it is so small that there is no occasion to take it into account.

112. *To find a general expression for the aberration.* We have by Trigonometry (Fig. 23),

$\sin A a C : \sin C A a :: C A : C a :: \text{vel. of earth} : \text{vel. of light};$   
whence,

$$\sin A a C = \sin C A a \frac{C A}{C a}, \text{ or since } A a C = S C S',$$

$$\sin \text{aberr.} = \sin C A a \frac{\text{vel. of earth}}{\text{vel. of light}} \dots (15).$$

When  $C A a$  is  $90^\circ$ , the aberration has its maximum value, and this has been found by observation to be about  $20''$  ( $20''.36$ ),

whence, 
$$\sin 20'' = \frac{\text{vel. of earth}}{\text{vel. of light}} \dots (16).$$

substituting, and taking  $\sin B C a$  for  $\sin C A a$ , to which it is very nearly equal, we have,

$$\sin \text{aberr.} = \sin B C a \sin 20'' \dots (17).$$

We may conclude from this equation that the aberration increases with the angle  $B C a$  made by the direction of the star with the direction of the earth's motion ; that it is equal to zero, when this angle is zero, and has its maximum value of  $20''$  (more accurately  $20''.36$ ) when this angle is  $90^\circ$ .

113. Let us now inquire into the entire effect of aberration in the course of a year. Let  $S$  (Fig. 24) be the sun ;  $E$  the earth ;  $Efg$  its orbit ;  $ZTV$  that orbit extended to the fixed stars, or the ecliptic (p. 11, def. 17) ;  $ET$  a tangent to the earth's orbit at  $E$  ;  $\odot$  the place of  $S$  among the fixed stars or in the ecliptic, as seen from the earth ;  $s$  a fixed star ;  $sVT$  the arc of a great circle passing through  $s$  and  $T$ . Then, by what has preceded (Art. 109), the earth moving in the direction  $Efg$ , the apparent place of the star may be represented by  $s'$  and the aberration by  $sEs'$ . Thus, the effect of aberration at any one time is to displace the star by a small amount, directly towards the point  $T$  of the ecliptic, which is  $90^\circ$  behind the sun. As the earth moves, the position of the point  $T$  will vary ; and in the course of a year, while the earth describes its entire orbit in the direction  $Efg$ , this point will move in the same direction entirely around the ecliptic. In this period of time, therefore,  $ss'$  the small arc of aberration will revolve entirely around  $s$  the true position of the star ; from which we conclude, that in consequence of aberration a star appears to describe a closed curve in the heavens around its true place.

As the inclination of the direction of the star to the direction of the earth's motion will vary during a revolution of the earth, the aberration will also vary during this period (Art. 112), and hence the curve in question will not be a circle. It appears upon investigation that it is an ellipse, having the true place of the star for its centre, and of which the semi-major axis is constant and equal to  $20''.36$ , and the semi-minor axis variable and expressed by  $20''.36 \sin \lambda$  ( $\lambda$  denoting the latitude of the star). Each star, then, describes an ellipse which is the more eccentric in proportion as the star is the nearer to the ecliptic ; for, the expression for the minor axis shows, that the smaller the latitude the less will be this axis. For a star situated in the ecliptic, the minor axis will be zero, and the ellipse will be reduced to a right line. For a star in the pole of the ecliptic, the

minor axis is equal to the major, and the ellipse therefore becomes a circle.

114. Since aberration causes the apparent place of a star to differ slightly from its true place, the true and apparent co-ordinates will, in consequence, differ somewhat from each other. The effects of the aberration of light upon the apparent right ascension and declination of a star, are called, respectively, the *Aberration in Right Ascension*, and the *Aberration in Declination*. In like manner its effects upon the longitude and latitude are called the *Aberration in Longitude*, and the *Aberration in Latitude*. Formulæ, for computing these corrections to be applied to the apparent co-ordinates to obtain the true, or the reverse, are investigated in the Appendix.\*

115. Since the motion of the earth is at all times in a direction perpendicular, or nearly so, to the line followed by the light which comes from the sun to the earth, the aberration of the sun, which takes place only in longitude, is continually equal to  $20''.36$ , (Art. 112.) Thus, the sun's apparent place is always about  $20''.36$  behind its true place.

116. For a planet, the aberration is different from what it is for a fixed star. As a planet changes its place during the time that the light is passing from it to the earth, it would, if the earth were stationary, appear to be as far behind its true place as it has moved during this interval. This aberration due to the motion of the planet, combined with that due to the earth's motion, will give the real aberration of the planet.

117. For the moon, the aberration occasioned by its motion around the earth is very small. The earth's motion produces no lunar aberration, for the reason that the moon, and consequently the light emitted from it, partakes of this motion.

118. If the apparent places of a star, found at various times, be corrected for aberration, the same result for the true place of the star is obtained. Again, the deductions of Art. 113 agree in every particular with the observed phenomena of the apparent displacement of the stars first discovered by Dr. Bradley. These facts show that the aberration of light is the true cause

---

\* For the practical method of determining and applying these corrections, see Probs. XIX, XXI, XXII, XXIII.



of these phenomena, and consequently at the same time establish the fact of the earth's orbital motion, as well as that of the progressive motion of light.

119. It may be worth while to state, that the first discovery of the progressive motion of light preceded the detection and explanation by Bradley of the phenomena of aberration. The discovery was made by Roemer, a Danish astronomer, in the year 1667, from a comparison of observations upon the eclipses of Jupiter's satellites.

120. As to the actual velocity of light, we have by equation (16) vel. of earth : vel. of light ::  $\sin 20'' : 1 :: 1 : 10314$ . As determined from observations upon Jupiter's satellites, it is very nearly the same. The time employed by light in coming from the sun to the earth is 8m. 13s.

*Precession and Nutation.*

121. In the investigations that follow, we shall take it for granted that it is possible to find the obliquity of the ecliptic and the place of the equinox. Methods of determining them will be given, when we come to treat of the apparent motion of the sun.

122. By comparing the longitudes and latitudes of the same fixed stars, obtained at different periods (Art. 60), it is found that their latitudes continue very nearly the same, but that all their longitudes increase at the mean rate of about  $50''$  per year. The longitude of a star being the arc of the ecliptic, intercepted in the order of the signs between the vernal equinox and a circle of latitude passing through the star (p. 13, def. 30), it follows from the last mentioned circumstance, that the vernal equinox must have a motion along the ecliptic in a direction contrary to the order of the signs, amounting to about  $50''$  in a year. As it has been found that the autumnal equinox is always at the distance of  $180^\circ$  from the vernal, it must have the same motion. This retrograde motion of the equinoctial points, is called the *Precession of the Equinoxes*.

123. As the latitude of a star is its distance from the ecliptic (p. 13, def 31), it follows from the circumstance of the latitudes of all the stars continuing very nearly the same, that the ecliptic remains fixed, or very nearly so, with respect to the situations of the fixed stars.



124. The ecliptic being stationary, it is plain that the precession of the equinoxes must result from a continual slow motion of the equator in one direction. It appears from observation, that the obliquity of the ecliptic, or the inclination of the equator to the ecliptic, remains, in the course of this motion, very nearly the same.

125. Since the equator is in motion, its pole must change its place in the heavens. Let  $V L A$  (Fig. 25) represent the ecliptic,  $K$  its pole, which is stationary,  $P$  the position of the north pole of the equator or of the heavens at any given time, and  $V E A$  the corresponding position of the line of the equinoxes :  $K P L$  represents the circle of latitude passing through  $P$ , or the solstitial colure. Now, the point  $V$  being at the same time in the ecliptic and equator, is  $90^\circ$  distant from the two points  $K$  and  $P$  the poles of these circles ; therefore, it is the pole of the circle  $K P L$  passing through these points, and hence  $V L = 90^\circ$ . It follows from this, that when the vernal equinox has retrograded to any point  $V'$ , the pole of the equator, originally at  $P$ , will be found in the circle of latitude  $K P' L'$  for which  $V' L'$  equals  $90^\circ$  : it will also be at the distance  $K P'$  from the pole of the ecliptic equal to  $K P$ . Whence it appears, that the pole of the equator has a retrograde motion in a small circle about the pole of the ecliptic, and at a distance from it equal to the obliquity of the ecliptic. As the motion of the equator which produces the precession of the equinoxes, is uniform, the motion of the pole must be uniform also ; and as the pole will accomplish a revolution in the same time with the equinox, its rate of motion must be the same as that of the equinox, that is,  $50''$  of its circle in a year. The period of the revolution of the equinox and of the pole of the equator is 25920 years  $\left( = \frac{360^\circ}{50''} \right)$ .

126. The ecliptic, although very nearly stationary, as stated in Art. 123, is not strictly so. By comparing the values of the obliquity of the ecliptic, found at distant periods, it is ascertained that it is subject to a gradual diminution from century to century. A comparison of the results of observations made by Flamsteed in 1690, and by Dr. Maskelyne in 1769, gives for the mean secular diminution  $50''$ , and for the mean annual diminution

0".50. A more accurate determination of the mean annual diminution is 0".46.

It appears from observation, that there are minute secular changes in the latitudes of the stars, which establish that the diminution of the obliquity of the ecliptic arises from a slow displacement of the plane of the ecliptic (or of the earth's orbit) in space.

127. If the ecliptic slowly changes its position in the heavens, its pole must likewise ; and since the obliquity of the ecliptic is continually diminishing, its pole must be gradually approaching the pole of the equator.

128. The motion of the ecliptic alters somewhat the precession of the equinoxes, making it a little less than it would be, if the equator only was in motion : for, let  $E L$  (Fig. 26) represent the position of the ecliptic, and  $V Q$  that of the equator, at any assumed date, and  $E L'$ ,  $V' Q'$  the positions of the same circles at some later date ; the obliquity  $L' V'' Q'$  at the second epoch being less than that ( $L V Q$ ) at the first epoch : also let  $v$  be the physical point of the moveable ecliptic, which at the first epoch coincided with the point  $V$ . If the ecliptic remained stationary in the position  $E L$ , the precession during the interval of the epochs would be  $V V'$ . But, by reason of its motion, the actual precession is  $v V''$ , and it is obvious from an inspection of the figure, (the angle  $V V' Q'$  being an acute angle,) that this is less than  $V V'$  the precession on the fixed ecliptic. We learn by the aid of Physical Astronomy, that the amount of annual precession would, if the ecliptic were fixed, be 50".35. As we have already seen, the actual precession on the moveable ecliptic is 50" (more accurately, 50".23).

129. The motion of the equator which produces the precession of the equinoxes, must also produce changes in the right ascensions and declinations of the stars. These changes will be different, according to the situations of the stars with respect to the equator and equinoctial points.

130. It remains for us now to take notice of a minute *inequality* in the motion of the equator and its pole, which we have thus far overlooked. Dr. Bradley, in observing the polar distance of a certain star, with the view of verifying his theory of aberration, discovered that the observed polar distance did not

agree with the apparent polar distance, as computed from the results of previous observation, by allowing for precession, aberration, and refraction ; and hence inferred the existence of a new cause of variation in the co-ordinates of a star. On continuing his observations, he found that the polar distance alternately increased and diminished, and that it returned to the same value in about 19 years. These phenomena led him to suppose that the pole, instead of moving uniformly in a circle around the pole of the ecliptic, revolved around a point conceived to move in this manner.

If the pole has such a motion, it is plain that (allowing the fact of the earth's rotation) it must result from a vibratory motion of the earth's axis. To this supposed vibration of the axis of the earth, and consequently of that of the heavens, Dr. Bradley gave the name of *Nutation*. The term Nutation is also applied to the changes of the co-ordinates of a star's place, which are produced by the nutation of the earth's axis. The point about which the pole was conceived to revolve, is the mean position of the pole, or the *Mean Pole*.

Dr. Bradley discovered, from his observations, that the curve described by the pole must be an ellipse, having its major axis in the solstitial colure ; and estimated the value of the major axis at about  $18''$ , and that of the minor axis at about  $16''$ . He also discovered that a connection existed between the position of the pole in its ellipse, and the position of the moon at the time its latitude was zero (Art. 60), and changing from south to north, or of the point in which the moon crossed the plane of the ecliptic, in passing from the south to the north side of it, called the ascending node of the moon's orbit ; for, he found that the pole retrograded in like manner with the node ; that it completed its revolution in the same time, namely, in about 19 years ; and that its position was determinable from the place of the node by a geometrical construction. Let P (Fig. 27) represent the mean pole, and  $p$  the true pole ;  $pfg$  represents the ellipse described by the true pole around P as a centre ;  $g\ g'$ , lying in the solstitial colure K P L, being its major axis, and  $f\ f'$  its minor axis. It is to be observed, that the pole P is not stationary, but revolves in the circle N P P', carrying with it the ellipse  $pfg$ .

This theory of a nutation of the earth's axis has been completely verified by subsequent observations, and Physical Astronomy has revealed the cause of the phenomenon.

131. As the equator must move with the axis of the earth or heavens, nutation will change the position of the equinox and the obliquity of the ecliptic. It is plain that its effect upon the position of the equinox will be to make it oscillate periodically, and by equal degrees, from one side to the other of the position which corresponds to the mean pole; and that its effect upon the obliquity of the ecliptic, will be to make it alternately greater and less than the obliquity corresponding to the mean pole. The position of the equinox which corresponds to the mean pole, is called the *Mean Equinox*. The obliquity corresponding to the mean pole, is termed the *Mean Obliquity*. *Mean Equator* has a like signification. The real equinox and the real equator are called, respectively, the *True Equinox* and the *True Equator*. The actual obliquity of the ecliptic is termed the *Apparent Obliquity*. Right ascension and declination, as estimated from the true equator and true equinox, are called, respectively, *True Right Ascension*, and *True Declination*; and longitude, as reckoned from the true equinox, is called *True Longitude*. Right ascension, declination, and longitude, reckoned from the mean equinox and mean equator, are called, respectively, *Mean Right Ascension*, *Mean Declination*, and *Mean Longitude*. The true and mean co-ordinates differ by reason of nutation. The effect of nutation upon the right ascension is called the *Nutation in Right Ascension*; upon the declination, *Nutation in Declination*; and upon the longitude, *Nutation in Longitude*. Its effect upon the obliquity of the ecliptic is called *Nutation of Obliquity*. The distance of the true from the mean equinox in longitude, which is the same as the nutation in longitude, is sometimes termed the *Equation of the Equinoxes in Longitude*; and the distance, in right ascension, the *Equation of the Equinoxes in Right Ascension*. The precession of the mean equinox is equal to the *Mean Precession* of the true equinox, which is  $50''.2$ .

132. Formulæ, for computing the nutation in right ascension, declination, &c. at any given time, are investigated in the Appendix. These formulæ cannot be used without a knowledge of



the moon's motions. In practice, the nutations in right ascension, &c. are found by the aid of tables. (See Probs. XX, XXIII.) If these be applied to the true co-ordinates, the results will be the mean co-ordinates. If the mean co-ordinates be known, the same corrections will furnish the true.

133. Physical Astronomy has made known the existence of another nutation of the earth's axis, too small to be detected by observation. It is called *Solar Nutation*. The nutation discovered by Dr. Bradley is frequently called *Lunar Nutation*.

*To reduce the co-ordinates of a star from one epoch to another.*

134. This problem is resolved by first converting the true co-ordinates into the mean, then transferring the mean co-ordinates from the one epoch to the other, and finally converting the reduced mean co-ordinates into the true. The mode of performing the first and last mentioned operations has already been considered (Art. 132). It remains now for us to show how to *reduce mean co-ordinates from one epoch to another*.

135. 1. *When the interval of time between the epochs comprises but a few years.* In this case the changes from precession, of the mean right ascension and declination in the course of a year, called the *Annual Variation in right ascension*, and the *Annual Variation in declination*, are determined, then multiplied by the number of years in the interval, and applied as corrections to the given right ascension and declination.

For this purpose formulæ have been investigated, in which the annual variations in right ascension and declination are expressed in terms of the right ascension and declination of the star, and the obliquity of the ecliptic. Let  $V L A$  (Fig. 28) be the ecliptic,  $K$  its pole,  $P P' P''$  the circle described by the mean pole,  $P$  the mean pole and  $V Q A$  the mean equator at any given time,  $P'$  the mean pole and  $V' Q' A'$  the mean equator a year afterwards, and  $s$  a star. Draw  $P' r$  perpendicular to the declination circle  $P s a$ . We have,

$$\text{an. var. in dec.} = s a' - s a = P s - P' s = P r ;$$

but since  $P P' r$  may be considered as a right angled plane triangle,

$$P r = P P' \cos P' P r = P P' \sin Q P a . . . (18).$$

Regarding  $K P P'$  as a right angled isocles triangle, we obtain,

$$\sin K P P' \text{ or } 1 : \sin K P' :: \sin P K P' : \sin P P' ;$$



whence,

$$\sin P P' = \sin P K P' \sin K P', \text{ or } P P' = P K P' \sin K P' \\ (\text{nearly}) \dots (19);$$

substituting in equation (18), there results,

$$P r = P K P' \sin K P' \sin Q P a.$$

$P K P' = 50''.2$  (Art. 125);  $K P' = \text{obliquity of the ecliptic} = \omega$ ;  
 $Q P a = V Q - V a = 90^\circ - R$  ( $R$  designating the right ascension of the star  $s$ ). Thus, finally,

$$\text{an. var. in dec.} = 50''.2 \sin \omega \cos R \dots (20).$$

Next, we have,

$$\text{an. var. in r. asc.} = V' a' - V a = V' a' - m b = V' m + b a' \dots (21),$$

but,  $V' m = V V' \cos V V' m = 50''.2 \cos \omega$ ;

and since the right angled triangles  $s P' r$  and  $s b a'$  are similar,

$$\sin s r \text{ or } \sin s P' (\text{nearly}) : \sin P' r :: s a' : \sin b a';$$

whence,

$$\sin b a' = \sin P' r \frac{\sin s a'}{\sin P' s}, \text{ or } b a' = P' r \frac{\sin s a'}{\sin P' s} (\text{nearly}).$$

The triangle  $P P' r$  gives  $P' r = P P' \sin P' P r = P P' \cos Q P a = P K P' \sin K P' \cos Q P a$  (equa. 19); and  $\sin P' s = \cos s a'$ . Substituting, we obtain

$$b a' = P K P' \sin K P' \cos Q P a \frac{\sin s a'}{\cos s a'} = P K P' \sin K P' \cos \\ Q P a \text{ tang } s a'.$$

Replacing  $P K P'$ ,  $K P'$ , and  $Q P a$  by their values, as above, and taking the declination  $s a$  for  $s a'$  and denoting it by  $D$ , there results,

$$b a' = 50''.2 \sin \omega \sin R \text{ tang } D.$$

Now, substituting in equation (21) the values of  $V' m$  and  $b a'$ , we have,

$$\text{an. var. in r. asc.} = 50''.2 \cos \omega + 50''.2 \sin \omega \sin R \text{ tang } D \dots (22).$$

The results of formulæ (20, 22) are to be used with their algebraic signs, if the reduction is from an earlier to a later epoch, otherwise with the contrary signs. The declination is always to be considered *positive*, if *North*, and *negative*, if *South*.

$$V' m = 50''.2 \cos \omega = 50''.2 \cos 23^\circ 28' = 46''.0$$

is the annual retrograde motion of the equinoctial points along the equator.

— 136. 2. *When the interval of the epochs is of considerable or great length.* If the epochs are separated by an interval of more

than 10 or 12 years, the foregoing process will not answer: for in a period of ten years the annual variations will have sensibly altered.\* In this case we may proceed as follows: Convert the right ascension and declination into longitude and latitude, add to the longitude (or if the reduction be to an earlier epoch, subtract from it) the precession in longitude, which will be the product of  $50''.23$  by the interval of the epochs, expressed in years and parts of a year, and then with the longitude thus obtained, and the latitude, calculate the right ascension and declination, using the mean obliquity of the ecliptic.

When the period is of great length, or very great precision is desired, the precession on the fixed ecliptic should be used, which is  $50''.35$  per year (Art. 128); and the right ascension should be corrected for the change of the position of the equinox on the equator, produced by the motion of the ecliptic; which correction is —  $0''.13$  (per year) for later epochs.

*Remarks on the Corrections.—Verification of the Hypothesis that the Diurnal Motion of the Stars is Uniform and Circular.*

137. It appears from what we have stated on the subject of the Corrections: 1. That Refraction varies during the day with the altitude of the body, and changes for all altitudes with the state of the atmosphere; 2. That Parallax varies, like the Refraction, with the altitude of the body, and changes from one day to another with its distance; 3. That Aberration remains sensibly the same for two or three days, and depends for its absolute value on the time of the year; 4. That Precession and Nutation do not perceptibly alter the co-ordinates of a star, unless it be a circum-polar star, under several days, and that the former increases uniformly with the time, while the latter varies periodically, its effects entirely disappearing in about 19 years; and 5. That the absolute value of the Nutation depends entirely upon the longitude of the moon's ascending node.

138. In the determination of the amount and laws of the corrections, it was taken for granted by astronomers, that the diurnal

---

\* It is to be understood that we are here giving methods of obtaining very accurate results. The process just explained, except for stars near the pole, will furnish results sufficiently accurate for most purposes, even when the interval comprises 20 years or more.

motion of the stars ~~was~~ uniform and circular. This hypothesis may be verified in the following manner: Let the zenith distance and azimuth of the same star be measured at various times during a revolution, and corrected for refraction (the other corrections being insensible, Art. 137). Then, if the latitude of the place be known (Art. 59), in the triangle  $ZPS$  (Fig. 13) we shall have  $ZP$  the co-latitude,  $ZS$  the zenith distance of the star, and  $PZS$  its azimuth, whence we may compute  $PS$ . If this calculation be made for the time of each observation, it will be found that the same value for  $PS$  is obtained in every instance; which proves the diurnal motion to be circular. Again, let the angle  $ZPS$  be computed for the time of each observation, with the same data, and it will be found that it varies proportionally to the time; which establishes that the diurnal motion is also uniform, or, at least, sensibly so during one revolution.

139. When the transits of a circumpolar star are observed at intervals of several days, and allowance is made for the error of the rate of the clock, as determined from observations upon stars in the vicinity of the equator, and for the aberration in right ascension, it is found that the times of the transits differ slightly from each other; from which it appears, that the diurnal motion of the stars is not strictly uniform. When, however, allowance is made for the precession and nutation in right ascension, this difference disappears. We may hence conclude that the motion of rotation of the earth is uniform, and that the motions of the earth and of its axis, which produce the phenomena of precession and nutation, alter the times of the transits of the stars, thereby making the period of the apparent revolution of a star to differ slightly from the period of the earth's rotation.

It may be observed, that the greatest difference obtains in the case of the pole star, and is half a second.

## CHAPTER V.

## OF THE EARTH;—ITS FIGURE AND DIMENSIONS:—LATITUDE AND LONGITUDE OF A PLACE.

140. Although it is in general sufficient for astronomical purposes, to regard the earth as a sphere, still it is necessary in some cases of astronomical observation and computation, when accurate results are desired, to take notice of its deviation from the spherical form. No account need, however, be taken of the irregularities of its surface, occasioned by mountains and valleys, as they are exceedingly minute when compared with the whole extent of the earth. It is to be understood, then, that by the figure of the earth is meant the general form of its surface, supposing it to be smooth, or that the surface of the land corresponded with that of the sea.

141. The figure of the earth is ascertained from an examination of the form of the terrestrial meridians.

A *Degree* of a terrestrial meridian is an arc of it corresponding to an inclination of  $1^\circ$  of the verticals at the extremities of the arc. It is also called a *Degree of Latitude*.

142. The length of a degree at any place will serve as a measure of the curvature of the meridian at that place; for it is obvious from considerations already presented (Art. 2), that the earth, if not strictly spherical, must be nearly so, and therefore that a degree  $ab$  (Fig. 29) may, with but little, if any error, be considered as an arc of  $1^\circ$  of a circle, which has its centre at  $C$  the point of intersection of the verticals  $Ca$ ,  $Cb$  at the extremities of the arc. The curvature will then decrease in the same proportion as the radius of this circle increases, and therefore in the same proportion as the length of a degree increases. Wherefore, the form of a meridian may be determined by measuring the length of a degree at various latitudes.

143. *To determine the length of a degree of a terrestrial meridian.* To accomplish this, we have,

1. *To run a meridian line*; an operation which is performed in

the following manner: An altitude and azimuth instrument (or some other instrument adapted to meridian observations) is first placed at the point of departure, and accurately adjusted to the meridian. A new station is then established by sighting forward with the telescope. To this station the instrument is removed, and is there adjusted to the meridian by sighting back to the first station. A third station is then established by sighting forward with the telescope as before, to which the instrument is removed. By thus continually establishing new stations, and carrying the instrument forward, the meridian line may be marked out for any required distance. The meridian adjustments may be corrected from time to time by astronomical observations (Arts. 45, 62).

2. *To find the length of the arc passed over.* When the ground is level, the length of the arc may be directly measured. In case the nature of the ground is such as not to allow of a direct measurement, it may be calculated with equal precision, by means of a base line, and a chain of triangles the angles of which are measured.

3. *To find the inclination of the verticals at the extreme stations.* This angle may be obtained by measuring the meridian zenith distances of the same fixed star at the two stations, correcting them for refraction if they are observed about the same time; and for refraction, aberration, precession, and nutation, if they are observed at different times, and taking their difference. For, let O, O' (Fig. 29) be the two stations in question, Z, Z' their zeniths, and O S, O' S the directions of a fixed star, and we shall have,

$O c O' = Z O I - O I c = Z O S - Z' I S = Z O S - Z' O' S$ ;  
that is, the angle comprised between the verticals equal to the difference of the meridian zenith distances of the same star.

4. *The length of an arc of the meridian, either somewhat greater or less than a degree, having been found by the foregoing operations, thence to compute the length of a degree.* Let N denote the number of degrees and parts of a degree in the measured arc, A its length, and  $x$  the length of a degree. Then, allowing that the earth for an extent of several degrees does not differ sensibly from a sphere, we may state the proportion,

$$N : A :: 1^\circ : x, \text{ whence } x = \frac{1^\circ \times A}{N} \dots (23).$$

144. Degrees have been measured with the greatest possible



care, at various latitudes and on various meridians. Upon a comparison of the measured degrees, it appears that *the length of a degree increases as we proceed from the equator towards either pole*. It follows, therefore, (Art. 132,) that the curvature of a meridian is greatest at the equator, and diminishes as we go towards the poles; and, consequently, that *the earth is flattened at the poles*.

145. The fact of the decrease of the curvature of a terrestrial meridian from the equator to the poles, leads to the supposition that it is an ellipse, having its major axis in the plane of the equator, and its minor axis coincident with the axis of the earth. Analytical investigations, founded on the lengths of a degree in different latitudes and on different meridians, prove that a meridian is, in fact, very nearly an ellipse, and that the earth has very nearly the form of an *oblate spheroid*. The same investigations also make known the dimensions of the earth. The amount of the oblateness at the poles is measured by the ratio of the difference of the equatorial and polar diameters to the equatorial diameter, which is technically termed the *Oblateness*.

146. The form of the earth has also been determined by other methods, which cannot here be explained. All the results, taken together, indicate an oblateness of  $\frac{1}{305}$ .

The following are the dimensions of the earth in miles :

Radius at the equator . . . . .	3962.6	miles.
Radius at the pole . . . . .	3949.6	"
Difference of equatorial and polar radii	13.0	"
Mean radius, or at 45° latitude, . . .	3956.1	"
Mean length of a degree . . . . .	69.05	"
The fourth part of a meridian . . .	6214.2	"

147. Owing to the elliptical form of a terrestrial meridian, the radius and vertical at a place do not coincide. Let E N Q S (Fig. 30) represent a terrestrial meridian. For any point O situated on this meridian, C O will be the radius, and the normal line Z O N the vertical. The position of the vertical will always be such, that the apparent zenith Z will lie between the true zenith  $z$  and the elevated pole P. The inclination of the radius to the vertical, or the angle C O N, called the reduction of latitude, is greatest at the latitude 45°, and is there equal to about 11½'.

148. The oblateness of the earth occasions some slight modifications in the effects of parallax, which are in some instances to be taken into account in computing the apparent azimuth and zenith distance of a body, from the known co-ordinates of its true place. (These are investigated in the Appendix.)

*Determination of the Latitude and Longitude of a Place.*

149. The latitude and longitude of a place ascertain its situation upon the earth's surface, and are essential elements in many astronomical investigations.

150. *To find the latitude of a place.*

1. *By the zenith distances or altitudes of a circumpolar star at its upper and lower transits.* The principle of this method has already been demonstrated (Art. 59), and shown to be a particular case of a well known principle of arithmetical proportions; the following is a more complete proof of it. Let Z (Fig. 31) represent the zenith, H O R the horizon, P the pole, and S, S' the points at which the upper and lower transits of a circumpolar star take place; H P will be equal to the latitude, and Z P will be equal to the co-latitude. Now, we have,

$$H P = H S + P S, \text{ and } H P = H S' - P S' = H S' - P S$$

$$\text{whence, } 2 H P = H S + H S', \text{ or, } H P = \frac{H S + H S'}{2} \dots (24).$$

In like manner we obtain,

$$Z P = \frac{Z S + Z S'}{2} \dots (25).$$

Wherefore, let the altitudes of a circumpolar star at its upper and lower transits, be measured and corrected for refraction, and their half sum will be the latitude; or, let the zenith distances be measured, and corrected for refraction, and their half sum subtracted from  $90^\circ$  will be the latitude. Stars should be selected that have a considerable altitude at their inferior transit, for, the greater is the altitude the less is the uncertainty as to the amount of the refraction. On this principle the pole star is to be preferred to all others.

2. *By a single meridian altitude or zenith distance.* Let  $s, s', s''$  (Fig. 6) be the points of meridian passage of three different stars, the first to the north of the zenith, the second between the zenith and equator, and the third to the south of the equator: Z E = the latitude, and we have for the three stars,

$$Z E = s E - Z s, \quad Z E = s' E + Z s', \quad Z E = Z s'' - s'' E.$$

Thus, if the zenith distance be called north or south according as the zenith is north or south of the star when on the meridian, in case the zenith distance and declination are of the same name their sum will be equal to the latitude; but if they are of different names, their difference will be the latitude, of the same name with the greater.

This method supposes the declination of the body to be known. The declination of a star or of the sun at any time is, *in practice*, obtained for the solution of this and other problems, by the aid of tables, or is taken by inspection from the English Nautical Almanac or other similar work. If the time of the meridian transit be known, the altitude may be measured by a sextant (Art. 66). The observed altitude must be corrected for refraction, and also for parallax if the body observed is the sun, or moon, or either one of the planets.

This method of finding the latitude is the one most generally employed at sea, the sun being the object observed. As the time of noon is not known with accuracy, several altitudes about the time of noon are taken, and the meridian altitude is *deduced* from these.

151. The astronomical latitude being known, the *reduced latitude* (p. 15, def. 4) may be obtained by subtracting from it the reduction of latitude. For, if  $OC$  (Fig. 30) represents the radius and  $ON$  the vertical, at any place  $O$ , and  $ECQ$  represents the terrestrial equator,  $ONQ$  will be the astronomical latitude,  $OCQ$  the reduced latitude, and  $CON$  the reduction of latitude; and we have

$ONQ = OCQ + CON$ , and  $OCQ = ONQ - CON$  . . (26).  
(For the practical method of resolving this problem, see Prob. XV).

152. There are various methods of finding the longitude of a place, nearly all of which rest upon the following principle:

*The difference at any instant between the local times, (whether sidereal or solar), at any place and on the first meridian, is the longitude of the place, expressed in time; and consequently, also, the difference between the local times at any two places, is their difference of longitude, in time.*

The truth of this principle is easily established. In the first place, we remark that the longitude of a place contains the same number of degrees and parts of a degree as the arc of the celes-

tial equator comprised between the meridian of Greenwich and the meridian of the place. Now, it is 0h. 0m. 0s. of mean solar time, or mean noon at any place, when the mean sun (Art. 39) is on the meridian of that particular place. Therefore, as the mean sun, moving in the equator, recedes from the meridian towards the west at the rate of  $15^{\circ}$  per mean solar hour, when it is mean noon at a place to the *west* of Greenwich, it will be as many hours and parts of an hour *past* mean noon at Greenwich, as is expressed by the quotient of the division of the arc of the celestial equator, or its equal the longitude, by 15. If the place be to the *east*, instead of to the west of Greenwich, when it is mean noon there, it will be as much *before* mean noon at Greenwich as is expressed by the longitude of the place converted into time (as above). In either situation of the place, then, the principle just stated will be true.

It is plain that the equality between the differences of the times and of the longitudes will subsist equally, if sidereal instead of solar time be used.

153. *To find the longitude of a place.*

1. *Let two observers, stationed one at Greenwich, and the other at the given place, note the times of the occurrence of some phenomenon which is seen at the same instant at both places; the difference of the observed times will be the longitude in time. These same observations made at any two places will make known their difference of longitude. If the stations are not distant from each other, a signal, as the flashing of gunpowder, or the firing of a rocket, may be observed. When they are remote from each other, celestial phenomena must be taken. Eclipses of the satellites of Jupiter and of the moon are phenomena adapted to the purpose in question. However, as in these eclipses the diminution of the light of the body is not sudden, but gradual, the longitude cannot be obtained with very great accuracy from observations made upon them.*

2. *Transport a chronometer which has been carefully adjusted to the local time at Greenwich, to the place whose longitude is sought, and compare the time given by the chronometer with the local time of the place. In the same way, by transporting a chronometer from any one place to another, their difference of longitude may be obtained. The error and rate of the*



chronometer must be determined at the outset, and as often afterwards as circumstances will admit, that the error at the moment of the observation may be known as accurately as possible. To insure greater certainty and precision in the knowledge of the time, three or four chronometers are often taken, instead of one only.

This method is much used at sea; the local time being obtained from an observation upon the sun or some other heavenly body, in a manner to be hereafter explained.

3. *Let the Greenwich time of the occurrence of some celestial phenomenon be computed, and note the time of its occurrence at the given place.*

Eclipses of the sun and moon, and of Jupiter's satellites, occultations of the stars by the moon, and the angular distance of the moon from some one of the heavenly bodies, are the phenomena employed. The Greenwich times of the beginning and end of the eclipses of Jupiter's satellites, are published for the solution of the problem of the longitude, in the English Nautical Almanac. Eclipses of the sun and occultations of the stars furnish the most exact determinations of the longitude, but they cannot be used for this purpose unless the longitude is already approximately known.

The explanation, in detail, of the *method of lunar distances*, which is chiefly used at sea, may be found in treatises on Navigation and Nautical Astronomy.

---

## CHAPTER VI.

### OF THE PLACES OF THE FIXED STARS.

154. The place of a fixed star in the sphere of the heavens, is found by ascertaining its true right ascension and declination, which are the co-ordinates of its place. The process of finding the true right ascension and declination of a heavenly body has



already been detailed : the apparent right ascension and declination are found as explained in Arts. 48, 59, and to these are applied the several corrections of refraction, parallax (when sensible), and aberration (Arts. 75, 106, 114).

When right ascensions and declinations found at different times are to be compared together, or employed in the same calculations, as often becomes necessary, they are to be reduced to the same epoch by correcting for precession and nutation (p. 57).

155. It is important to observe, however, that the places of the fixed stars, as at present known, were not obtained by the direct process just referred to, that is, by observing the right ascension and declination, and applying to them at once all the corrections of which we have treated. They were arrived at by successive approximations. The respective corrections were applied in succession as they came to be discovered ; and more accurate results were obtained, as, by the improvement of the instruments, the observations became more and more exact, and as the amount of the corrections came to be known with greater and greater precision.

156. In order to distinguish the fixed stars from each other, they are arranged into groups, called *Constellations*, which are imagined to form the outlines of figures of men, animals, or other objects, from which they are named. Thus one group is conceived to form the figure of a Bear, another of a Lion, a third of a Dragon, and a fourth of a Lyre. The division of the stars into constellations is of very remote antiquity ; and the names given by the ancients to individual constellations are still retained.

The constellations are divided into three classes : *Northern Constellations*, *Southern Constellations*, and *Constellations of the Zodiac*. Their whole number is 91 : Northern 34, Southern 45, and Zodiacal 12. The number of the ancient constellations was but 48. The rest have been formed by modern astronomers, from southern stars not visible to the ancient observers, and others variously situated, which escaped their notice, or were not attentively observed.

157. The stars of a constellation are distinguished from each other by the letters of the Greek alphabet, and in addition to these, if necessary, the Roman letters, and the numbers 1, 2, 3, &c. ; the characters, according to their order, denoting the rela-

tive magnitudes of the stars. Thus,  $\alpha$  Arietis designates the largest star in the constellation Aries;  $\beta$  Draconis, the second star of the Dragon, &c.

Some of the fixed stars have particular names, as *Sirius*, *Aldebaran*, *Arcturus*, *Regulus*, &c.

158. The stars are also divided into classes, or *magnitudes*, according to the degrees of their apparent brightness. The largest or brightest are said to be of the *first magnitude*; the next in order of brightness, of the *second magnitude*; and so on to stars of the *sixth magnitude*, which includes all those that are barely perceptible to the naked eye. All of a smaller kind are called *telescopic* stars, being invisible without the assistance of the telescope. The classification according to apparent magnitude is continued with the telescopic stars down to stars of the *sixteenth magnitude*.

159. The places of the fixed stars are generally expressed by their right ascensions and declinations, but sometimes also by their longitudes and latitudes. A table containing a list of fixed stars, designated by their proper characters, and giving their mean right ascensions and declinations, or their mean longitudes and latitudes, is called a *Catalogue* of those stars.\*

Table ~~LXI~~ is a catalogue of fifty principal fixed stars, and gives xc. their mean right ascensions and declinations for the beginning of the year 1840, as well as their annual variations in right ascension and declination. The annual variations serve to extend the use of the catalogue about 10 years (Art. 135) before and after the epoch for which it is constructed. (See Prob. XVIII.) Every ten years, or thereabouts, a new catalogue must be formed.

160. If the *true* right ascension and declination of a star at a given time be required, correct the mean right ascension and declination found by the catalogue, for *nutation*. (See Art. 132.) And if the *apparent* right ascension and declination be required, correct also for *aberration*. (See Art. 114.)

161. The latitude and longitude of a fixed star or other hea-

---

\* Various catalogues have at different periods been published. The first catalogue was begun by Hipparchus 120 years before the Christian era. The most modern and most accurate catalogues, although not the most extensive, are the catalogues of Lacaille, Bradley, Mayer, and Maskelyne.

venly body are obtained originally by computation from its right ascension and declination.

*To convert the right ascension and declination of a body into its longitude and latitude.*

Let  $E Q$  (Fig. 32) represent the equator,  $E C$  the ecliptic,  $P, K$  the poles of the equator and ecliptic,  $E$  the vernal equinox,  $P S R$  a circle of declination, and  $K S L$  a circle of latitude, both passing through a body  $S$ . The right ascension of the body is  $E R = R$ ; the declination  $R S = D$ ; the longitude  $E L = L$ ; and the latitude  $L S = \lambda$ .  $R E L = \omega$  is the obliquity of the ecliptic, which is one of the essential data of the problem.  $R E S = x$  and  $L E S = y$  are employed as auxiliary angles. In the right angled triangle  $L E S$ , we have by Napier's rules for the solution of right angled triangles,

$$\sin (\text{co. } L E S) = \tan E L \tan (\text{co. } E S);$$

whence,

$$\tan E L = \cos L E S \tan E S, \text{ or, } \tan L = \cos (R E S - \omega) \tan E S;$$

but,

$$\sin (\text{co. } R E S) = \tan E R \tan (\text{co. } E S), \text{ or, } \tan E S = \frac{\tan E R}{\cos R E S};$$

thus,

$$\tan L = \cos (R E S - \omega) \frac{\tan E R}{\cos R E S} = \frac{\cos (x - \omega) \tan R}{\cos x} \dots (27).$$

And to find  $x$ , we have

$$\sin E R = \tan (\text{co. } R E S) \tan R S, \text{ or, } \cot x = \sin R \cot D \dots (28).$$

Again,

$$\sin E L = \tan (\text{co. } L E S) \tan L S, \text{ or } \tan L S = \tan L E S \sin E L, \\ \text{which gives, } \tan \lambda = \tan (x - \omega) \sin L \dots (29).$$

Equation (28) makes known the value of  $x$ , with which we derive the values of  $L$  and  $\lambda$ , by means of equations (27) and (29). In resolving the equations, attention must be paid to the signs of the quantities, which are determined according to the usual trigonometrical rules, it being understood that the declination  $D$  is to be regarded as *negative* when it is *south*.  $x$  is to be taken always less than  $180^\circ$ , and greater or less than  $90^\circ$  according as its cotangent is negative or positive.  $L$  will always be in the same quadrant with  $R$ . The latitude  $\lambda$  will be north or south, according as  $\tan \lambda$  comes out positive or negative.

The apparent or mean obliquity is used, according as the case

refers to true or mean co-ordinates. (For exemplifications of this problem, see Prob. XXIV.)

162. It is now frequently necessary to resolve the converse problem, that is, *to convert the longitude and latitude of a body into its right ascension and declination.*

The triangle R E S (Fig 32) gives,

$$\sin (\text{co. R E S}) = \tan \text{E R} \tan (\text{co E S}) ;$$

whence,

$$\tan \text{E R} = \cos \text{R E S} \tan \text{E S}, \text{ or, } \tan \text{R} = \cos (\text{L E S} + \omega) \tan \text{E S} ;$$

but,

$$\sin (\text{co. L E S}) = \tan \text{E L} \tan (\text{co. E S}), \text{ or } \tan \text{E S} = \frac{\tan \text{E L}}{\cos \text{L E S}} ;$$

thus,

$$\tan \text{R} = \cos (\text{L E S} + \omega) \frac{\tan \text{E L}}{\cos \text{L E S}} = \frac{\cos (y + \omega) \tan \text{L}}{\cos y} \quad (30) ;$$

and to find  $y$ , we have

$$\sin \text{E L} = \tan (\text{co. L E S}) \tan \text{L S}, \text{ or } \cot y = \sin \text{L} \cot \lambda . \quad (31).$$

For the declination, we have

$$\sin \text{E R} = \tan (\text{co. R E S}) \tan \text{R S}, \text{ or, } \tan \text{R S} = \tan \text{R E S} \sin \text{E R} ;$$

or,  $\tan \text{D} = \tan (y + \omega) \sin \text{R} \quad . . . \quad (32).$

The value of  $y$  being derived from equation (31), and substituted in equations (30) and (32), these equations will then make known the values of R and D. The signs of the quantities are determined by the usual trigonometrical rules, the latitude  $\lambda$  being taken *negative* when *south*.  $y$  is always less than  $180^\circ$ , and greater or less than  $90^\circ$  according as its cotangent comes out negative or positive. R will be in the same quadrant as L. The declination will be north or south, according as its tangent comes out positive or negative. (For exemplifications of this problem, see Prob. XXV.)

163. Table XCII contains the mean longitudes and latitudes of some of the principal fixed stars for the beginning of the year 1840, together with their annual variations which serve to make known the mean longitudes and latitudes at any other epoch. (See Prob. XVIII.)

164. There are two principal modes of representing the stars; the one by delineating them on a globe, where each star occupies the spot in which it would appear to an eye placed in the centre of the globe, and where the situations are reversed



when we look down upon them ; the other mode is by a chart, where the stars are generally so arranged as to represent them in positions similar to their natural ones, or as they would appear on the internal concave surface of the globe. The construction of a globe or chart is effected by means of the right ascensions and declinations of the stars. The point which represents the place of a star is found by marking off the right ascension and declination of the star, upon the globe.

All the fixed stars visible to the naked eye, together with some of the telescopic stars, are represented on celestial globes of 12 or 18 inches in diameter.

165. The fixed stars (so called) are not all of them, rigorously speaking, fixed or stationary in the heavens. It has been discovered that many of them have a very slow motion from year to year. These motions of the stars are called *Proper Motions*. The annual variations in right ascension and declination, and in longitude and latitude, given in Tables XC and XCII, are the variations due both to the precession of the equinoxes and the proper motions of the stars.

---

## CHAPTER VII.

### OF THE APPARENT MOTION OF THE SUN IN THE HEAVENS.

166. The sun's declination, and the difference of right ascension of the sun and some fixed star, found from day to day throughout a revolution, are the elements from which the circumstances of the sun's apparent motion are derived.

The motion of the sun as at present known, has been arrived at in the same approximative manner as the places of the fixed stars (Art. 155). It would in fact be theoretically impossible to correct the co-ordinates of the sun's apparent place for precession, nutation, and aberration, in the original determination of the



sun's motion ; for, the knowledge of these corrections pre-supposes some knowledge of the motion of the sun.

167. The curve on the sphere of the heavens passing through the successive positions determined as above from day to day, is the ecliptic. If we suppose it to be a circle, as it appears to be, its position will result from the position of the equinoctial points, and its obliquity to the equator.

168. *To find the obliquity of the ecliptic.*

Let  $EQA$  (Fig. 33) represent the equator,  $ECA$  the ecliptic, and  $OC, OE$  lines drawn through  $O$  the centre of the earth and perpendicular to  $OE$  the line of the equinoxes ; then the angle  $COQ$  will be the obliquity of the ecliptic. This angle has for its measure the arc  $CQ$ , and therefore *the obliquity of the ecliptic is equal to the greatest declination of the sun*. It can but rarely happen that the time of the greatest declination will coincide with the instant of noon at the place where the observations are made, but it must fall within at least twelve hours of the noon for which the observed declination is the greatest. In this interval the change of declination cannot exceed  $4''$ , and therefore the greatest observed declination cannot differ more than  $4''$  from the obliquity. A formula has been investigated, which gives in terms of determinable quantities, the difference between any of the greater declinations and the maximum declination. By *reducing* by means of this formula a number of the greater declinations to the maximum declination, and taking the mean of the individual results, a very accurate value of the obliquity may be found.

169. *To find the position of the vernal or autumnal equinox.*

1. On inspecting the observed declinations of the sun, it is seen that about the 21st of March the declination changes in the interval of two successive noons from south to north. The vernal equinox occurs at some moment of this interval. Let  $RS, R'S'$  (Fig. 34) represent the declinations at the noons between which the equinox occurs : as one is north and the other south, their sum ( $S$ ) will be the daily change of declination at the time of the equinox. Denote the time from noon to noon by  $T$ . Now, to find the interval ( $x$ ) between the noon preced-

ing the equinox and the instant of the equinox, state the proportion

$$S : RS :: T : x = \frac{T \times RS}{S}.$$

Next, take the daily change in right ascension ( $RR'$ ) on the day of the equinox and compute the value of  $RE$ , by the proportion

$$T : x, \text{ or } \frac{T \times RS}{S} :: RR' : RE;$$

on the principle that the declination changes for a day or more proportionally to the time. Add  $RE$  to  $MR$  the observed difference of right ascension (Art. 166) on the day preceding the equinox, and the sum  $ME$  will be the distance of the equinox from the meridian of the star observed in connection with the sun.\*

The position of the autumnal equinox may be found by a similar process, the only difference in the circumstances being that the declination changes from north to south instead of from south to north.

If the value of  $x$  which results from the first proportion, be added to the time of noon on the day preceding the equinox, the result will be the *time* of the equinox.

2. In the triangle  $RES$  (Fig. 33) we have the angle  $RES = \omega$  the obliquity of the ecliptic, and  $RS = D$  the declination of the sun, both of which we may suppose to be known, and we have by Napier's rules,

$$\sin ER = \tan(\text{co. } RES) \tan RS = \cot \omega \tan D \dots (33);$$

whence we can find  $ER$ . And, by taking the sum or difference of  $ER$  and  $MR$ , according as the star is on the opposite side of the sun from the equinox, or the same side, we obtain  $ME$  as before. If this calculation be effected for a number of positions  $S, S', S'', \&c.$  of the sun on different days, and a mean of all the individual results be taken, a more exact value of  $ME$  will be obtained.

$ME$  being accurately known, the precise time of the equinox may readily be deduced from the observed daily variation of right ascension on the day of the equinox.

\* The star is here supposed to be to the west of the sun.

170. The calculations just mentioned rest upon the hypothesis that the ecliptic is a great circle. The close agreement which is found to subsist between the values of  $ME$  deduced from observations upon the sun in different positions  $S, S', S''$ , &c., establishes the truth of this hypothesis. It is also confirmed by the fact, that the right ascensions of the vernal and autumnal equinox differ by  $180^\circ$ , since we may infer from this that the line of the equinoxes passes through the centre of the earth.

171. The *mean obliquity* of the ecliptic is derived from the apparent obliquity, as well as the mean equinox from the true equinox, by correcting for nutation.

172. The mean obliquity at any one epoch having been found, its value at any assumed time may be deduced from it by allowing for the annual diminution of  $0''.46$  (see Table XXII). In like manner, the place of the mean equinox at any given time may be derived from its place once found, by allowing for the annual precession of  $50''.2$ .

The mean obliquity having thus been found for any assumed time, the apparent obliquity at the same time becomes known, by applying the nutation of obliquity. (See Prob. X.)

173. The longitude of the sun may be expressed in terms of the obliquity of the ecliptic and the right ascension or declination. In the triangle  $ERS$ , (Fig. 33)  $ES (= L)$  represents the longitude of the sun supposed to be at  $S$ ,  $ER (= R)$  its right ascension, and  $RS (= D)$  its declination. Now, by Napier's rules,

$$\cos RES = \text{tang } ER \cot ES, \text{ or } \cot ES = \frac{\cos RES}{\text{tang } ER} = \cos RES \cot ER;$$

thus,

$$\cot L = \cos \omega \cot R, \text{ or } \text{tang } L = \frac{\text{tang } R}{\cos \omega} \dots (34).$$

Also,

$$\sin RS = \cos(\text{co. } RES) \cos(\text{co. } ES); \text{ whence, } \sin ES = \frac{\sin RS}{\sin RES};$$

$$\text{or, } \sin L = \frac{\sin D}{\sin \omega} \dots (35).$$

With these formulæ the longitude of the sun may be computed from either its right ascension or declination. (See Prob. XII.)

Formulae (34) and (35) may be written thus,

$$\text{tang } R = \text{tang } L \cos \omega; \sin D = \sin L \sin \omega \dots (36).$$

These formulae will make known the right ascension and declination of the sun, when his longitude is given. (See Prob. XI.) It will be seen in the sequel, that in the present advanced state of astronomical science, the longitude of the sun at any assumed time may be computed from the ascertained laws and rate of the sun's motion.

174. The interval between two successive returns of the sun to the same equinox, or to the same longitude, is called a *Tropical Year*.

And the interval between two successive returns of the sun to the same position with respect to the fixed stars, is called a *Sidereal Year*.

175. It appears from observation that the length of the tropical year is subject to slight periodical variations. The period from which it deviates periodically and equally on both sides, is called the *Mean Tropical Year*. As the changes in the length of the true tropical year are very minute, the length of the mean tropical year is obviously very nearly equal to the mean length of the true tropical year, in an interval during which it passes one or more times through all its different values. In point of fact, it may be found with a very close approximation to the truth, by comparing two equinoxes observed at an interval of 60 or 100 years.

Theory shows that the variation in the length of the tropical year arises from the periodical inequality in the precession of the equinoxes which results from nutation, and certain periodical inequalities in the sun's yearly rate of motion; and thus establishes also, that the mean tropical year, as above defined, is the same as the interval between two successive returns of the sun, supposed to have its mean motion, to the same mean equinox. According to the most accurate determinations, the length of the mean tropical year, expressed in mean solar time, is 365d. 5h. 48m. 47.58s.

176. In a mean tropical year the sun's mean motion in longitude is  $360^\circ$ ; hence, to find his *mean daily motion in longitude*, we have only to state the proportion

$$365\text{d. } 5\text{h. } 48\text{m. } 47\text{s.} : 1\text{d.} :: 360^\circ : x = 59' 8''.33.$$

177. *The sidereal year is longer than the tropical.* For, since the equinox has a retrograde motion of  $50''.23$  in a year, when the sun has returned to the equinox it will not have accomplished a sidereal revolution, into  $50''.23$ . The excess of the sidereal over the tropical year results from the proportion

$$59' 8''.3 : 50''.23 :: 1d. : x = 20m. 23.1s.$$

Thus the length of the mean sidereal year, expressed in mean solar time, is 365d. 6h. 9m. 10.7s.

178. If from the right ascensions and declinations of the sun, found on two successive days, the corresponding longitudes be deduced (equa. 34, 35), and their difference taken, the result will be the sun's daily motion in longitude at the time of the observations. The sun's daily motion in longitude is not the same throughout the year, but, on the contrary, is continually varying. It gradually increases during one half of a revolution, and gradually decreases during the other half, and at the end of the year has recovered its original value. Thus, the greatest and least daily motions occur at opposite points of the ecliptic. They are, respectively,  $61' 10''$  and  $57' 11''$ .

179. The exact laws of the sun's unequable motion cannot be obtained, without a knowledge of the law of variation of the sun's distance, or, what amounts to the same, of the form of the orbit apparently described by the sun in space.

## CHAPTER VIII.

OF THE MOTIONS OF THE SUN, MOON, AND PLANETS,  
IN THEIR ORBITS.

### *Kepler's Laws.*

180. The celebrated astronomer, Kepler, discovered from observation, that the motions of the planets, including the earth, are in conformity with the three following laws:

1. *The areas described by the radius vector of a planet [or*



the line drawn from the sun to the planet] *are proportional to the times.*

2. *The orbit of a planet is an ellipse, of which the sun occupies one of the foci.*

3. *The squares of the times of revolution of the planets are proportional to the cubes of their mean distances from the sun, or of the semi-major axes of their orbits.*

These laws are known by the denomination of *Kepler's Laws*. They were first announced by Kepler as the fundamental laws of the planetary motions, from a *partial* examination only of these motions. The first two he assumed as hypotheses, after that he had discovered that the radius vector and angular motion of a planet were variable, and afterwards verified them, or rather *partially* verified them. They have since been *completely* verified by other astronomers. We shall accordingly adopt these two laws for the present as *hypotheses*, and show in the sequel that they are verified by the results deducible from them.

These laws being established, the third is obtained by simply comparing the known major axes and times of revolution.

181. The apparent motion of the sun in space must be subject to Kepler's first two laws; for, the apparent orbit of the sun is of the same form and dimensions as the actual orbit of the earth, and the law and rate of the sun's motion in its apparent orbit, are the same as the law and rate of the earth's motion. To establish these two facts, let  $E E' A$  (Fig. 35) represent the elliptic orbit of the earth, and  $S$  the position of the sun in space. If the earth move from  $E$  to any point  $E'$ , as it seems to remain stationary at  $E$ , it is plain that the sun will appear to move from  $S$  to a position  $S'$ , on the line  $E S'$  drawn parallel to  $E' S$  the actual direction of the sun from the earth, and at a distance  $E S'$  equal to  $E' S$  the actual distance of the sun from the earth. Thus, for every position of the earth in its orbit, the corresponding apparent position of the sun is obtained by drawing a line parallel to the radius vector of the earth, and equal to it. It follows, therefore, that the area  $S E S'$  apparently described by the radius vector of the sun (or the line drawn from the sun to the earth) in any interval of time, is equal to the area  $E S E'$  actually described by the radius vector of the earth in the same

time ; and, consequently, that the arc  $S S'$  apparently described by the sun in space, is equal to the arc  $E E'$  actually described in the same time by the earth. Whence we conclude, that the apparent motion of the sun in space, and the actual motion of the earth, are the same in every particular.

182. It has been discovered that the motion of the moon in its revolution around the earth, is subject to the same laws as the motion of a planet in its revolution around the sun. We shall assume this to be a fact, and show that our hypothesis is verified by the results to which it leads.

183. That point of the orbit of a planet, which is nearest to the sun, is called the *Perihelion*, and that point which is most distant from the sun, the *Aphelion*. The corresponding points of the moon's orbit, or of the sun's apparent orbit, are called, respectively, the *Perigee* and the *Apogee*.

These points are also called *Apsides* ; the former being termed the *Lower Apsis*, and the latter the *Higher Apsis*. The line joining them is denominated the *Line of Apsides*.

The orbits of the sun, moon, and planets, being regarded as ellipses, the perigee and apogee, or the perihelion and aphelion, are the extremities of the major axis of the orbit.

184. The law of the *angular* motion of a planet about the sun may be deduced from Kepler's first law. Let  $P p A p''$  (Fig. 36) represent the orbit of a planet, considered as an ellipse, and  $p, p'$  two positions of the planet at two instants separated by a short interval of time ; and let  $n$  be the middle point of the arc  $p p'$ . With the radius  $S n$  describe the small circular arc  $l n l'$ , and with the radius  $S b$  equal to unity, describe the arc  $a b$ . It is plain that the two positions  $p, p'$  may be taken so near to each other, that the area  $S p p'$  will be sensibly equal to the circular sector  $S l l'$ . If we suppose this to be the case, as the measure of the sector is  $\frac{1}{2} l n l' \times S n = \frac{1}{2} a b \times \overline{S n}^2$  (substituting for  $l n l'$  its value  $a b \times S n$ ), we shall have

$$\text{area } S p p' = \frac{1}{2} a b \times \overline{S n}^2.$$

When the planet is at any other part of its orbit, as  $n'$ , if  $S p'' p'''$  be an area described in the same time as before, we shall have

$$\text{area } S p'' p''' = \frac{1}{2} a' b' \times \overline{S n'}^2.$$

But these areas are equal according to Kepler's first law :  
hence,  $\frac{1}{2} a b \times S \overline{n}^2 = \frac{1}{2} a' b' \times S \overline{n'}^2 \dots (37);$

and  $a b : a' b' :: S \overline{n'}^2 : S \overline{n}^2,$

that is, *the angular motion of a planet about the sun for a short interval of time, is inversely proportional to the square of the radius vector.*

It results from this that the angular motion is greatest at the perihelion, and least at the aphelion, and the same at corresponding points, on either side of the major axis : also, that it decreases progressively from the perihelion to the aphelion, and increases progressively from the aphelion to the perihelion.

185. Now to compare the true with the mean angular motion, suppose a body to revolve in a circle around the sun, with the mean angular motion of a planet, and to set out at the same instant with it from the perihelion. Let P M A M' represent the elliptic orbit of the planet, and P B a B' the circle described by the body. The position B of this fictitious body at any time will be the *mean place* of the planet as seen from the sun. The two bodies will accomplish a semi-revolution in the same period of time, and therefore be, respectively, at A and a at the same instant ; for, it is obvious that the fictitious body will accomplish a semi-revolution in half the period of a whole revolution, and by Kepler's law of areas, the planet will describe a semi-ellipse in half the time of a revolution. At the onset, the motion of the planet is the most rapid (Art. 184), but it continually decreases until the planet reaches the aphelion, while the motion of the body remains constantly equal to the mean motion. The planet will therefore take the lead, and its angular distance p S B from the body, will increase until its motion becomes reduced to an equality with the mean motion, after which it will decrease until the planet has reached the aphelion A, where it will be zero. In the motion from the aphelion to the perihelion, the angular velocity of the planet will at first be less than that of the body (Art. 184), but it will continually increase, while that of the body will remain unaltered : thus, the body will now get in advance of the planet, and their angular distance p' S B' will increase, as before, until the motion of the planet again attains to an equality with the mean motion, after which it will de-

crease, as before, until it again becomes zero at the perihelion.

It appears, then, that *from the perihelion to the aphelion the true place is in advance of the mean place*, and that *from the aphelion to the perihelion, on the contrary, the mean place is in advance of the true place*.

The angular distance of the true place of a planet from its mean place, as it would be observed from the sun, is called the *Equation of the Centre*. Thus,  $p S B$  is the equation of the centre corresponding to the particular position  $p$  of the planet. It is evident, from the foregoing remarks, that the equation of the centre is zero at the perihelion and aphelion, and greatest at the two points, as  $M$  and  $M'$ , where the planet has its mean motion. The greatest value of the equation of the centre is called the *Greatest Equation of the Centre*.

186. As the laws of the motion of the moon (Art. 182) and of the apparent motion of the sun (Art. 181) are the same as those of a planet, the principles established in the two preceding articles are as applicable to these bodies in their revolution around the earth, as to a planet in its revolution around the sun.

### *Definitions of Terms.*

187. 1. The *Geocentric Place* of a body, is its place as seen from the earth.

2. The *Heliocentric Place* of a body, is its place as it would be seen from the sun.

3. *Geocentric Longitude* and *Latitude* appertain to the geocentric place, and *Heliocentric Longitude* and *Latitude* to the heliocentric place.

4. Two heavenly bodies are said to be *in Conjunction*, when their longitudes are the same, and to be *in Opposition*, when their longitudes differ by  $180^\circ$ . When any one heavenly body is in conjunction with the sun, it is, for the sake of brevity, said to be *in Conjunction*; and when it is in opposition to the sun, to be *in Opposition*.

The planets Mercury and Venus, allowing that their distances from the sun are each less than the earth's distance (Art. 19), can never be in opposition. But they may be in conjunction, either by being between the sun and earth, or by being on the



opposite side of the sun. In the former situation they are said to be in *Inferior Conjunction*, and in the latter, in *Superior Conjunction*.

5. A *Synodic Revolution* of a body, is the interval between two consecutive conjunctions or oppositions.

For the planets Mercury and Venus, a synodic revolution is the interval between two consecutive inferior or superior conjunctions.

6. The *Periodic Time* of a planet, is the period of time in which it accomplishes a revolution around the sun.

7. The *Nodes* of a planet's orbit, or of the moon's orbit, are the points in which the orbit cuts the plane of the ecliptic. The node at which the planet passes from the south to the north side of the ecliptic is called the *Ascending Node*, and is designated by the character  $\Omega$ . The other is called the *Descending Node*, and is marked  $\omega$ .

8. The *Eccentricity* of an elliptic orbit, is the distance between the centre of the orbit and either focus.

#### *Elements of the Orbit of a Planet.*

188. To have a complete knowledge of the motions of the planets, so as to be able to calculate the place of any one of them at any assumed time, it is necessary to know for each planet, in addition to the laws of its motion discovered by Kepler, the position and dimensions of its orbit, its mean motion, and its place at a specified epoch. These necessary particulars of information are subdivided into seven distinct elements, called the *Elements of the Orbit of a Planet*, which are as follows :

1. The longitude of the ascending node.
2. The inclination of the plane of the orbit to the plane of the ecliptic, called the inclination of the orbit.
3. The mean distance of the planet from the sun, or the semi-major axis of its orbit.
4. The eccentricity of the orbit.
5. The heliocentric longitude of the perihelion.
6. The epoch of the planet being at its perihelion, or instead, its mean longitude at a given epoch.
7. The periodic time of the planet.

The first two ascertain the *position of the plane* of the



planet's orbit ; the third and fourth, the *dimensions* of the orbit ; the fifth, the *position of the orbit in its plane* ; the sixth, the *place of the planet at a given epoch* ; and the seventh, its *mean rate of motion*.

189. The elements of the earth's orbit, or of the sun's apparent orbit, are but *five* in number ; the first two of the above mentioned elements being wanting, as the plane of the orbit is coincident with the plane of the ecliptic.

190. The elements of the moon's orbit are the same with those of a planet's orbit, it being understood that the perigee of the moon's orbit answers to the perihelion of a planet's orbit, and that the *geocentric* longitude of the perigee and the *geocentric* longitude of the node of the moon's orbit answer respectively to the heliocentric longitude of the perihelion, and the heliocentric longitude of the node of a planet's orbit.

191. The linear unit adopted, in terms of which the semi-major axes, eccentricities, and radii vectores of the planetary orbits, are expressed, is the mean distance of the sun from the earth, or the semi-major axis of the earth's orbit. When thus expressed, these lines are readily obtained in known measures whenever the mean distance of the sun becomes known. The lines of the moon's orbit are found in terms of the moon's mean distance from the earth, as unity.

*Methods of Determining the Elements of the Sun's Apparent Orbit, or of the Earth's Real Orbit.*

*Mean motion.*

192. The sun's mean daily motion in longitude results from the length of the mean tropical year obtained from observation (Art. 176).

*Semi-major axis.*

193. As we have just stated, the semi-major axis of the sun's apparent orbit is the linear unit, in terms of which the dimensions of the planetary orbits are expressed. Its absolute length is computed from the mean horizontal parallax of the sun.

194. *The horizontal parallax of a body being given, to find its distance from the earth.* We have (equation 9, p. 43),

$$D = \frac{R}{\sin H} ;$$

where *H* represents the horizontal parallax of the body, *D* its dis-

tance from the centre of the earth, and  $R$  the radius of the earth. The parallax of all the heavenly bodies, with the exception of the moon, is so small, that it may, without material error, be taken in this equation in place of its sine. Thus,

$$D = \frac{R}{\sin H} = R \times \frac{1}{H} \dots (38).$$

Again, since 6.2831853 is the length of the circumference of a circle, of which the radius is 1, and 1296000 is the number of seconds in the circumference, we have  $6.2831853 : 1 :: 1296000'' : x = 206264''.8$  = the length of the radius (1) expressed in seconds. Hence, if the value of  $H$  be expressed in seconds,

$$D = R \frac{206264.8}{H} \dots (39).$$

195. In the determination of the sun's parallax, by the process of Arts. 100 and 101, an error of  $2''$  or  $3''$ , equal to about one fourth of the whole parallax, may be committed, so that the distance of the sun, as deduced by equation (39) from his parallax found in that manner, may be in error by an amount equal to one fourth or more of the true distance. There is a much more accurate method of obtaining the sun's parallax, which will be noticed hereafter. It has been found by the method to which we allude, that the horizontal parallax of the sun at the mean distance is  $8''.58$ , which may be relied upon as exact to within a small fraction of a second. We have, then, for the sun's mean distance, or the mean semi-major axis of his orbit,

$$D = R \frac{206264.8}{8''.58} = 24040.19 R = 95,102,992 \text{ miles};$$

taking for  $R$  the mean radius of the earth = 3956 miles.

#### *Eccentricity.*

196. First Method. *By the greatest and least daily motions in longitude.* We have already explained (Art. 179) the mode of deriving from observation the sun's motion in longitude from day to day. Now, let  $v$  = the greatest daily motion in longitude;  $v'$  = the least daily motion in longitude;  $r$  = the least or perigean distance of the sun; and  $r'$  the greatest or apogean distance; and we shall have, by the principle of Art. 174, 124

$$r : r' :: \sqrt{v'} : \sqrt{v};$$

whence,  $r' + r : r' - r :: \sqrt{v} + \sqrt{v'} : \sqrt{v} - \sqrt{v'}$ ;

$$\text{or,} \quad \frac{r' + r}{2} : r' - r :: \frac{\sqrt{v} + \sqrt{v'}}{2} : \sqrt{v} - \sqrt{v'};$$

but,

$$\frac{r' + r}{2} = \text{semi-major axis} = 1; \text{ and } r' - r = 2 (\text{eccentricity}) = 2e.$$

$$\text{Thus,} \quad 1 : 2e :: \frac{\sqrt{v} + \sqrt{v'}}{2} : \sqrt{v} - \sqrt{v'};$$

$$\text{and} \quad e = \frac{\sqrt{v} - \sqrt{v'}}{\sqrt{v} + \sqrt{v'}} \dots (40).$$

The greatest and least daily motions are, respectively, (at a mean)  $61'.165$  and  $57'.192$ . Substituting, we have,

$$e = 0.016791.$$

The eccentricity may also be obtained from the *greatest and least apparent diameters* by a process similar to the foregoing, on the principle that the distances of the sun at different times are inversely proportional to his corresponding apparent diameters.

The greatest apparent diameter of the sun is  $32' 35''.6$ , and the least apparent diameter  $31' 31''.0$ .

197. Second Method. *By the greatest equation of the centre.*

1. *To find the greatest equation of the centre.* Let  $L$  = the true longitude, and  $M$  = the mean longitude, at the time the true and mean motions are equal between the perigee and apogee;  $L'$  = the true longitude and  $M'$  = the mean longitude, when the motions are equal between the apogee and perigee; and  $E$  = the greatest equation of the centre. Then (Art. 185),

$$L = M + E, \text{ and } L' = M' - E;$$

$$\text{whence,} \quad L' - L = M' - M - 2E;$$

$$\text{and} \quad E = \frac{(M' - M) - (L' - L)}{2} \dots (41).$$

About the time of the greatest equation, the sun's true motion and consequently the equation of the centre, continues very nearly the same for two or three days; we may, therefore, with but slight error, take the noon, when the sun is on either side of the line of apsides, that separates the two days on which the motions in longitude are most nearly equal to  $59' 8''$ , as the epoch of the greatest equation.

The longitude  $L$  or  $L'$  at either epoch thus ascertained, results

from the observed right ascension and declination.  $M' - M$  = the mean motion in longitude in the interval of the epochs, and is found by multiplying the number of mean solar days and fractions of a day comprised in the interval, by  $59' 8''.330$ , the mean daily motion in longitude.

For example. From observations upon the sun, made by Dr. Maskelyne in the year 1775, it is ascertained in the manner just explained, that the sun was near its greatest equation at noon, or at 0h. 3m. 35s. mean solar time on the 2d April, and at noon on the 31st, or at 23h. 49m. 35s. mean solar time on the 30th of September. The observed longitudes were, at the first period,  $12^\circ 33' 39''.06$ , and at the second  $188^\circ 5' 44''.45$ . The interval of time between the two epochs is 182d. — 14m.

Mean motion in 182d. — 14m. . . . .  $179^\circ 22' 41''.56$

Difference of two longitudes . . . . .  $175 \quad 32 \quad 5.39$

Diff. . . . .  $2 \quad ) \quad 3 \quad 50 \quad 36.17$

Greatest equation of centre . . . . .  $1 \quad 55 \quad 18.08$

More accurate results are obtained, by reducing observations made during several days before and after the epoch of the greatest equation, and taking the mean of the different values of the greatest equation thus obtained. According to M. Delambre, the greatest equation was in 1775,  $1^\circ 55' 31''.66$ .

2. The eccentricity of an orbit may be derived from the greatest equation of the centre by means of the following formula :

$$e = \frac{K}{2} - \frac{11 K^3}{3.2^8} - \frac{587 K^5}{3.5.2^{16}} - \&c. \dots (42),$$

in which K stands for the expression  $\frac{E}{57^\circ.2957795}$  (E being the greatest equation of the centre). In the case of the sun's orbit, K being a small fraction, all its powers beyond the first may be omitted. Thus, retaining only the first term of the series, and taking  $E = 1^\circ 55' 31''.66$  the greatest equation in 1775, we have,

$$e = \frac{K}{2} = \frac{1^\circ 55' 31''.66}{2 \times 57^\circ.2957795} = .016803.$$

198. From observations made at distant periods, it is discovered that the equation of the centre, and consequently the eccentricity, is subject to a continual slow diminution. The amount of the

diminution of the greatest equation in a century, called the *secular* diminution, is estimated by Delambre, at  $17''.2$ .

*Longitude and epoch of the perigee.*

199. As the sun's angular velocity is the greatest at the perigee, the longitude of the sun at the time its angular velocity is greatest, will be the longitude of the perigee. The time of the greatest angular velocity may easily be obtained within a few hours, by means of the daily motions in longitude, derived from observation (Art. 178).

200. The more accurate method of determining the longitude and epoch of the perigee, rests upon the principle that the apogee and perigee are the only two points of the orbit whose longitudes differ by  $180^\circ$ , in passing from one to the other of which the sun employs just half a year. This principle may be inferred from Kepler's law of areas, for it is a well known property of the ellipse, that the major axis is the only line drawn through the focus, that divides the ellipse into equal parts, and by the law in question equal areas correspond to equal times.

201. By a comparison of the results of observations made at distant epochs, it is discovered that the longitude of the perigee is continually increasing at a mean rate of  $61''.5$  per year. As the equinox retrogrades  $50''.2$  in a year, the perigee must then have a direct motion in space of  $11''.3$  per year.

It will be seen, therefore, that the interval between the times of the sun's passage through the apogee and perigee is not, strictly speaking, half a sidereal year, but exceeds this period by the interval of time employed by the sun in moving through an arc of  $5''.6$  the motion of the apogee and perigee in longitude in half a year.

202. According to the most exact determinations, the mean longitude of the perigee of the sun's orbit at the beginning of the year 1800, was  $279^\circ 30' 8''.39$ .

203. The heliocentric longitude of the perihelion of the earth's orbit, is equal to the geocentric longitude of the perigee of the sun's apparent orbit minus  $180^\circ$ . For, let AEP (Fig. 35) be the earth's orbit, and PV the direction of the vernal equinox. When the earth is in its perihelion P, the sun is in its perigee S, and we have the heliocentric longitude of the perihelion,  $VSP = VPL =$



angle  $a b c - 180^\circ =$  geocentric longitude of the sun's perigee —  $180^\circ$ .\*

204. The epoch and mean longitude of the perigee of the sun's orbit being once found, the sun's mean longitude at any assumed epoch is easily obtained by means of the mean motion in longitude.

*Methods of determining the Elements of the Moon's Orbit.*

*Longitude of the node.*

205. In order to obtain the longitude of the moon's ascending node, we have only to find the longitude of the moon at the time its latitude is zero and the moon is passing from the south to the north side of the ecliptic; and this may be deduced from the longitudes and latitudes of the moon, derived from observed right ascensions and declinations, by methods precisely analogous to those by which the right ascension of the sun, at the time its declination is zero and it is passing from the south to the north of the equator, or the position of the vernal equinox, is ascertained. (See Art. 169.)

*Inclination of the orbit.*

206. Amongst the latitudes computed from the moon's observed right ascensions and declinations, the greatest measures the inclination of the orbit. It is found to be about  $5^\circ$ ; sometimes a little greater, and at other times a little less.

*Mean motion.*

207. With the longitudes of the moon, found from day to day, it is easy to obtain the interval from the time at which the moon has any given longitude till it returns to the same longitude again. This interval is called a *Tropical Revolution* of the moon. It is found to be subject to considerable periodical variations, and thus one observed tropical revolution may differ materially from the mean period. In order to obtain the mean tropical revolution, we must compare two longitudes found at distant epochs. Their difference, augmented by the product of  $360^\circ$  by the number of revolutions performed in the interval of the epochs, will be the mean motion in longitude in the interval, from which the mean

---

\* It is plain that the same relation subsists between the heliocentric longitude of the earth and the geocentric longitude of the sun in every other position of the earth in its orbit.

motion in 100 years or 36525 days, called the *Secular* motion, may be obtained by a simple proportion. The secular motion being once known, it is easy to deduce from it the period in which the motion is  $360^\circ$ , which is the mean topical revolution.

It should be observed, however, that to find the precise mean secular motion in longitude, it is necessary to compare the mean longitudes instead of the true. Now, the true longitude of the moon at any time having been found, the mean longitude at the same time is derived from it, by correcting for the equation of the centre, and certain other periodical inequalities of longitude hereafter to be noticed. But this cannot be done, even approximately, until the theory of the moon's motions is known with more or less accuracy.

208. The longitude of the moon, at certain epochs, may be very conveniently deduced from observations upon lunar eclipses. For, the time of the middle of the eclipse is very near the time of opposition when the longitude of the moon differs  $180^\circ$  from that of the sun, and the longitude of the sun results from the known theory of its motions. The recorded observations of the ancients upon the times of the occurrence of eclipses, are the only observations that can now be made use of, for the direct determination of the longitude of the moon at an ancient epoch.

209. The mean tropical revolution of the moon is found to be  
 $27.321582\text{d. or } 27\text{d. } 7\text{h. } 43\text{m. } 4.7\text{s.}$

Hence  $27.321582\text{d.} : 1\text{d.} :: 360^\circ : 13^\circ.17639. = 13^\circ 10' 35''.0 =$   
 moon's mean daily motion in longitude.

210. Since the equinox has a retrograde motion, the *sidereal* revolution of the moon must exceed the tropical revolution, as the sidereal year exceeds the tropical year. The excess will be equal to the time employed by the moon, in describing the arc of precession answering to a revolution of the moon. Thus,

$$365.25\text{d.} : 50''.2 :: 27.3\text{d.} : 3''.75 = \text{arc of precession,}$$

and  $13^\circ.17 :: 1\text{d.} :: 3''.75 : 6.9\text{s.} = \text{excess.}$

Wherefore, the mean sidereal revolution of the moon is  $27\text{d. } 7\text{h. } 43\text{m. } 11.6\text{s.}$

211. It has been found, by determining the moon's mean rate of motion for periods of various lengths, that it is subject to a continual slow acceleration. This acceleration will not, however, be indefinitely progressive: Laplace has investigated its physical

cause, and shown from the principles of Physical Astronomy, that it is really a periodical inequality in the moon's mean motion, which requires an immense length of time to go through its different values.

The mean motion given in Art. 209 answers to the commencement of the present century.

*Longitude of the perigee, eccentricity, and semi-major axis.*

212. The methods of determining these elements of the moon's orbit, are similar to those by which the corresponding elements of the sun's orbit are found. It is to be observed, however, that for the longitudes of the sun, which are laid off in the plane of the ecliptic, in the case of the moon, corresponding angles are laid off in the plane of its orbit. These angles are reckoned from a line drawn making an angle with the line of nodes equal to the longitude of the ascending node, and are called *Orbit Longitudes*. The orbit longitude is equal to the moon's angular distance from the ascending node plus the longitude of the ascending node.

213. *The ecliptic longitude of the moon at any time being given, to find the orbit longitude.*

Let  $V'NM$  (Fig. 38) represent the moon's orbit referred to the sphere of the heavens,  $VNm$  the ecliptic,  $V$  the vernal equinox, and  $V'$  the corresponding point from which the orbit longitudes are reckoned. The orbit longitude  $V'NM = V'N + NM = VN + NM = \text{long. of node} + NM$ . Now, by Napier's rules,

$$\cos MNm = \cot NM \tan N m ;$$

$$\text{or,} \quad \cot NM = \cos MNm \cot N m . . . (42).$$

$Nm$  = ecliptic long. — long. of node ; and  $MNm$  = inclination of orbit.

214. The horizontal parallax of the moon, like almost every other element of astronomical science, is subject to periodical changes of value. It varies not only during one revolution, but also from one revolution to another. The fixed and mean parallax about which the true parallax may be conceived to oscillate, answers to the mean distance, that is, the distance about which the true distance varies periodically, and is called the *Constant of the Parallax*. It is, for the equatorial radius of the earth,  $57' 0''.9$  ; from which we find by equation (39) the

mean distance of the moon from the earth to be 60.3 radii, or about 240,000 miles.

The greatest and least parallaxes of the moon are  $61' 24''$  and  $53' 48''$ .

215. The eccentricity of the moon's orbit is more than three times as great as that of the sun's orbit. Its greatest equation exceeds  $6^\circ$ .

*Mean longitude at an assigned epoch.*

216. We have already explained (Art. 207) the principle of the determination of the mean longitude of the moon from an observed true longitude. Now, when the mean longitude at any one epoch whatever becomes known, the mean longitude at any assigned epoch is easily deduced from it by means of the mean motion in longitude.

*Methods of determining the Elements of a Planet's Orbit.*

217. The methods of determining the elements of the planetary orbits suppose the possibility of finding the heliocentric longitude and the radius vector of the earth for any given time. Now, the elements of the earth's orbit having been found by the processes heretofore detailed, the longitude may be computed by means of Kepler's first law, and the radius vector from the polar equation of the elliptic orbit. The manner of effecting such computation will be considered hereafter, at present the possibility of effecting it will be taken for granted.

*Heliocentric longitude of the ascending node.*

218. When the planet is in either of its nodes, its latitude is zero. It follows, therefore, that the longitude of the planet at the time its latitude is zero, is the geocentric longitude of the node at the time the planet is passing through it. Now, if the right ascension and declination of the planet be observed from day to day, about the time it is passing from one side of the ecliptic to the other, and converted into longitude and latitude, the time at which the latitude is zero, and the longitude at that time may be obtained by a proportion. When the planet is again in the same node, the geocentric longitude of the node may again be found in the same manner as before. On account of the different position of the earth in its orbit, this longitude will differ from the former.

Now, if two geocentric longitudes of the same node be found, its heliocentric longitude may be computed.

Let S (Fig. 39) be the sun, N the node, and E one of the positions of the earth for which the geocentric longitude of the node V E N is known. Denote this angle by G, the sun's longitude V E S by S, and the radius vector S E by  $r$ . Also, let E' be the other position of the earth, and denote the corresponding quantities for this position, V E' N, V E' S, and S E', respectively, by  $G'$ ,  $S'$ , and  $r'$ . Let the radius vector of the planet when in its node, or S N = V; and the heliocentric longitude of the node, or V S N = X. The triangle S N E gives,

$$\sin S N E : \sin S E N :: S E : S N;$$

but  $S E N = V E S - V E N = S - G;$

and,  $S N E = V A N - V S N = V E N - V S N = G - X;$

hence,  $\sin (G - X) : \sin (S - G) :: r : V;$

or,  $r \sin (S - G) = V \sin (G - X) \dots (43).$

In like manner,  $r' \sin (S' - G') = V \sin (G' - X);$

dividing,  $\frac{r \sin (S - G)}{r' \sin (S' - G')} = \frac{\sin (G - X)}{\sin (G' - X)};$

or,  $\frac{r \sin (S - G)}{r' \sin (S' - G')} = \frac{\sin G \cos X - \sin X \cos G}{\sin G' \cos X - \sin X \cos G'} = \frac{\sin G - \cos G \tan X}{\sin G' - \cos G' \tan X};$

whence,

$$\tan X = \frac{r \sin (S - G) \sin G' - r' \sin (S' - G') \sin G}{r \sin (S - G) \cos G' - r' \sin (S' - G') \cos G} \dots (44).$$

Equation (43) gives  $V = \frac{r \sin (S - G)}{\sin (G - X)} \dots (45).$

#### *Inclination of the orbit.*

219. The longitude of the node having been found by the preceding or some other method, compute the day on which the sun's longitude will be the same or nearly the same: the earth will then be on the line of the nodes. Observe on that day the planet's right ascension and declination, and deduce the geocentric longitude and latitude. Let E N p (Fig. 40) be the plane of the ecliptic, V the vernal equinox, S the sun, N the node, E the earth on the line of nodes, and P the planet as referred to



the celestial sphere, from the earth. Let  $\lambda$  denote the geocentric latitude  $Pp$ ;  $E$  the arc  $Np = Vp - VN =$  geo. long. of planet — long. of node; and  $I$  the inclination  $PNp$ . The right angled triangle  $PNp$  gives

$$\sin Np = \tan Pp \cot PNp = \tan \lambda \cot I;$$

hence,  $\cot I = \frac{\sin E}{\tan \lambda}$ , and  $\tan I = \frac{\tan \lambda}{\sin E} \dots (46).$

220. It will be understood, that to obtain an exact result, we must compute the precise time of the day at which the longitude of the sun is the same as that of the node, and then, by means of their observed daily variations, correct the longitude and latitude of the planet for the variations in the interval between the time thus ascertained and the time of the observation above mentioned.

#### *Periodic time.*

221. The interval from the time the planet is in one of its nodes till its return to the same, gives the periodic time or sidereal revolution.

222. Another and more accurate method is to observe the length of a synodic revolution (p. 82), and to compute the periodic time from this. If we compare the time of a conjunction which has been observed in modern times, with that of a conjunction observed by the earlier astronomers, and divide the interval between them by the number of synodic revolutions contained in it, we shall have the mean synodic revolution with great exactness, from which the mean periodic time may be deduced.\*

*To find the heliocentric longitude and latitude, and the radius vector, for a given time.*

223. The earth being in constant motion in its orbit, and being thus at different times very differently situated with regard to the other planets, as well in respect to distance as direction, it is necessary for the purpose of comparing the observations made upon these bodies with each other, to refer them all to one common point of observation. As the sun is the fixed centre about which the revolutions of the planets are performed, it is

---

\* We shall, in the sequel, investigate the equation that expresses the relation between the synodic revolution and the periodic time.

the point best suited to this purpose, and accordingly it is to the sun that the observations are in reality referred. The reduction of observations from the earth to the sun, as it is actually performed, consists in the deduction of the heliocentric longitude and latitude from the geocentric longitude and latitude, these being derived from the observed right ascension and declination. We will now show how to effect this deduction, supposing that the longitude of the node and the inclination of the orbit are known. Let  $N P$  (Fig. 41) be part of the orbit of a planet,  $S N C$  the plane of the ecliptic,  $N$  the ascending node,  $S$  the sun,  $E$  the earth, and  $P$  the planet; also, let  $P \pi$  be a perpendicular let fall from  $P$  upon the plane of the ecliptic, and  $E V$ ,  $S V$  the direction of the vernal equinox. Let  $\lambda = P E \pi$  the geocentric latitude of the planet;  $l = P S \pi$  its heliocentric latitude;  $G = V E \pi$  its geocentric longitude;  $L = V S \pi$  its heliocentric longitude;  $S = V E S$  the longitude of the sun;  $N = V S N$  the heliocentric longitude of the node;  $I = P N C$  the inclination of the orbit;  $r = S E$  the radius vector of the earth; and  $v = S P$  the radius vector of the planet.

The point  $\pi$  is called the *reduced place* of the planet, and  $S \pi$  its *curtate distance*. All the angles of the triangle  $S E \pi$  have also received particular appellations:  $S \pi E$  the angle subtended at the reduced place of the planet by the radius of the earth's orbit, is called the *Annual Parallax*,  $S E \pi$  the *Elongation*, and  $E S \pi$  the *Commutation*. Let  $A = S \pi E$ ,  $E = S E \pi$ , and  $C = E S \pi$ . Draw  $S \pi'$  parallel to  $E \pi$ : then  $A = \pi S \pi' = V S \pi - V S \pi' = V S \pi - V E \pi = L - G$ ;  $E = V E \pi - V E S = G - S$ ;  $C = V S E - V S \pi = 180^\circ + V S E' - V S \pi = 180^\circ + V E S - V S \pi = 180^\circ + S - L = T - L$  (putting  $T = 180^\circ + S$ ).

1. *For the latitude.* The triangles  $E P \pi$ ,  $S P \pi$  give

$$E \pi \tan \lambda = P \pi = S \pi \tan l, \text{ whence } \frac{\tan \lambda}{\tan l} = \frac{S \pi}{E \pi};$$

$$\text{but, } S \pi : E \pi :: \sin E : \sin C, \text{ or, } \frac{S \pi}{E \pi} = \frac{\sin E}{\sin C};$$

$$\text{substituting, } \frac{\tan \lambda}{\tan l} = \frac{\sin E}{\sin C};$$

$$\text{whence, } \tan \lambda \sin C = \tan l \sin E \dots (46).$$

$$\text{or, } \tan \lambda \sin (T - L) = \tan l \sin (G - S) \dots (47).$$

Again, the triangle  $N P p$  gives, by Napier's rules,

$$\sin N p = \cot P N P \tan P p, \text{ or, } \sin (L - N) = \cot I \tan l. \quad (48).$$

Either of the equations (47) and (48) will give the value of  $l$ , when the longitude  $L$  is known.

2. *For the longitude.* If we substitute in equation (47) the value of  $\tan g l$ , given by equation (48), and replace  $(G - S)$  by  $E$ , we have,

$$\tan g \lambda \sin (T - L) = \sin (L - N) \tan g I \sin E ;$$

but  $T - L = (T - N) - (L - N) = D - (L - N)$ , (denoting  $(T - N)$  by  $D$ ) ; substituting and designating  $L - N$  by  $x$ ,

$$\tan g \lambda \sin (D - x) = \sin x \tan g I \sin E ;$$

whence,

$$\tan g \lambda \sin D \cos x - \tan g \lambda \cos D \sin x = \tan g I \sin E \sin x ;$$

$$\text{or, } \tan g \lambda \sin D - \tan g \lambda \cos D \tan g x = \tan g I \sin E \tan g x ;$$

$$\text{which gives, } \tan g x = \frac{\tan g \lambda \sin D}{\tan g \lambda \cos D + \tan g I \sin E} \dots (49.)$$

Substituting the values of  $x$ ,  $D$  and  $E$ , we have finally,

$$\tan g (L - N) = \frac{\tan g \lambda \sin (T - N)}{\tan g \lambda \cos (T - N) + \tan g I \sin (G - S)} \quad (50).$$

As  $N$  is known, the value of  $L$  will result from this equation.

224. The co-ordinates employed to fix the position of a planet in the plane of its orbit, are its orbit longitude (Art. 212) and its radius vector, both of which result from the heliocentric longitude and latitude, the longitude of the node and the inclination of the orbit being known. In Fig. 41,  $V' N P$  represents the orbit longitude, and  $S P (=v)$  the radius vector, for the position  $P$ . Now, the triangle  $P S \pi$  gives,

$$S P = \frac{S \pi}{\cos P S \pi}, \text{ or, } v = \frac{S \pi}{\cos l} ;$$

and the triangle  $E S \pi$  gives,

$$\sin A : \sin E : S E : S \pi = \frac{S E \sin E}{\sin A} = \frac{r \sin E}{\sin A} ;$$

$$\text{whence, by substitution, } v = \frac{r \sin E}{\sin A \cos l} = \frac{r \sin (G - S)}{\sin (L - G) \cos l} \quad (51).$$

The orbit longitude  $L' = N P + \text{long. of node} \dots (52).$

And to find  $N P$ , the triangle  $N P p$  gives,

$$\cos P N p = \cot N P \tan g N p, \text{ or } \tan g N P = \frac{\tan g N p}{\cos I} \dots (52) ;$$

and,  $N p = \text{long. of planet} - \text{long. of node} \dots (52).$

225. The heliocentric longitude may be obtained in a very simple manner, if the observations be made upon the planet at the time of *conjunction* or *opposition*; for, it will then either be equal to the geocentric longitude or differ  $180^\circ$  from it.

When the heliocentric longitude is found, the latitude may be computed by equation (47) or (48). Equation (51) will disappear in conjunctions and oppositions; but the radius vector (S P) may be computed from the triangle E S P (Fig. 42): for, the side S E the radius vector of the earth, is known, as well as the angle S E P the geocentric latitude of the planet, and the angle E S P =  $180^\circ - P S p = 180^\circ - \text{heliocentric lat.}$

226. The radius vector of either of the inferior planets\* at the time of maximum elongation, may be approximately deduced from the amount of the maximum elongation, determined from observation. The elongation of an inferior planet at any time, is equal to the difference of the geocentric longitudes of the planet and sun, and is therefore easily obtained. Let N P P' (Fig. 43) represent the orbit of an inferior planet, supposed to lie in the plane of the ecliptic. The line E P drawn from the earth to the planet, will at the time of maximum elongation be tangent to the orbit; and thus, if the greatest value of the elongation be observed, we shall have in the right angled triangle E P S, the line E S, and the angle S E P, from which the radius vector S P may be computed.

As the earth and planet are in motion, the greatest elongation will occur at different points of the planet's orbit, and therefore we may find by the foregoing process different radii vectores.

*Longitude of the perihelion, eccentricity, and semi-major axis.*

227. The longitude of the perihelion, the eccentricity, and the semi-major axis, may be derived from the heliocentric orbit longitude and the radius vector found for three different times.

Let S P, S P', S P'' (Fig. 44) be the three given radii vectores, V' S P, V' S P', V' S P'', the three given longitudes, and A B the line of apsides of the planet's orbit. Let the angles P S P', P S P'', which are known, be represented by  $m$ ,  $n$ , and the

---

\* An *Inferior* planet is one whose orbit lies within the orbit of the earth.

angle B S P, which is unknown, by  $x$ ; and let the three radii vectores S P, S P', S P'' be denoted by  $v, v', v''$ ; the semi-major axis A C by  $a$ , and the ratio of the eccentricity to the semi-major axis by  $e$ : then, the three unknown quantities which are to be determined, are  $a, e$ , and the angle  $x$ , and the general polar equation of the ellipse furnishes for their determination the three equations:

$$v = \frac{a(1 - e^2)}{1 + e \cos x} \dots (53).$$

$$v' = \frac{a(1 - e^2)}{1 + e \cos(x + m)} \dots (54).$$

$$v'' = \frac{a(1 - e^2)}{1 + e \cos(x + n)} \dots (55).$$

Equating the values of  $a(1 - e^2)$  obtained from equations (53, and (54), we have,

$$v + v e \cos x = v' + v' e \cos(x + m),$$

or, 
$$e = \frac{v' - v}{v \cos x - v' \cos(x + m)} \dots (56).$$

In like manner from (53) and (55),

$$e = \frac{v'' - v}{v \cos x - v'' \cos(x + n)} \dots (57).$$

Let  $v' - v = p$ , and  $v'' - v = q$ ; then, by equating the second members of equations (56), (57), and transforming, we obtain,

$$\begin{aligned} \frac{p}{q} &= \frac{v \cos x - v' \cos(x + m)}{v \cos x - v'' \cos(x + n)} \\ &= \frac{v \cos x - v' \cos m \cos x + v' \sin m \sin x}{v \cos x - v'' \cos n \cos x + v'' \sin n \sin x} \\ &= \frac{v - v' \cos m + v' \sin m \tan x}{v - v'' \cos n + v'' \sin n \tan x}, \end{aligned}$$

$$\text{whence, } \tan x = \frac{p(v - v'' \cos n) - q(v - v' \cos m)}{q v' \sin m - p v'' \sin n} \dots (58).$$

The value of  $x$  being found by this equation, and subtracted from the orbit longitude of the planet in the first position P, the result will be the orbit longitude of the perihelion. Also,  $x$  being known,  $e$  may be computed from either of the equations (56) and (57). And hence again, the semi-major axis from equation (53), (54), or (55).



228. The semi-major axis or mean distance from the sun, may also be had by taking the mean of a great number of radii vectores found for every variety of position of the planet in its orbit.

229. Now that Kepler's third law has been established by investigations in Physical Astronomy, it furnishes the most accurate method of finding the mean distance of a planet from the sun. Thus, let  $P$  = the periodic time of the planet, and  $a$  = its mean distance; then, the length of the sidereal year being 365.256374 days (Art. 177),

$$(365.256374 \text{ d.})^2 : P^2 :: 1^3 : a^3;$$

whence, 
$$a = \left( \frac{P}{365.256374 \text{ d.}} \right)^{\frac{2}{3}} \dots (59).$$

*Epoch of a planet being at the perihelion of its orbit.*

229. From several observations upon the planet, about the time it has the same longitude as the perihelion, the correct time of its being at the perihelion may be easily determined by proportion.

230. *The mean longitude at an assigned epoch* is obtained upon the same principles as the mean longitude of the sun or moon (Arts. 204, 207.)

#### *Remarks.*

231. The foregoing methods of determining the elements of a planet's orbit suppose observations to be made at two or more successive returns of the planet to its node. It is possible, by certain methods of trial and conjecture, to derive the elements of a planetary orbit, with tolerable accuracy, from observations continued during only a part of a revolution, and without waiting for a passage of the planet through its node. It was thus that Lalande determined the elements of the orbit of Uranus, with a near approximation to the truth, within one year of the period of the first discovery of that planet by Sir W. Herschel.

#### *Mean Elements and their Variations.*

232. The elements of the planetary orbits, obtained by the foregoing processes, are the true elements at the periods when the observations are made. Upon determining them at different periods, it appears that they are subject to minute variations. A comparison of the values found at various distant epochs

shows that they are slowly changing from century to century, and that the changes experienced during equal long periods of time are very nearly the same. The amount of the variation of an element in a period of 100 years is called its *Secular Variation*. Upon reducing the elements, found at different times, to the same epoch, by allowing for the proportional parts of the secular variations, the different results for each element are found to differ slightly from each other, which shows that the elements are also subject to slight periodical variations. These variations being very minute, the true elements can never differ much from the mean, or those from which they deviate periodically and equally on both sides.

The mean elements at an assigned epoch may be had by finding the true elements at various times, and reducing them to the given epoch, by making allowance for the proportional parts of the secular variations, and then taking for each element the mean of all the particular values obtained for it.

233. A comparison of the mean values of the same element, found at distant epochs, makes known the variation of its mean value in the interval between them, from which the *secular variation* may be deduced by simple proportion.

234. The elements of the moon's orbit are also subject to continual variations. These are, for the most part, periodic, and are far greater than the variations of the corresponding elements of a planet's orbit. It will be seen then, that in determining the mean elements, a much greater number of observations will be required than in the case of a planetary orbit. The mean node and perigee have a rapid and nearly uniform progressive motion. Theory shows that the other mean elements, with the exception of the semi-major axis, are subject to secular variations, but their effect has hitherto been very inconsiderable.

235. The mean elements which have been derived as above directly from observation, have subsequently been verified and corrected, by comparing the computed with the observed places of the planet; and for this purpose many thousands of observations have been made.

236. Tables II and III contain the elements of the orbits of the principal planets, and of the moon's orbit, together with their secular variations, for the beginning of the year 1801; and

also the elements of the orbits of the four small planets, Vesta, Juno, Ceres and Pallas, for the beginning of the year 1820.

If an element be desired for any time different from the epoch of the table, we have only to allow for the proportional part of the secular variation in the interval between the given time and the epoch of the table.

237. It will be seen, on inspecting Table II, that the mean distances of the planets from the sun, or the semi-major axes of their orbits, are the only elements that are invariable. The rest are subject to minute secular variations. The nodes have all retrograde motions. The perihelia, on the contrary, have direct motions, with the single exception of the perihelion of the orbit of Venus, which has a retrograde motion. The eccentricities of some of the orbits are increasing, of others diminishing. That of the earth's orbit is diminishing.

The node of the moon's orbit has a retrograde motion, and the perihelion a direct motion. The former accomplishes a tropical revolution in 6788.50982 days, or about 18 years 214 days; and the latter in 3231.4751 days, or in about 8 years 309 days. The mean motion of the node, and the mean motion of the perigee, are both subject to a slow secular diminution.

---

## CHAPTER IX.

OF THE DETERMINATION OF THE PLACE OF A PLANET, OR OF THE SUN, OR MOON, FOR A GIVEN TIME, BY THE ELLIPTICAL THEORY; AND OF THE VERIFICATION OF KEPLER'S LAWS.

*Place of a Planet, or of the Sun, or Moon, in its Orbit.*

238. The angle contained between the line of apsides of a planet's orbit and the radius vector, as reckoned from the perihelion towards the east, is called the *True Anomaly*. Thus, let B P A P' (Fig. 45) represent the orbit, B the perihelion, and

P the position of the planet; then, B S P is its true anomaly. The angle contained between the line of apsides and the mean place of the planet, also reckoned from the perihelion towards the east, is called the *Mean Anomaly*. Thus, let M be the mean place of a planet at the time P is its true place, and B S M will be its mean anomaly.

Describe a circle B *p* A on the line of apsides as a diameter; through P draw *p* P D perpendicular to the line of apsides, and join *p* and C: the angle B C *p*, which the line thus determined makes with the line of apsides, is called the *Eccentric Anomaly*.

The corresponding angles appertaining to the sun's apparent orbit, and to the moon's orbit, have received the same appellations.

239. The interval between two consecutive returns of a body to either apsis of its orbit, is called the *Anomalistic Revolution*. The anomalistic revolution of the earth, or of the sun in its apparent orbit, is termed also the *Anomalistic Year*.

240. The periodic time, or the mean motion of a body, and the motion of the apsis of its orbit, being known, the anomalistic revolution may be easily computed. Let *m* = the sidereal motion of the apsis answering to the periodic time, and M = the mean daily motion of the planet; then,

M : 1 d. :: *m* : *x* = diff. of anomalistic rev. and periodic time.

241. When the epoch of any one passage of a planet through its perihelion, or of the sun or moon through its perigee, has been found, we may, by means of the anomalistic revolution, deduce from it the epoch of every other passage.

242. The length of the anomalistic year exceeds that of the sidereal year by 4 m. 43.9 s.

243. From the anomalistic revolution, and the epoch of the last passage through the perihelion or perigee (as the case may be), we may derive the mean anomaly for any given time. Let T = the anomalistic revolution, *t* = the time that has elapsed since the last passage through the perihelion or perigee, and A = the mean anomaly: then,

$$T : 360^\circ :: t : A = 360^\circ \frac{t}{T} \dots (60).$$

244. The place of a body in its elliptical orbit is ascertained

by finding its true anomaly. The problem which has for its object the determination of the true anomaly from the mean, was first resolved by Kepler, and is called *Kepler's Problem*. The solution of it may be found in the Appendix. Another and more convenient method of obtaining the true anomaly, is to compute the equation of the centre from the mean anomaly, and add it to the mean anomaly, or subtract it from it, according to the position of the body in its orbit (Art. 185).

### *Heliocentric Place of a Planet.*

245. The place of a planet in the plane of its orbit is designated by its orbit longitude, and radius vector. To find the orbit longitude we have the equation,

$$\text{long.} = \text{long. of perihelion} + \text{true anomaly.}$$

The orbit longitude may also be deduced from the mean longitude, by adding or subtracting the equation of the centre.

The radius vector results from the polar equation of the elliptic orbit (Art. 227), viz :

$$V = \frac{a(1 - e^2)}{1 + e \cos x} \dots (61).$$

in which  $x$  denotes the true anomaly,  $e$  the eccentricity, and  $a$  the semi-major axis.

246. Now to find the heliocentric longitude and latitude, which ascertain the position of the planet with respect to the ecliptic, the triangle  $N P p$  (Fig. 41) gives,

$$\sin P p = \sin N P \sin P N p ;$$

or,  $\sin \text{lat.} = \sin (\text{orbit long.} - \text{long. of node}) \times \sin (\text{inclin.}) \dots (62);$   
and

$\cos P N p = \tan g N p \cot N P$ , or  $\tan g N p = \tan g N P \cos P N p$  ;  
or,

$\tan g (\text{long.} - \text{long. of node}) = \tan g (\text{orbit long.} - \text{long. of node})$   
 $\times \cos (\text{inclination}) \dots (63).$

### *Geocentric Place of a Planet.*

247. *From the heliocentric longitude and latitude and the radius vector of a planet, to find the geocentric longitude and latitude.* Let  $S$  (Fig. 41) be the sun,  $E$  the earth,  $P$  the planet,  $\pi$  its reduced place, and  $V$  the vernal equinox. Denote the heliocentric longitude  $V S \pi$  by  $L$ , the heliocentric latitude  $P S \pi$  by  $l$ , and the radius vector  $S P$  by  $v$  ; and denote the geocentric



longitude by  $G$ , and the geocentric latitude by  $\lambda$ . Also let  $E = S E \pi$  the elongation;  $C = E S \pi$  the commutation;  $A = S \pi E$  the annual parallax; and  $r = S E$  the radius vector of the earth. Now,

$$V E \pi = S E \pi + V E S,$$

or,  $G = E + \text{long. of sun.}$

This equation will make known the geocentric longitude, when the value of  $E$  is found. In the triangle  $S E \pi$ , the side  $S \pi = S P \cos P S \pi = v \cos l$ , and is therefore known, the side  $E S$  is given by the elliptical theory (Art. 245), and the angle  $C$  may be derived from the following equation:  $C = V S E - V S \pi = \text{long. of earth} - \text{long. of planet}$ : and to find  $E$  we have, by Trigonometry,

$$E S + S \pi : E S - S \pi :: \tan \frac{1}{2}(E \pi S + S E \pi) : \tan \frac{1}{2}(E \pi S - S E \pi),$$

or,  $r + v \cos l : r - v \cos l :: \tan \frac{1}{2}(A + E) : \tan \frac{1}{2}(A - E)$ ;

$$\text{whence, } \tan \frac{1}{2}(A - E) = \frac{r - v \cos l}{r + v \cos l} \tan \frac{1}{2}(A + E);$$

$$= \frac{1 - \frac{v \cos l}{r}}{1 + \frac{v \cos l}{r}} \tan \frac{1}{2}(A + E).$$

Let  $\tan \theta = \frac{v \cos l}{r}$ . Then,

$$\tan \frac{1}{2}(A - E) = \frac{1 - \theta}{1 + \theta} \tan \frac{1}{2}(A + E);$$

$$\text{or, } \tan \frac{1}{2}(A - E) = \tan (45^\circ - \theta) \tan \frac{1}{2}(A + E) \dots (64).$$

But,  $A + E = 180^\circ - C$ , and  $E = \frac{1}{2}(A + E) - \frac{1}{2}(A - E)$ .

Next, to find the *geocentric latitude*.

$$S \pi \tan l = P \pi = E \pi \tan \lambda,$$

$$\text{whence, } \frac{S \pi}{E \pi} = \frac{\tan \lambda}{\tan l}$$

$$\text{but, } S \pi : E \pi :: \sin E : \sin C, \text{ or } \frac{S \pi}{E \pi} = \frac{\sin E}{\sin C},$$

$$\text{and therefore, } \frac{\sin E}{\sin C} = \frac{\tan \lambda}{\tan l};$$

$$\text{or, } \tan \lambda = \frac{\sin E \tan l}{\sin C} \dots (65).$$

248. When a planet is in conjunction or opposition, the sines of the angles of elongation and commutation are each nothing. In these cases, then, the geocentric latitude cannot be found by the preceding formula, it may however be easily determined in a different manner. Suppose the planet to be in conjunction at P (Fig. 42); then,

$$\text{tang } \lambda = \frac{P \pi}{E \pi} = \frac{P \pi}{E S + S \pi};$$

but the triangle S P  $\pi$  gives,

$$P \pi = v \sin l, \text{ and } S \pi = v \cos l; \text{ and } ES = r;$$

hence, 
$$\text{tang } \lambda = \frac{v \sin l}{r + v \cos l} \dots (66).^*$$

249. To find the distance of the planet from the earth, represent the distance by D; then, from the triangles S P  $\pi$  and E P  $\pi$  (Fig. 41), we have,

$$P \pi = E P \sin P E \pi = D \sin \lambda,$$

and

$$P \pi = S P \sin P S \pi = v \sin l;$$

whence,

$$D = \frac{v \sin l}{\sin \lambda} \dots (67).$$

250. The distance of a planet being known, its horizontal parallax may be computed from the equation

$$\sin H = \frac{R}{D} \dots (68). \quad (\text{Art. 98}).$$

### *Places of the Sun and Moon.*

251. The place of the sun, as seen from the earth, may be easily deduced from the heliocentric place of the earth; for, the longitude of the sun is equal to the heliocentric longitude of the earth plus  $180^\circ$ , and the radius vector of the earth's orbit is the same as the distance of the sun from the earth. But it is more convenient to regard the sun as describing an orbit around the earth, and to compute its true anomaly, (Art. 244), and thence the longitude and radius vector by the equation

$$\text{long.} = \text{true anomaly} + \text{long. of perigee},$$

and the polar equation of the orbit.

252. The orbit longitude and the radius vector of the moon

---

\* For opposition and inferior conjunction, the sign of  $\cos l$  must be changed.

are found by the same process as the longitude and radius vector of the sun. The orbit longitude being known, the ecliptic longitude and the latitude may be determined by a process precisely similar to that by which the heliocentric longitude and latitude of a planet are found (Art. 245).

*Verification of Kepler's Laws.*

253. If Kepler's first two laws be true, then the geocentric places of the planets, computed by the process that we have described (Art. 246), which is founded upon them, ought to agree with the true geocentric places as obtained for the same time by direct observation: or, the heliocentric places computed from the observed geocentric places (Art. 223), ought to agree with the same as computed by the elliptic theory (Arts. 245, 246). Now, a great number of comparisons have been made between the observed and computed places, and in every instance a close agreement between the two has been found to subsist. We infer, therefore, that the motions of the planets must be very nearly in conformity with these laws.

The truth of the third law has been established by a direct comparison of the mean distances of the different planets with their periodic times.

254. Kepler's laws have been verified for the sun and moon, in a similar manner.

255. The relative distances of the sun, or moon, at different times, result from observations upon the apparent diameter, upon the principle that any two distances are inversely proportional to the corresponding apparent diameters. Let  $\Delta$  = semi-diameter corresponding to the mean distance, and  $\delta$  = semi-diameter, corresponding to any distance  $D$ : then

$$\delta : \Delta :: 1 : D ; \text{ whence, } D = \frac{\Delta}{\delta} \dots (69) ;$$

an equation which, when  $\Delta$  has been found, will make known the distance corresponding to any observed semi-diameter  $\delta$ , in terms of the mean distance as a unit.

Now, to find  $\Delta$ , denote the greatest and least semi-diameters respectively by  $\delta'$ ,  $\delta''$ , and the corresponding distances by  $D'$  and  $D''$ , and we have,

$$D' = \frac{\Delta}{\delta'}, \quad D'' = \frac{\Delta}{\delta''} ;$$

and thence,  $\frac{1}{2} (D' + D'') = \frac{1}{2} \left( \frac{\Delta}{\delta'} + \frac{\Delta}{\delta''} \right)$  or,  $1 = \frac{1}{2} \left( \frac{\Delta}{\delta'} + \frac{\Delta}{\delta''} \right)$  ;

whence, 
$$\Delta = \frac{2 \delta' \delta''}{\delta' + \delta''} . . . (70).$$

256. The distance of the sun or moon in terms of the mean distance as a unit, may be found in a similar manner ; but it may be had more accurately by means of a principle which has been discovered from observation, namely, that the distance is inversely proportional to the square root of the daily angular motion.

## CHAPTER X.

OF THE INEQUALITIES OF THE MOTIONS OF THE PLANETS  
AND OF THE MOON ; AND OF THE CONSTRUCTION OF TA-  
BLES FOR FINDING THE PLACES OF THESE BODIES.

257. It is a general law of nature, discovered by Sir Isaac Newton, that bodies tend, or *gravitate* towards each other, with a force directly proportional to their masses and inversely proportional to the square of their distance. The force which causes one body to gravitate towards another, is supposed to arise from a mutual attraction existing between the particles of the two bodies, and is hence called the *Attraction of Gravitation*. This force of attraction, common to all the bodies of the Solar System, is the general physical cause of their motions. The sun's attraction retains the planets in their orbits, and the planets by their mutual attractions slightly alter each other's motions. The reasoning by which *Newton's Theory of Universal Gravitation* is established, appertains to Physical Astronomy, and will be presented in another part of the work.

258. If a planet were acted on by no other force than the attraction of the sun, it is proved that its orbit would be accu-

rately an ellipse, and that the areas described by its radius vector in equal times, would be precisely equal. But, it is in reality attracted by the other planets, as well as the sun, and therefore its actual motions cannot be in strict conformity with the laws of Kepler. In fact, if we descend to great accuracy, the agreement between the observed and computed places noticed in Art. 253, is found not to be exact. The deviations from the elliptic motion which are produced by the attractions of the planets, are called *Perturbations*, or, in Plane Astronomy, *Inequalities*. Although, as we have just seen, the fact of the existence of inequalities in the motions of the planets is discoverable from observation, their laws cannot be determined without the aid of theory.

259. In treating of the perturbations in the motions of one planet, resulting from the attractions of another, the attracting planet is called the *Disturbing Body*, and the force which produces the perturbations the *Disturbing Force*. To find the disturbing force, let P (Fig. 46) be the planet, S the sun, and M the disturbing body; and let PD represent the attraction of M for the planet. Decompose PD into two forces, PE and PF, one of which, PE, is equal and parallel to SG, the attraction of M for the sun; the other, PF, will be known in position and intensity. The two forces, PE and SG, being equal and parallel, they cannot alter the relative motion of the sun and planet, and accordingly may be left out of account: there remains, therefore, the component PF, which will be wholly effectual in disturbing this motion. This, then, is the disturbing force.

It happens in the case of each planet, that the distances of some of the other planets are so great, that their disturbing forces are insensible. The attractions of these bodies for the sun and planet are sensibly equal and parallel. Owing to the great distance of the planets from each other, and the smallness of their mass compared with that of the sun, the disturbing force is in every instance very minute in comparison with the sun's attraction.

260. It is plain that the disturbing force will, in general, be obliquely inclined to the perpendicular to the plane of the orbit, PK; the tangent to the orbit, PT; and the radius vector, PS; and may, therefore, be decomposed into forces



acting along these lines. The component along the perpendicular will alter the latitude, and the two others both the longitude and radius vector; that along the tangent by changing the velocity of the planet; and that along the radius vector by changing the gravity towards the sun. It appears, therefore, that the disturbing force produces at the same time perturbations or *inequalities of longitude, of latitude, and of radius vector*.

261. Let us now consider how these inequalities may be determined. And, in the first place, the inequalities produced by each disturbing body may be separately investigated upon mechanical principles, as if the other bodies did not exist, for the reason that the effect of each disturbing body is sensibly the same that it would be if the other bodies did not act. That this is very nearly, if not quite true, may be at once inferred from the minuteness of the whole disturbance produced by the joint action of all the disturbing forces of the system. The problem which has for its object the determination of the inequalities in the motions of one body, in its revolution around a second, produced by the attraction of a third, is called the *Problem of the Three Bodies*. If, in the case of any one planet, this problem be resolved for each of the other bodies of the system which occasion sensible perturbations, all the inequalities to which the motion of the planet is subject will become known.

262. The general solution of the problem of the three bodies, that is for any mass and distance of the disturbing body, or any intensity of the disturbing force, cannot be effected in the existing state of the mathematical sciences. But the problem has been resolved for the case that presents itself in nature, in which the disturbing force is very minute in comparison with the central attraction. The results obtained by the analysis, are certain analytical expressions for the perturbations in longitude, latitude, and radius vector, involving variables and constants.

263. The general expression for the whole perturbation in longitude, due to the action of any one disturbing body, is

$$C \sin (P' - P) + C' \sin 2 (P' - P) + C'' \sin 3 (P' - P) + \&c. (71),$$

in which  $C$ ,  $C'$ , &c. are constants,  $P$  the heliocentric longitude of the body disturbed, and  $P'$  that of the disturbing body. The number of terms is, strictly speaking, indefinite, but they form a

decreasing series ; and the value of the first term never amounts to more than a few seconds ; so that only a small number of the first terms (which will be different in different cases) need to be used.

264. The constants  $C$ ,  $C'$ , &c. are to be determined from observation ; they may, however, be determined in the case of some of the planets from theory alone. The process of finding them from observation is as follows : Suppose that the earth is the body whose perturbations are under consideration, and let  $D$  denote the perturbation in longitude, produced by the joint action of all the disturbing forces. Then, supposing, for the sake of simplicity, that the expression for the perturbation due to each disturbing body, consists of but two terms, we have,

$$D = C \sin (P' - P) + C' \sin 2 (P' - P) + c \sin (P'' - P) + c' \sin 2 (P'' - P) + \&c. \dots (72).$$

Find, by observation, the heliocentric longitude of the earth, and take the difference between this and the longitude as computed for the same time by the elliptical theory. This difference will be the value of  $D$  at the time of the observation.  $P$ ,  $P'$ ,  $P''$ , &c. the heliocentric longitudes of the earth and of the disturbing bodies, and consequently  $P' - P$ ,  $P'' - P$ , &c. are given by the elliptical theory. Thus, in the above equation all will be known but  $C$ ,  $C'$ ,  $c$ ,  $c'$ , &c. By repetitions of this process, as many equations may be obtained as there are constants to be determined, and from these the values of the constants may be computed. It is usual, however, to obtain a much greater number of equations than there are constants ; as, by combining them according to certain rules, much more exact values of the constants may be derived.

265. In the expression,

$$C \sin (P' - P) + C' \sin 2 (P' - P) + \&c.,$$

for the perturbation in longitude, due to the action of a disturbing body, each term,  $C \sin (P' - P)$ ,  $C' \sin 2 (P' - P)$ , &c., is technically termed an *Equation*, and is considered as representing a specific inequality. The angle  $P' - P$ , or  $2 (P' - P)$ , or other multiple of  $P' - P$ , the sine of which enters into the equation of an inequality, is called the *Argument* of the inequality ; and the constant is called the *Coefficient* of the inequality. As

the greatest value of the sine of the argument is unity, the coefficient is equal to the greatest value of the inequality.

266. The coefficient being known, the value of the inequality at any particular time will become known, if that of the argument be found. Now, the argument is the difference between the longitudes of the disturbing body and disturbed body, or some multiple of this difference, and may be found by the elliptical theory. In practice, the mean longitudes may be taken, without material error, in place of the true, and these are easily deduced from the mean longitudes at a given epoch, by means of the mean motions in longitude of the two bodies. When the values of all the inequalities in longitude have been separately determined, by taking their algebraic sum, we shall have the *correction* to be applied to the elliptic longitude, in order to find the exact longitude.

267. The general expression for the total perturbation of radius vector, due to the action of one body, is

$$C \cos (P' - P) + C' \cos 2 (P' - P) + C'' \cos 3 (P' - P) + \&c. (73).$$

As in the expression for the perturbation of longitude, each term is called an equation, and represents a distinct inequality, the constant being the coefficient, and the variable angle, the cosine of which enters into the equation, the argument of the inequality. The amounts of the different inequalities, at an assumed time, are computed after the same manner as those of the inequalities of longitude, and being added together with their algebraical signs, will give the correction to be applied to the elliptic radius vector.

268. The perturbation in latitude is very minute. The inequalities of latitude, as of longitude and radius vector, are represented by equations, composed of a constant coefficient and the sine or cosine of a variable argument, or of the form  $C \sin A$  or  $C \cos A$ .

269. The arguments of the inequalities we have been considering, are angles depending upon the configurations of the disturbing and disturbed planets with respect to each other and the sun, and also, in some cases, with respect to the nodes and perihelia of their orbits. Whenever these configurations become the same, as they will periodically, the arguments, and therefore

the inequalities themselves, will have the same value. It follows, therefore, that the inequalities in question are *periodic*.

The interval of time in which an inequality passes through all its gradations of positive and negative value, is called the *Period* of the inequality. It is manifestly equal to the interval of time employed by the argument in increasing from zero to  $360^\circ$ ; for, in this interval  $\sin A$  or  $\cos A$  takes all its values, both positive and negative, and at the expiration of it recovers the same value again.

270. It has been stated, that the elements of the elliptic orbits of the planets are, for the most part, subject to a slow variation from century to century. Investigations in Physical Astronomy have established that the variations of the elements are due to the action of the disturbing forces of the planets, and that they are not progressive (except in the cases of the longitude of the node and the longitude of the perihelion), but are really periodic inequalities, whose periods comprise many centuries. From the great lengths of their periods these inequalities are termed *Secular Inequalities*, in order to distinguish them from the inequalities of the elliptic motion, denominated *Periodic Inequalities*, the periods of which are comparatively short.

Physical Astronomy furnishes expressions called *Secular Equations*, which give the value of an element at any assumed time.

271. The inequalities of the moon's motions arise from the disturbing action of the sun. The attractions of the planets for the moon and earth are sensibly equal and parallel. The lunar inequalities are investigated upon the same principles as the planetary, and are represented by *equations* of the same general form, that is, consisting of a constant coefficient and the sine or cosine of a variable argument. They far exceed in number and magnitude those of any single planet.

272. There are three lunar inequalities of longitude which are prominent above the rest, and were early discovered by observation.

The most considerable is called the *Evection*, and was discovered by Ptolemy in the first century of the Christian era. It has for its argument, double the angular distance of the moon

from the sun minus the mean anomaly of the moon, and amounts when greatest to  $1^{\circ} 20' 30''$ .

The second is called the *Variation*, and was discovered in the sixteenth century by Tycho Brahé. Its argument is double the angular distance of the moon from the sun, and its maximum value is  $35' 42''$ .

The third is denominated the *Annual Equation*, from the circumstance of its period being an anomalistic year. Its argument is the mean anomaly of the sun. When greatest, it amounts to  $11' 12''$ .

273. The discovery of the other lunar inequalities (with the exception of one inequality of latitude) is due to Physical Astronomy.

The whole number of lunar inequalities of longitude, according to Burekhardt, is 32.

274. To present now at one view, the entire process of finding the exact heliocentric place of a planet, or the geocentric place of the moon, at any assumed time.

1. Seek the elements of the elliptic orbit from a table of elements, such as Table II or III, allowing for the proportional part of the secular variation, or (more exactly) obtain them from their secular equations (Art. 270).

2. Compute the longitude, latitude, and radius vector, by the elliptic theory (Arts. 245, 246).

3. Compute the values of the inequalities in longitude and latitude and of radius vector by means of their equations (Art. 266), and apply them individually with their proper signs, as corrections to the elliptic values of the longitude, latitude and radius vector.

275. If we suppose the sun to be in motion, instead of the earth, its inequalities will be the same as those to which the motion of the earth is actually subject.

276. When the heliocentric place of a planet has been found, its geocentric place, if required, may be determined by the process explained in Art. 247.

### *Construction of Tables.*

277. The determination of the place of the sun or moon, or of a planet, may be greatly facilitated by the use of tables. The principles and modes of construction of tables adapted to



this purpose are nearly the same for each body. We will first explain the mode of constructing tables for facilitating the computation of the sun's longitude. We have the equation,

$$\text{True long.} = \text{mean long.} + \text{equa. of centre} + \text{inequalities} + \text{nutations.}$$

If, then, tables can be constructed that will furnish by inspection the mean longitude, the equation of the centre, the amounts of the various inequalities in longitude, and the nutation in longitude, at any assumed time, we may easily find the true longitude at the same time.

278. 1. *For the mean longitude.* The sun's mean motion in longitude in a mean tropical year, is  $360^{\circ}$ . From this we may find by proportion, the mean motions in a common year of 365 days and a bissextile year of 366 days.

With these results and the mean longitude for the epoch of Jan. 1, 1801, we may easily derive the mean longitude at the beginning of each of the years prior and subsequent to the year 1801. The second column of Table XVIII contains the mean longitude of the sun at the beginning of each of the years inserted in the first column. The third column of this table contains the mean longitude of the perigee at the same epochs: it was constructed by means of the mean longitude of the perigee found for the beginning of the year 1800, and its mean yearly motion in longitude, which is  $61''.52^*.$

Having the sun's mean daily motion in longitude (Art. 176), we obtain by proportion the motion in any proposed number of months, days, hours, minutes, or seconds. Table XIX contains the respective amounts of the sun's motion from the commencement of the year to the close of each month; Table XX, the sun's mean motion for days from 1 to 31, and for hours from 1 to 24; and Table XXI, the same for minutes and seconds from 1 to 60. With these tables, the sun's mean motion in longitude in the interval between any given time in any year and the beginning of the year, may be had: and if this be added to the epoch for the given year, taken out from Table XVIII, the

---

\* The quantities in Table XVIII are called *Epochs*. The *Epoch* of a quantity is its value at some chosen epoch.

result will be the mean longitude at any given time. (See Problem IX.)

279. Tables XIX and XX also contain the motions of the sun's perigee, from which and the epoch given by Table XVIII, results the longitude of the perigee at any proposed time. The longitude of the perigee is given in the Solar Tables, for the purpose of making known the mean anomaly, the mean anomaly being equal to the mean longitude minus the longitude of the perigee.

280. 2. *For the equation of the centre.* To find the equation of the centre of an orbit we have the following equation :

$$\text{Equa. of centre} = A \sin \theta + B \sin 2 \theta + C \sin 3 \theta + \&c. ;$$

in which A, B, C, &c. are constants that rapidly decrease in value, and which may be determined for any particular orbit, and  $\theta$  the mean anomaly. Now, by giving to the mean anomaly  $\theta$  in this equation, a series of values increasing by small equal differences (of  $1^\circ$ , for instance,) from zero to  $360^\circ$ , and computing the corresponding values of the equation of the centre; then registering in a column the different values assigned to  $\theta$ , and in another column to the right of this, the computed values of the equation of the centre, we shall obtain a table which will give on inspection the equation of the centre corresponding to any particular mean anomaly. In this manner was constructed Table XXV. In this table, however, for the sake of compactness, the values of the equation, instead of being registered in one column, are put in as many different columns as there may be different numbers of signs in the value of the mean anomaly; each column answering to the particular number of signs placed at the head of it.

If the equation of the centre at an assumed time be required, find the mean anomaly by the tables (Art. 279), and with the value found for it take out the equation of the centre from Table XXV.

The given quantity with which a quantity is taken from a table, is called the *Argument*. Accordingly the mean anomaly is the argument of the equation of the centre in Table XXV.

281. 3. *For the inequalities.* The equations of the inequalities, as we have already stated, are of the form  $C \sin A$ , the

argument A being the difference between the longitude of the disturbing planet and that of the earth, or some multiple of this difference. With the equations of the inequalities, a table of each inequality may be constructed, upon the same principles as Table XXV. But, as the expression for the whole perturbation in longitude (Art. 263), produced by any one planet, involves only two variables, the longitude of the earth and the longitude of the planet, it is thought to be more convenient to have a table of *double entry*, which will give the amount of the perturbation by means of the two variables as arguments. Such a table may be constructed, by assigning to the longitude of the earth and the longitude of the disturbing planet a series of values increasing by a common difference, and computing with each set of the values of these quantities, the corresponding amount of the perturbation.

In connection with the tables of the perturbations, we must have tables that make known the values of the arguments at any given time. Now, the mean longitude of the sun may be found by the solar tables (Art. 278), and thence the mean heliocentric longitude of the earth by subtracting  $180^\circ$ ; and the mean longitude of the disturbing planet may be had from similar tables. The columns of Table XVIII, marked I, II, III, IV, V, VI, VII, contain the arguments of all the perturbations, for the beginning of each of the years registered in the first column, expressed in thousandth parts of a circle. Tables XIX and XX contain the variations of the arguments for months and hours. Their variations for minutes and seconds are too small to be taken into account. With these tables and Table XVIII, the values of the arguments at any given time may be found, and by means of the arguments the perturbations may be taken from Tables XXVIII, XXIX, XXX, XXXI, XXXII, and XXXIII.

282. 4. *For the nutation.* The formula for the lunar nutation in longitude, is  $17''.3 \sin N$ , —  $0''.2 \sin 2 N$ , in which N denotes the supplement of the longitude of the moon's ascending node. With this formula the second column of Table XXVII was constructed. The value of N, in thousandth parts of a circle, results from Tables XVIII, XIX, and XX. The solar nutation is also given by Table XXVII.

283. Tables may also be constructed that will facilitate the computation of the radius vector. We have,

True rad. vector = elliptic rad. vector + perturbations.

A table of the elliptic radius vector may be formed by means of the polar equation of the orbit, and tables of the perturbations from their analytical expressions (Art. 265). The tables of the perturbations will have the same arguments as the tables of the perturbations of longitude.

284. Lunar and planetary tables are constructed upon the same principles as the solar tables we have been describing, which serve to make known the orbit longitude and radius vector. But other tables are necessary in the case of these bodies, for the computation of the ecliptic longitude and the latitude.

285. The difference between the orbit longitude and the ecliptic longitude, is called the *Reduction* to the ecliptic. A formula for the reduction has been investigated, in which the variable is the difference between the orbit longitude and the longitude of the node. If this formula be reduced to a table, by taking the reduction from the table and adding it to the orbit longitude we shall have the ecliptic longitude. Table LIII is a table of reduction, for the moon.

286. *For the latitude*, we have the equation

True lat. = lat. in orbit + perturbations.

We have already seen (Art. 246), that

$\sin (\text{lat. in orbit}) = \sin (\text{orbit long.} - \text{long. of node}). \sin \text{inclina.}$

A table constructed from this formula, will have for its argument the orbit longitude minus the longitude of the node, which is also the argument of reduction. (See Table LV).

The tables of the perturbations in latitude are constructed upon the same principles as the tables of the perturbations in longitude and radius vector.

287. A table exhibiting the longitude and latitude, right ascension and declination, distance, parallax, semi-diameter, &c., of the sun or other body, at stated periods of time, as at noon of each day throughout the year, is called an *Ephemeris* of the body. An ephemeris of the sun, of the moon, and of each of the planets, is

published for each year in advance, in the English Nautical Almanac, and in the *Connaissance des Temps*.

---

## CHAPTER XI.

### OF THE MOTIONS OF THE COMETS.

288. When first seen, a comet is ordinarily at some distance from the sun, and moving towards him. After this it continues to approach the sun for a certain time, and then recedes from him to a greater or less distance, and finally disappears. In many instances comets have come so near the sun as to be for a time lost in his beams.

289. Comets resemble the planets in their changes of apparent place amongst the fixed stars, but they differ from them in never having been observed to perform an entire circuit of the heavens. Their apparent motions are also more irregular than those of the planets, and they are confined to no particular region of the heavens, but traverse indifferently every part.

290. Sir Isaac Newton, from observations that had been made upon the remarkable comet of 1680, ascertained that this comet described a parabolic orbit, having the sun at its focus, or an elliptic orbit of so great an eccentricity as to be undistinguishable from a parabola, and that its radius vector described equal areas in equal times. Since then, the orbits of about 140 comets have been computed, and found to be, with a few exceptions, of a parabolic form, or sensibly so.

291. It was demonstrated by Newton, on the theory of gravitation, that a body projected in space may describe about the sun as a focus, either one of the conic sections, and that the form of the orbit will depend upon the projectile velocity alone. With one particular velocity the orbit will be a parabola; with any less velocity, it will be an ellipse or circle; and with any



greater velocity, it will be a hyperbola. Now, as there is but one velocity from which a parabolic orbit will result, and as any comet, which may have originally moved in a hyperbola, must have passed its perihelion, and receded beyond the limits of the solar system, it may be inferred, with great probability, that the orbits of the comets whose observed courses are not distinguishable from parabolic arcs, are in fact ellipses of great eccentricity. This is the theory of the cometary motions proposed by Newton.

The orbits of some of the comets are known from observation to be very eccentric ellipses.

292. The *elements of a comet's orbit* are the longitude of the ascending node, the inclination of the orbit, the longitude of the perihelion, the perihelion distance, and the epoch of the perihelion passage. These make known the position and dimensions of the orbit, on the supposition that it is a parabola, and thus appertain only to the motions of the comet for the period during which it is visible.

293. Assuming that the radius vector of a comet describes areas proportional to the times, the elements of its orbit may be computed from three observed geocentric places. The problem is, however, one of considerable difficulty.

294. Astronomers do not seek to deduce from the observations made during one appearance of a comet, its entire elliptic orbit. It is impossible, from such observations, to compute the major axis of its orbit and its period with any accuracy, inasmuch as in the interval during which they are made the comet describes but a small portion of its entire orbit.

The only mode of obtaining the period of a comet's revolution, is by noting the time of its return to the perihelion of its orbit. A comet cannot be recognized at a second appearance by its aspect, for, this is liable to great alterations. It may, however, be identified by means of the elements of its orbit, as it is extremely improbable that the elements of the orbits of two different comets will agree throughout. This method of identifying a comet on a second appearance may sometimes fail of application, inasmuch as the orbit of a comet may experience great alterations from the attractions of the planets.

295. Owing to the great lengths of the periods of most of the

comets, and the comparatively short interval during which their motions have been carefully observed, there are but *three* comets, the periods and entire orbits of which have been determined. These are denominated *Encke's Comet*, *Biela's Comet*, and *Halley's Comet*. The two former are small telescopic objects, but the latter, when near its perihelion, is distinctly visible to the naked eye.

296. Encke's Comet is so called from Professor Encke, of Berlin, who first ascertained its periodical return. It accomplishes its revolution in the short period of 1207 days, or about  $3\frac{1}{3}$  years, and moves in an orbit inclined under a small angle to the plane of the ecliptic, and whose perihelion is at the distance of the planet Mercury, and aphelion nearly at the distance of Jupiter. This discovery was made on the occasion of its fourth recorded appearance, in 1819. Since then, it has returned several times to its perihelion, and in every instance very nearly as predicted. Its last return took place in 1835, its next will happen in the fall of the present year (1838).

297. Biela's Comet, as it is called, was discovered by M. Biela, of Johannisberg, on the 27th February, 1836. Its period is about  $6\frac{3}{4}$  years. Its orbit is inclined under a small angle to the plane of the ecliptic, and lies mostly between the orbits of the earth and of Jupiter. By a remarkable coincidence, the orbit of this comet very nearly intersects the orbit of the earth. Its last appearance took place, according to prediction, in 1832; the next will be in 1838.

298. Halley's Comet is so called from Sir Edmund Halley, who discovered its period, and correctly predicted its return. From a comparison of the elements of the orbits described by the comets of 1531, 1607, and 1682, he concluded that the same comet had made its appearance in these several years, and predicted that it would again return to its perihelion in the year 1759, as it actually did. Assuming the earth's mean distance from the sun to be unity, the perihelion distance of this comet is 0.58, and aphelion distance 35.32. Accordingly, it approaches the sun to within one half the distance of the earth, and recedes from him far beyond the orbit of Uranus. Its period is about 76 years, but is liable to a variation of a year or more, from the effect of the attractions of the planets. The last perihelion passage took place

on the 16th of November, 1835, within a few days of the predicted time. The next will occur about the year 1912.

299. Of the comets which have been observed, some have a direct and others a retrograde motion. The perihelia of their orbits, for the most part, lie within the orbit of the earth, and the aphelia far without the orbit of Uranus. Many of them come into close proximity to the sun. The great comet of 1680, according to the computation of Newton, came 166 times nearer the sun than the earth is. The planes of the orbits are inclined under every variety of angle to the plane of the ecliptic.

300. The motions of the comets are liable to great derangements from the attractions of the planets. As their orbits cross the orbits of the planets, they may come into proximity with these bodies, and be strongly attracted by them. The comet of 1770, commonly called Lexel's Comet, offers a striking example of the disturbances to which the cometary motions are exposed. From observations made upon this comet in the year 1770, Lexel made out that its period was  $5\frac{1}{2}$  years; still it has not since been seen. According to Burckhardt, this comet, previous to the year 1767, moved in an orbit which answered to a period of 50 years, and never approached near enough to the earth and sun to become visible. Early in the year 1767 it came so near the planet Jupiter, that his attraction changed its orbit to one of  $5\frac{1}{2}$  years. It thus became visible in 1770, and would have again been seen in 1776, had it not been so situated with regard to the earth as to be entirely hid by the sun's rays. In the year 1779 it again met with Jupiter, and its orbit was so much enlarged by his attraction, that it now employs twenty years in completing a revolution, and no longer comes near enough to the earth to be visible.

## CHAPTER XII.

## OF THE MOTIONS OF THE SATELLITES.

301. As it has already been remarked, the planets which have satellites are Jupiter, Saturn, and Uranus. The number of Jupiter's satellites is four ; of Saturn's, seven ; of Uranus', six.

302. The satellites of Jupiter are perceptible with telescopes of moderate power. It is found by repeated observations, that they are continually changing their positions with respect to one another and the planet, being sometimes all to the right of the planet, and sometimes all to the left of it, but more frequently some on each side. They are distinguished from each other by the distance to which they recede from the planet, that which recedes to the least distance being called the *First* Satellite, that which recedes to the next greater distance the *Second*, and so on.

The satellites of Jupiter were discovered by Galileo, in the year 1610.

303. The satellites of Saturn and of Uranus cannot be seen except through excellent telescopes. They experience changes of apparent position, similar to those of Jupiter's satellites.

304. The apparent motion of Jupiter's satellites alternately from one side to the other of the planet, leads to the supposition that they actually revolve around the planet. This inference is confirmed by other phenomena. While a satellite is passing from the eastern to the western side of the planet, a small dark spot is frequently seen crossing the disc of the planet in the same direction : and again, while the satellite is passing from the western to the eastern side, it often disappears, and after remaining for a time invisible, re-appears at another place. These phenomena are easily explained, if we suppose that the planet and its satellites are opaque bodies, illuminated by the sun, and that the satellites revolve around the planet from west to east. On this hypothesis, the dark spot seen traversing the disc of the

planet, is the shadow cast upon it by the satellite on passing between the planet and the sun, and the disappearance of the satellite is an *eclipse*, occasioned by its entering the shadow of the planet.

As the transit of the shadow occurs during the passage of the satellite from the eastern to the western side of the planet, and the eclipse of the satellite during its passage from the western to the eastern side, the direction of the motion must be from west to east.

305. Analogous conclusions may be drawn from similar phenomena exhibited by the satellites of Saturn. The satellites of Uranus also revolve around their primary, but the direction of their motion is from east to west.

306. Let us now examine into the principal circumstances of the eclipses of Jupiter's satellites, and of the transits of their shadows across the disc of the primary. Let  $E E' E''$  (Fig. 47) represent the orbit of the earth,  $P P' P''$  the orbit of Jupiter, and  $s s' s''$  that of one of its satellites. Suppose that  $E$  is the position of the earth, and  $P$  that of the planet, and conceive two lines,  $a a'$ ,  $b b'$ , to be drawn tangent to the sun and planet: then, while the satellite is moving from  $s$  to  $s'$  it will be eclipsed, and while it is moving from  $f$  to  $f'$  its shadow will fall upon the planet. Again, if  $E c$ ,  $E c'$  represent two lines drawn from the earth tangent to the planet on either side, the satellite will, while moving from  $g$  to  $g'$ , traverse the disc of the planet, and while moving from  $h$  to  $h'$ , be behind the planet, and thus concealed from view. It will be seen on an inspection of the figure, that during the motion of the earth from  $E''$ , the position of opposition, to  $E'$ , that of conjunction, the disappearances or *immersions* of the satellite will take place on the western side of the planet; and that the *emersions*, if visible at all, can be so only when the earth is so far from opposition and conjunction that the line  $E s'$ , drawn from the earth to the point of emersion, will lie to the west of  $E c$ . It will also be seen, that during the passage of the earth from  $E'$  to  $E''$  the emersions will take place on the eastern side of the planet, and that the immersions cannot be visible, unless the line  $F s$ , drawn from the earth to the point of immersion, passes to the east of the planet. It appears from observation, that the immersion and emersion are never both visible at the



same period, except in the case of the third and fourth satellites.

If the orbits of the satellites lay in the plane of Jupiter's orbit, an eclipse of each satellite would occur every revolution, but, in point of fact, they are somewhat inclined to this plane, from which cause the fourth satellite sometimes escapes an eclipse.

307. The periods and other particulars of the motions of the satellites, result from observations upon their eclipses. The middle point of time between the satellite entering and emerging from the shadow of the primary, is the time when the satellite is in the direction, or nearly so, of a line joining the centres of the sun and primary. If the latter continued stationary, then the interval between this and the succeeding central eclipse would be the periodic time of the satellite. But, the primary planet moving in its orbit, the interval between two successive eclipses is a synodic revolution. The synodic revolution, however, being observed, and the period of the primary being known, the periodic time of the satellite may be computed.

308. The mean motions of the satellites differ but little from their true motions : and hence the forms of their orbits must be nearly circular. The orbit, however, of the third satellite of Jupiter has a small eccentricity ; that of the fourth, a larger.

309. The distances of the satellites from their primary are determined from micrometrical measurements of their apparent distances at the times of their greatest elongations.

A comparison of the mean distances of Jupiter's satellites with their periodic times, proves that Kepler's third law with respect to the planets applies also to these bodies ; or, that the squares of their sidereal revolutions are as the cubes of their mean distances from the primary.

The same law also has place with the satellites of Saturn and Uranus.

310. The computation of the place of a satellite for a given time, is effected upon similar principles with that of the place of a planet. The mutual attractions of Jupiter's satellites occasion sensible perturbations of their motions, of which account must be taken when it is desired to determine their places with accuracy.

311. Laplace has shown from the theory of gravitation, that, by reason of the mutual attractions of the first three of Jupiter's satel-

lites, their mean motions and mean longitudes are permanently connected by the following remarkable relations.

1. *The mean motion of the first satellite plus twice that of the third is equal to three times that of the second.*

2. *The mean longitude of the first satellite plus twice that of the third minus three times that of the second is equal to  $180^\circ$ .*

312. It follows from this last relation, that the longitudes of the three satellites can never be the same at the same time, and consequently that they can never be all eclipsed at once.

## CHAPTER XIII.

### ON THE MEASUREMENT OF TIME.

#### *Different Kinds of Time.*

313. In Astronomy, as we have already stated, three kinds of time are used: *Sidereal*, *True or Apparent Solar*, and *Mean Solar Time*. Sidereal time being measured by the diurnal motion of the vernal equinox, true or apparent solar time by that of the sun, and mean solar time by that of an imaginary sun called the *Mean sun*, conceived to move uniformly in the equator with the real sun's mean motion in right ascension or longitude.

314. The sidereal day and the mean solar day are each of uniform duration, but the length of the true solar day is variable, as we will now proceed to show.

The sun's daily motion in right ascension, expressed in time, is equal to the excess of the solar over the sidereal day. Now this arc, and therefore the true solar day, varies from two causes, viz :

1. *The inequality of the sun's daily motion in longitude.*
2. *The obliquity of the ecliptic to the equator.*

If the ecliptic were coincident with the equator, the daily arc of right ascension would be equal to the daily arc of longitude,

and therefore would vary between the limits  $57' 11''$  and  $61' 10''$ , which would answer respectively to the apogee and perigee. But, owing to the obliquity of the ecliptic, the inclination of the daily arc of longitude to the equator is subject to a variation; and this, it is plain, will be attended with a variation in the daily arc of right ascension. The tendency of this cause is obviously to make the daily arc of right ascension least at the equinoxes, where the obliquity of the arc of longitude is greatest, and greatest at the solstices, where the obliquity is least.

315. As the length of the apparent solar day is variable, it cannot conveniently be employed for the expression of intervals of time; moreover, a clock, to keep apparent solar time, requires to be frequently adjusted. These inconveniences attending the use of apparent solar time, led astronomers to devise a new method of measuring time, to which they gave the name of mean solar time. By conceiving an imaginary sun to move uniformly in the equator with the real sun's mean motion, a day was obtained, of which the length is invariable, and equal to the mean length of all the apparent solar days in a tropical year; and by supposing the right ascension of this fictitious sun to be, at the instant of the sun's arrival at the perigee of his orbit, equal to the sun's true longitude, and consequently at all times equal to the sun's mean longitude, the time deduced from its position with respect to the meridian, was made to correspond very nearly with apparent solar time.

316. To find the excess of the mean solar day over the sidereal day, we have the proportion

$$360^\circ : 24 \text{ sid. hours} :: 59' 8''.33 : x = 3\text{m. } 56.555\text{s.}$$

A mean solar day, comprising 24 mean solar hours, is, therefore, 24h. 3m. 56.555s. of sidereal time. Hence, a clock regulated to sidereal time will gain 3m. 56.555s. in a mean solar day.

317. In order to find the expression for the sidereal day in mean solar time, we must use the proportion,

$$24\text{h. } 3\text{m. } 56.555\text{s.} : 24\text{h.} :: 24\text{h.} : x = 23\text{h. } 56\text{m. } 4.092\text{s.}$$

The difference between this and 24 hours is 3m. 55.908s.; and therefore, a mean solar clock will lose with respect to a sidereal

clock, or with respect to the fixed stars, 3m. 55.908s. in a sidereal day, and proportionally in other intervals. This is called the *daily acceleration* of the fixed stars.

318. To express any given period of sidereal time in mean solar time, we must subtract for each hour  $\frac{3\text{m. } 55.91\text{s.}}{24} = 9.83\text{s.}$ , and for minutes and seconds in the same proportion. And, on the other hand, to express any given period of mean solar time in sidereal time, we must add for each hour  $\frac{3\text{m. } 56.55\text{s.}}{24} = 9.86\text{s.}$ , and for minutes and seconds in the same proportion.

319. It is the practice of astronomers to adjust the sidereal clock to the motions of the *true* instead of the mean equinox. The inequality of the diurnal motion of this point is too small to occasion any practical inconvenience. Sidereal time, as determined by the position of the true equinox, will not deviate from the same as indicated by the position of the mean equinox, more than 2.3s. in 19 years.

320. Another species of time, called *Mean Equinoctial Time*, has recently been introduced to some extent into astronomical calculations. Mean equinoctial time signifies the mean time elapsed since the instant of the Mean Vernal Equinox. Its use is to afford an uniform date, which shall be independent of the different meridians, and of all inequalities in the sun's motion, and shall thus save the necessity, when speaking of the time of any event's happening, of mentioning at the same time the place where it was observed or computed. Thus, it is the same thing to say that a comet passed its perihelion on January 5th, 1837, at 5h. 47m. 0.0s., mean time at Greenwich; at 5h. 56m. 21.5s., mean time at Paris; or at 1836y. 289d. 6h. 16m. 40.96s., equinoctial time; but the former dates make the localities of Greenwich and Paris enter as elements of the expression; whereas the latter expresses the period elapsed since an epoch common to all the world, and identifiable independently of all localities. By this means all ambiguities in the reckoning of time are supposed to be avoided.

*Conversion of One Species of Time into Another.*

321. The difference between the apparent and mean time is called the *Equation of Time*. The equation of time, when

known, serves for the conversion of mean time into apparent, and the reverse.

322. *To find the equation of time.* The hour angle of the sun (p. 11, def. 16) varies at the rate of  $360^\circ$  in a solar day, or  $15^\circ$  per solar hour. If, therefore, its value at any moment be divided by 15, the quotient will be the apparent time at that moment. In like manner, the hour angle of the mean sun, divided by 15, gives the mean time. Now, let the circle V S D (Fig. 48) represent the equator, V the vernal equinox, M the point of the equator, which is on the meridian, and V S the right ascension of the sun, and we shall have,

$$\text{appar. time} = \frac{M S}{15} = \frac{V M - V S}{15}.$$

Again, if we suppose the circle V M D to represent the mean equator, V' the mean equinox, and S' the position of the mean sun, (V S' being equal to the mean longitude of the sun,) we shall have,

$$\text{mean time} = \frac{M S'}{15} = \frac{V M - V S'}{15} = \frac{V M - (V' S' + V V')}{15};$$

thus,

$$\text{equa. of time} = \text{mean time} - \text{ap. time} = \frac{V S - (V' S' + V V')}{15} (74);$$

or, *the equation of time is equal to the difference between the sun's true right ascension and mean longitude, corrected by the equation of the equinoxes in right ascension, and converted into time.*

A formula has been investigated, and reduced to a table, which makes known the equation of time by means of the sun's mean longitude as an argument. (See Table XII.) The value of the equation of time at noon, on any day of the year, is also to be found in the ephemeris of the sun, published in the Nautical Almanac, and other works. If its value for any other time than noon be desired, it may be obtained by simple proportion.

The value of the equation of time, determined from formula (74), is to be applied with its sign to the apparent time to obtain the mean, and with the opposite sign to the mean time to obtain the apparent.

323. The equation of time is zero, or mean and true time are the same four times in the year, viz: about the 15th of April,



the 15th of June, the 1st of September, and the 24th of December. Its greatest additive value (to apparent time) is about  $14\frac{1}{2}$  minutes, and occurs about the 11th of February; and its greatest subtractive value is about  $16\frac{1}{4}$  minutes, and occurs about the 3d of November.

324. *To convert sidereal time into mean time, and vice versa.* Making use of Fig. 48 already employed, the arc  $VM$ , called the *Right Ascension of Mid-Heaven*, expressed in time, is the sidereal time;  $VS'$  is the right ascension of the mean sun, estimated from the true equinox, or the mean longitude of the sun corrected for the equation of the equinoxes in right ascension (Art. 322); and  $MS'$  expressed in time, is the mean time. Let the arcs  $VM$ ,  $MS'$  and  $VS'$ , converted into time, be denoted respectively by  $S$ ,  $M$ , and  $L$ . Now,

$$VM = MS' + VS';$$

or,  $S = M + L$  . . (75); and  $M = S - L$  . . (76).

If  $M + L$  in equation (75) exceeds 24 hours, 24 hours must be subtracted; and if  $L$  exceeds  $S$  in equation (76), 24 hours must be added to  $S$ , to render the subtraction possible.

This problem may in practice be solved most easily by means of an ephemeris of the sun, which gives the value of  $S$ , or the sidereal time at the instant of mean noon of each day, together with a table of the acceleration of sidereal on mean solar time, and the corresponding table of the retardation of mean on sidereal time.

325. The conversion of apparent time into sidereal, or sidereal time into apparent, may be effected by first obtaining the mean time, and then converting this into sidereal or apparent time, as the case may be.

#### *Determination of the Time and Regulation of Clocks by Astronomical Observations.*

326. The regulation of a clock consists in finding its *error* and its *rate*.

327. The error of a mean solar clock is most conveniently determined, from observations with a transit instrument of the time, as given by the clock, of the meridian passage of the sun's centre. The time noted will be the *clock* time at apparent noon, and the exact mean time at apparent noon may be obtained by applying to the apparent time the equation of time with its proper

sign, which may for this purpose be taken from the Nautical Almanac by simple inspection. A comparison of the clock time with the exact mean time will give the error of the clock.

328. The daily rate of a mean solar clock may be ascertained by finding as above the error at two successive apparent noons. If the two errors are the same and lie the same way, the clock goes accurately to mean solar time; if they are different, their difference or sum, according as they lie the same or opposite ways, will be the daily gain or loss, as the case may be.

329. To find the error of a sidereal clock, compute the true right ascension of some one of the fixed stars, (see Prob. XXI,) and note the time of its transit; the difference between the time observed and the right ascension in time will be the error. The error of the daily rate is determined by observing two successive transits of the same star. The variation of the time of the second transit from that of the first will be the error in question.

The error and rate may be determined more accurately from observations upon several stars, taking a mean of the individual results. Stars at a distance from the pole are to be selected, for reasons which have been already assigned.

330. In default of a transit instrument, the time may be obtained, and time-keepers regulated, by observations made out of the meridian. There are two methods by which this may be accomplished, called, respectively, the method of *Single Altitudes*, and the method of *Double Altitudes* or of *Equal Altitudes*. These we will now explain.

1. *To determine the time from a measured altitude of the sun, or of a star, its declination and also the latitude of the place being given.*

Let us first suppose that the altitude of the sun is taken; correct the measured altitude for refraction and parallax, and also, if the sextant is the instrument used, for the semi-diameter of the sun. Then, if  $Z$  (Fig. 13) represents the zenith,  $P$  the elevated pole, and  $S$  the sun; in the triangle  $ZPS$  we shall know  $ZP$  = co-latitude,  $PS$  = co-declination, and  $ZS$  = co-altitude, from which we may compute the angle  $ZPS$  ( $= P$ ), which is the angular distance of the sun from the meridian, or, if expressed in time, the time of the observation from apparent noon, by the following equations (App., Resolution of oblique-angled spherical triangles, Case I),

$$2k = ZP + PS + ZS = \text{co-lat.} + \text{co-dec.} + \text{co-alt.} \dots (77).$$

$$\sin^2 \frac{1}{2}P = \frac{\sin(k - ZP) \sin(k - PS)}{\sin ZP \sin PS} \dots (78).$$

$$\text{or, } \sin^2 \frac{1}{2}P = \frac{\sin(k - \text{co-lat.}) \sin(k - \text{co-dec.})}{\sin(\text{co-lat.}) \sin(\text{co-dec.})} \dots (79).$$

The value of  $P$  being derived from these equations and converted into time (see Prob. III), the result will be the apparent time at the instant of the observation, if it was made in the afternoon; if not, what remains after subtracting it from 24 hours will be the apparent time. The apparent time being found, the mean time may be deduced from it by applying the equation of time.

A more accurate result will be obtained if *several* altitudes be measured, the time of each measurement noted, and the mean of all the altitudes taken and regarded as corresponding to the mean of the times. The correspondence will be sufficiently exact if the measurements be all made within the space of 10 or 12 minutes, and *when the sun is near the prime vertical*. If an even number of altitudes be taken, and alternately of the upper and lower limb, the mean of the whole will give the altitude of the sun's centre, without it being necessary to know his apparent semi-diameter. In practice, the declination of the sun may be taken for the solution of this problem from an ephemeris of the sun. For this purpose the time of the observation and the longitude of the place must be approximately known.

Example. On the 1st of June, 1838, at about 10h. 45m. A. M., the altitude of the sun's lower limb was measured at New York with a sextant, and found to be  $64^\circ 55' 5''$ . What was the correct time of the observation?

Measured alt. of sun's lower limb, . . .	64° 55' 5"
Sun's semi-diam., by Conn. des Tems,	15 48
<hr/>	
Appar. alt. of sun's centre, . . .	65 10 53
Parallax in alt. (Table X), . . .	+ 4
Refraction (Table VIII), . . .	—27
<hr/>	
True alt. of sun's centre, . . .	65 10 30

N. York approx. time of observation, 10h. 45m.

Diff. of long. of Paris and N. York, 5 5

Paris, approx. time of obs. . . . 4 20 P. M.

Sun's declin. June 1st, M. noon at Paris, 22° 2' 27"

" " June 2d, " " 22 10 31

Change of declin. in 24 hours, . . . 8 4

24h. : 8' 4" :: 4h. 20m. : 1' 27".

Declin. June 1st, M. noon at Paris, . 22° 2' 27"

Change of declin. in 4h. 20m. . +1 27

Declin. at time of obs. . . . 22 3 54

90° 0' 0"

Lat. of N. York, 40 42 40

logarithms.

Co-lat. . . 49 17 20 . ar. co. sin. 0.12033

Co-dec. . . 67 56 6 . ar. co. sin. 0.03303

Co-alt. . . 24 49 30

2 ) 142 2 56

k . . 71 1 28

k — co-lat. . 21 44 8 . . . sin. 9.56858

k — co-dec. . 3 5 22 . . . sin. 8.73154

2 ) 18.45348

 $\frac{1}{2}P = 9 42 13 . . . \sin. 9.22674$ 

P = 19 24 26

4

1h. 17m. 37s. 44"

10 42 22 A. M.

Equa. of time, — 2 34

M. time of obs. 10 39 48 A. M.

In case the altitude of a star is taken, the value of  $P$  derived from formula (79), when converted into time, will express the distance in time of the star from the meridian, and being added to the right ascension of the star, if the observation be made to the westward of the meridian, or subtracted from the right ascension (increased by 24h., if necessary,) if the observation be made to the eastward, will give the *sidereal* time of the observation.

2. *To determine the time of noon from equal altitudes of the sun, the times of the observations being given.*

If the sun's declination did not change while he is above the horizon, he would have equal altitudes at equal times before and after apparent noon. Hence, if to the time of the first observation one half the interval of time between the two observations should be added, the result would be the time of noon, as shown by the clock or watch employed to note the times of the observations. The deviation from 12 o'clock would be the error of the clock with respect to apparent time. The difference between this error and the equation of time would be the error of the clock with respect to mean time.

But, as in point of fact the sun's declination is continually changing, equal altitudes will not have place precisely at equal times before and after noon, and it is therefore necessary in order to obtain an exact result, to apply a correction to the time thus obtained. This correction is called the *Equation of Equal Altitudes*. Tables have been constructed, by the aid of which the equation is easily obtained. This is at the same time a very simple and very accurate method of finding the time, and the error of a clock.

If equal altitudes of a star should be observed, it is evident that half the interval of time elapsed would give the time of the star passing the meridian, without any correction. From this the error of the clock (if keeping sidereal time) may be found, as explained in Art. 329.

### *Of the Calendar.*

331. The sun naturally regulates the beginnings, ends, and durations of the seasons; and the calendar is constructed to distribute and arrange the smaller portions of the year.



332. The calendar divides the year into 12 months, containing in all 365 days. Now, it is desirable that it should always denote the same parts of the same season by the same days of the same months, that, for instance, the summer and winter solstices, if once happening on the 21st of June and 21st of December, should ever after be reckoned to happen on the same days ; that the date of the sun's entering the equinox, the natural commencement of spring, should, if once, be always on the 20th of March. For thus the labours of agriculture, which really depend on the situation of the sun in the heavens, would be simply and truly regulated by the calendar.

This would happen, if the civil year of 365 days were equal to the astronomical ; but the latter is greater ; therefore, if the calendar should invariably distribute the year into 365 days, it would fall into this kind of confusion, that in progress of time, and successively, the vernal equinox would happen on every day of the civil year. Let us examine this more nearly.

Suppose the excess of the astronomical year above the civil to be exactly 6 hours, and, on the noon of March 20th of a certain year, the sun to be in the equinoctial point ; then, after the lapse of a civil year of 365 days, the sun would be on the meridian, but not in the equinoctial point ; it would be to the west of that point, and would have to move 6 hours in order to reach it, and to complete the astronomical or tropical year. At the completions of a second and a third civil year, the sun would be still more and more remote from the equinoctial point, and would be obliged to move, respectively, for 12 and 18 hours before he could rejoin it and complete the astronomical year.

At the completion of a fourth civil year the sun would be more distant than on the two preceding ones, from the equinoctial point. In order to rejoin it, and to complete the astronomical year, he must move for 24 hours ; that is, for *one whole day*. In other words, the astronomical year would not be completed till the beginning of the next astronomical day ; till, in civil reckoning, the *noon of March 21st*.

At the end of four more common civil years, the sun would be in the equinox on the noon of March 22d. At the end of 8 and 64 years, on March 23d and April 6th, respectively ; at the end of 736 years, the sun would be in the vernal equinox on

September 20th ; and in a period of about 1508 years, the sun would have been in every sign of the zodiac on the same day of the calendar, and in the same sign on every day.

333. If the excess of the astronomical above the civil year were really what we have supposed it to be, 6 hours, this confusion of the calendar might be most easily avoided. It would be necessary merely to make every fourth civil year to consist of 366 days ; and, for that purpose, to interpose, or to *intercalate*, a day in a month previous to March. By this *intercalation*, what would have been March 21st is called March 20th, and accordingly the sun would be still in the equinox on the same day of the month.

334. This mode of correcting the calendar was adopted by Julius Cæsar. The fourth year into which the intercalary day is introduced was called *Bissextile* ; it is now frequently called the *Leap* year. The correction is called the *Julian* correction, and the length of a mean Julian year is 365d. 6h.

*By the Julian Calendar, every year that is divisible by 4 is a leap year, and the rest common years.*

335. The astronomical year being equal to 365d. 5h. 48m. 47.6s., it is less than the mean Julian by 11m. 12.4s. or 0.007783d. The Julian correction, therefore, itself needs correction. The calendar regulated by it, would, in process of time, become erroneous, and would require *reformation*.

The intercalation of the Julian correction being too great, its effect would be to *antedate* the happening of the equinox. Thus (to return to the old illustration) the sun, at the completion of the fourth civil year, now the Bissextile, would have passed the equinoctial point by a time equal to four times 0.007783d. ; at the end of the next Bissextile, by eight times 0.007783d. ; at the end of 130 years, by about one day. In other words, the sun would have been in the equinoctial point *24 hours previously*, or on the *noon of March 19th*.

In the lapse of ages this error would continue and be increased. Its accumulation in 1300 years would amount to 10 days, and then the vernal equinox would be reckoned to happen on March 10th.

336. The error into which the calendar had fallen, and would continue to fall, was noticed by Pope Gregory in 1582. At his

time the length of the year was known to greater precision than at the time of Julius Cæsar. It was supposed equal to 365d. 5h. 49m. 16.23s. Gregory, desirous that the vernal equinox should be reckoned on or near March 21st (on which day it happened in the year 325, when the Council of Nice was held), ordered that the day succeeding the 4th of October, 1582, instead of being called the 5th, should be called the 15th; thus suppressing 10 days, which, in the interval between the years 325 and 1582, represented nearly the accumulation of error arising from the *excessive intercalation of the Julian correction*.

This act *reformed* the calendar. In order to correct it in future ages, it was prescribed that, at certain convenient periods, the intercalary day of the Julian correction should be omitted. Thus the centurial years 1700, 1800, 1900, are, according to the Julian calendar, Bissextiles, but on these it was ordered that the intercalary day *should not be inserted*; inserted again in 2000, but not inserted in 2100, 2200, 2300; and so on for succeeding centuries. *By the Gregorian calendar, then, every centurial year that is divisible by 400 is a Bissextile or Leap year, and the others common years.* For other than centurial years, the rule is the same as with the Julian calendar.

337. This is a most simple mode of regulating the calendar. It corrects the insufficiency of the Julian correction, by omitting, in the space of 400 years, 3 intercalary days. And it is easy to estimate the degree of its accuracy. For, the real error of the Julian correction is 0.007783d. in 1 year, consequently  $400 \times 0.007783d.$  or 3.1132d. in 400 years. Consequently, 0.1132d. or 2h. 43m. 0.5s. in 400 years, or 1 day in 3533 years, is the measure of the degree of inaccuracy in the Gregorian correction.

338. The Gregorian calendar was adopted immediately on its promulgation, in all Catholic countries, but in those where the Protestant religion prevailed, it did not obtain a place till some time after. In England, "the change of style," as it was called, took place after the 2d of September, 1752, eleven nominal days being then struck out; so that the last day of *Old Style* being the 2d, the first of *New Style* (the next day) was called the 14th, instead of the 3d. The same legislative enactment which established the Gregorian calendar in England, changed the time of the beginning of the year from the 25th of March to the 1st of

January. Thus the year 1752, which by the old reckoning would have commenced with the 25th of March, was made to begin with the 1st of January : so that the number of the year is, for dates falling between the 1st of January and the 25th of March, one greater by the new than by the old style. In consequence of the intercalary day omitted in the year 1800, there is now, for all dates, 12 days difference between the old and new style.

Russia is at present the only Christian country in which the Gregorian calendar is not used.

339. The calendar months consist, each of them, of 30 or 31 days, except the second month, February, which, in a common year, contains 28 days, and in a Bissextile, 29 days ; the intercalary day being added at the last of this month.

340. To find the number of days comprised in any number of civil years, multiply 365 by the number of years, and add to the product as many days as there are Bissextile years in the period.

## PART II.

### ON THE PHENOMENA RESULTING FROM THE MOTIONS OF THE HEAVENLY BODIES, AND ON THEIR APPEARANCES, DIMENSIONS, AND PHYSICAL CONSTITUTION.

---

#### CHAPTER XIV.

##### OF THE SUN AND THE PHENOMENA ATTENDING ITS APPARENT MOTIONS.

###### *Inequality of Days.\**

341. AT all places north or south of the equator the observer is in an oblique sphere; the celestial equator, and the parallels of declination are oblique to the horizon; and the equator is bisected by the horizon, while the parallels are divided by it into unequal parts. This position of the sphere is represented in Fig. 7, where H O R is the horizon, Q O E the equator, and *n c r, s c t* &c. parallels of declination. Now, the length of the day is measured by the portion of the parallel to the equator described by the sun, which lies above the horizon. On inspecting Fig. 7, it will be evident that this diminishes continually while the sun is moving from the position of greatest northern declination to that of greatest southern declination, and increases continually while he is moving from the position of greatest southern to that of greatest northern declination. Whence it appears that the day will diminish in length from the summer to the winter solstice, and increase in length from the winter to the summer solstice.

---

\* The day, here considered, is the interval between sunrise and sunset.



342. As the equator is bisected by the horizon, at the equinoxes the day and night must be each 12 hours long.

343. When the sun is north of the equator, the greater part of its diurnal circle lies above the horizon, in northern latitudes; and therefore from the vernal to the autumnal equinox the day is, in the northern hemisphere, more than 12 hours in length. On the other hand, when the sun is south of the equator, the greater part of its circle lies below the horizon, and hence from the autumnal to the vernal equinox the day is less than 12 hours in length.

In the latter interval the nights will obviously, at corresponding periods, be of the same length as the days in the former.

344. The variation in the length of the day in the course of the year, will increase with the latitude of the place; for the greater is the latitude the more oblique are the circles described by the sun to the horizon, and the greater is the disparity between the parts into which they are divided by the horizon. This will be obvious, on referring to Fig. 7, where  $H O R$ ,  $H' O R'$ , represent the positions of the horizons of two different places with respect to these circles,  $H' O R'$  being the horizon for which the latitude or the altitude of the pole is the greatest.

For the same reason, the days will be the longer as we proceed from the equator northward, during the period that the sun is north of the equinoctial, and the shorter, during the period that he is south of this circle.

345. At the equator the horizon bisects all the diurnal circles, and consequently the day and night are there each 12 hours in length throughout the year.

346. At the arctic circle the day will be 24 hours long at the time of the summer solstice; for, the polar distance of the sun will then be  $66\frac{1}{2}^{\circ}$ , which is the same as the latitude of the arctic circle, whence it follows that the diurnal circle of the sun at this epoch, will correspond to the circle of perpetual apparition for the parallel in question.

On the other hand, when the sun is at the winter solstice, the night will be 24 hours long on the arctic circle.

347. To the north of the arctic circle, the sun will remain continually above the horizon, during the period that his north

polar distance is less than the latitude of the place, and continually below the horizon during the period that his south polar distance is less than the latitude of the place.

At the north pole, as the horizon is coincident with the equator, the sun will be above the horizon while passing from the vernal to the autumnal equinox, and below it while passing from the autumnal to the vernal equinox. Accordingly, at this locality there will be but one day and one night in the course of a year, and each will be of six months' duration.

348. The circumstances of the duration of light and darkness are obviously the same in the southern hemisphere as in the northern, for corresponding latitudes and corresponding declinations of the sun.

349. *The latitude of the place and the declination of the sun being given, to find the times of the sun's rising and setting and the length of the day.*

Let  $H P R$  (Fig. 49) be the meridian,  $H M R$  the horizon, and  $B s D$  the diurnal circle described by the sun. The hour angle  $E P t$ , or its measure  $E t$ , which converted into time expresses the interval between the rising or setting of the sun and his passage over the meridian, is called the *Semi-diurnal Arc*. Now,

$$E t = E M + M t = 90^\circ + M t;$$

which gives,  $\cos E t = - \sin M t;$

and we have by Napier's rules,

$$\sin M t = \cot t M s \tan g t s = \tan g P M H \tan g E B = \tan g P H \tan g E B;$$

whence,  $\cos E t = - \tan g P H \tan g E B,$

or,

$$\cos (\text{semi-diurnal arc}) = - \tan g \text{ lat.} \times \tan g \text{ dec.} \dots (80).$$

The semi-diurnal arc (in time) expresses the apparent time of the sun's setting; and subtracted from 12 hours, gives the apparent time of its rising. The double of it will be the length of the day.

In resolving this problem it will, in practice, generally answer to make use of the declination of the sun at noon of the given day, which may be taken from an ephemeris.

Exam. 1. Let it be required to find the apparent times of the

sun's rising and setting, and the length of the day at New York, at the summer solstice.

Log. tang lat. ( $40^{\circ} 42' 40''$ )	.	.	9.93474 —
Log. tang dec. ( $23^{\circ} 27' 40''$ )	.	.	9.63749
			<hr/>
Log. cos (semi-diurnal arc)	.	.	9.57223 —
Semi-diurnal arc	.	.	$111^{\circ} 55' 40''$
Time of sun's setting	.	.	7h. 27m. 43s.
Time of sun's rising	.	.	4 32 17
Length of day	.	.	14 55 26

Exam. 2. What are the lengths of the longest and shortest days at Boston; the latitude of that place being  $42^{\circ} 21' 15''$  N.

Ans. 15h. 6m. 28s. and 8h. 53m. 32s.

Exam. 3. At what hours did the sun rise and set on May 1st, 1837, at Charleston; the latitude of Charleston being  $32^{\circ} 47'$ , and the declination of the sun being  $15^{\circ} 6' 0''$  N.

Ans. Time of rising, 5h. 19m. 58s. Time of setting, 6h. 40m. 2s.

350. *To find the time of the sun's apparent rising or setting; the latitude of the place and the declination of the sun being given.*

At the time of his apparent rising or setting, the sun as seen from the centre of the earth will be below the horizon a distance  $sS$  (Fig. 49) equal to the refraction minus the parallax. The mean difference of these quantities is  $33' 42''$ . Let it be denoted by  $R$ . Now to find the hour angle  $ZPS$  ( $=P$ ), the triangle  $ZPS$  gives (see Appendix),

$$k = \frac{ZP + PS + ZS}{2} = \frac{\text{co. lat.} + \text{co. dec.} + (90^{\circ} + R)}{2} \dots (81),$$

$$\sin^2 \frac{1}{2} P = \frac{\sin (k - ZP) \sin (k - PS)}{\sin ZP \sin PS};$$

$$\text{or,} \quad \sin^2 \frac{1}{2} P = \frac{\sin (k - \text{co. lat.}) \sin (k - \text{co. dec.})}{\sin (\text{co. lat.}) \sin (\text{co. dec.})} \dots (82).$$

The value of  $P$ , (in time), will be the interval between apparent noon and the time of the apparent rising or setting.

If the time of the rising or setting of the upper limb of the sun, instead of its centre, be required, we must take for  $R$ ,  $33' 42'' +$  sun's semi-diameter, or  $49' 43''$ .

Unless very accurate results are desired, it will be sufficient to

take the declinations of the sun at 6 o'clock in the morning and evening. When the greatest precision is required, the times of true rising and setting must be computed by equation (80), and the declinations found for these times.

### *Twilight.*

351. When the sun has descended below the horizon, its rays still continue to fall upon a certain portion of the body of air that lies above it, and are thence reflected down upon the earth, so as to occasion a certain degree of light, which gradually diminishes, as the sun descends farther below the horizon and the portion of the air posited above the horizon, that is directly illuminated, becomes less. The same effect, though in a reverse order, takes place in the morning previous to the sun's rising. The light thus produced is called the *Crepusculum*, or *Twilight*.

352. The close of the evening twilight is marked by the appearance of faint stars over the western horizon, and the beginning of the morning twilight, by the disappearance of faint stars situated in the vicinity of the eastern horizon. It has been ascertained from numerous observations, that, at the beginning of the morning and end of the evening twilight, the sun is about  $18^\circ$  below the horizon.

353. *The latitude of the place and the sun's declination being given, to find the time of the beginning or end of twilight.*

The zenith distance of the sun at the beginning of morning or end of evening twilight, is  $90^\circ + 18^\circ$ : wherefore, we may solve this problem by means of equations (81) and (82), taking  $R = 18^\circ$ .

If the time of the commencement of morning twilight be subtracted from the time of sunrise, the remainder will be the duration of twilight.

At the latitude  $49^\circ$ , the sun at the time of the summer solstice is only  $18^\circ$  below the horizon, at midnight; for, the altitude of the pole at a place the latitude of which is  $49^\circ$ , differs only  $18^\circ$  from the polar distance of the sun at this epoch. At this latitude, therefore, twilight will continue all night, at the summer solstice. This will be true for a still stronger reason at greater latitudes.

354. The duration of twilight varies with the latitude of the place and with the time of the year. At all places in the northern hemisphere, the summer are longer than the winter twilights; and the longest twilights take place at the summer solstice; while the shortest occur when the sun has a small southern declination, different for each latitude.\* The summer twilights increase in length from the equator northward.

355. At the Pole twilight commences about a month and a half before the sun appears above the horizon, and lasts about a month and a half after he has disappeared.

### *The Seasons.*

356. The amount of heat received from the sun in the course of 24 hours, depends upon two particulars, the time of the sun's continuance above the horizon, and the obliquity of his rays at noon. By reason of the obliquity of the ecliptic, both of these circumstances vary materially in the course of the year; whence arises a variation of temperature or a change of seasons.

357. The tropics and the arctic circles divide the earth into five parts called *Zones*, throughout each of which the yearly change of the temperature is occasioned by a similar change in the circumstances upon which it depends.

The part contained between the two tropics, is called the *Torrid Zone*; the two parts between the tropics and polar circles, are called the *Temperate Zones*; and the other two parts, within the polar circles, are called *Frigid Zones*.

358. At all places in the north temperate zone the sun will always pass the meridian to the south of the zenith; for, the latitudes of all such places exceed  $23\frac{1}{2}^{\circ}$ , the greatest declination of the sun. The meridian zenith distance will be greatest at the winter solstice, when the sun has its greatest southern declina-

\* The duration of shortest twilight is given by the following formula :

$$\sin a = \frac{\sin 9^{\circ}}{\cos \text{lat.}}$$

Twice the angle  $a$ , converted into time, expresses the duration of shortest twilight. To find the sun's declination at the time of shortest twilight, we have,

$$\sin \text{dec.} = -\tan 9^{\circ} \sin \text{lat.}$$

(For the investigation of this and the preceding formula, see Gummere's Astronomy, pages 87 and 88.)



tion, and least at the summer solstice, when the sun has its greatest northern declination ; and it will vary continually between the values which obtain at these epochs. The day will be longest at the summer solstice, and the shortest at the winter solstice, and will vary in length progressively from the one date to the other.

We infer, therefore, that throughout the zone in question the greatest amount of heat will be received from the sun at the summer solstice, and the least at the winter solstice ; and that the amount received will gradually increase, or decrease, from one of these epochs to the other. The solstices are not, however, the epochs of maximum and minimum temperature, but are found from observation to precede these by about a month. The reason of this circumstance is, that the earth continues for a month, or thereabouts, after the summer solstice, to receive during the day more heat than it loses during the night, and for about the same length of time after the winter solstice, continues to lose during the night more heat than it receives during the day.

359. Within the torrid zone the length of the day varies after the same manner as in the temperate zone, though in a less degree ; but the motion of the sun with respect to the zenith, is different. At all places in the torrid zone the sun passes the meridian during a certain portion of the year to the south of the zenith, and during the remaining portion to the north of it ; for, all places so situated have their zeniths between the tropics in the heavens, and the sun moves from one tropic to the other, and back again to its original position, in a tropical year. Throughout the torrid zone, therefore, the sun will be in the zenith twice in the course of the year, and will be at its maximum distance from it on the one side and the other, at the solstices.

An inhabitant of the equator or its vicinity, will have summer at the two periods when the sun is in the zenith, and winter, (or a period of minimum temperature,) both at the summer and winter solstice. Near the tropics, there will be but little variation in the daily amount of heat received, during the period that the sun is north of the zenith.

360. At the frigid zone a new cause of a change of temperature exists ; the sun remains continually above the horizon, for a greater or less number of days about the summer solstice, and

continually below it for the same number of days about the winter solstice.

361. The amount of the yearly variation of temperature increases with the latitude of the place; for, the greater is the latitude, the greater will be the variation in the length of the day. Also, the mean yearly temperature is lower, as we recede from the equator and approach the poles; for, since the sun is, in the course of the year, the same length of time above the horizon, at all places, the mean yearly temperature must depend altogether upon the mean obliquity of the sun's rays at noon, and this increases with the latitude.

362. The yearly change in the sun's distance from the earth, has but little effect in producing a variation of temperature upon the earth's surface. The change of its heating power from this cause amounts to no more than  $\frac{1}{15}$ .

363. It is important to observe, that, although in the main climate varies with the latitude after the manner explained in the foregoing articles, it is still dependent more or less upon local circumstances, such as the vicinity of lakes, seas, and mountains, prevailing winds of some particular direction, &c.

364. In the north temperate zone, *Spring*, *Summer*, *Autumn*, and *Winter*, the four seasons into which the year is divided, are considered as respectively commencing at the times of the *Vernal Equinox*, *Summer Solstice*, *Autumnal Equinox*, and *Winter Solstice*.

365. Let V (Fig. 50) represent the vernal, and A the autumnal equinox; S the summer, and W the winter solstice. The perigee of the sun's apparent orbit is at present about  $10^{\circ} 10'$  to the east of the winter solstice. Let P denote its position. The lengths of the seasons are, agreeably to Kepler's law of areas, respectively proportional to the areas V E S, S E A, A E W, and W E V. Thus, the winter is the shortest season, and the summer the longest; and spring is longer than autumn. Spring and summer, taken together, are about 7 days longer than autumn and winter united.

366. Since the perigee of the sun's orbit has a progressive motion, the relative lengths of the seasons must be subject to a continual variation.

367. At the beginning of the year 1800, the longitude of the

sun's perigee was  $279^{\circ} 30' 8''.39$ . If from this we take  $180^{\circ}$ , the longitude of the autumnal equinox, the remainder,  $99^{\circ} 30' 8''.39$ , is the distance of the perigee from the autumnal equinox at that epoch. The motion of the perigee in longitude is at the rate of  $61''.52$  per year. Dividing  $99^{\circ} 30' 8''.39$  by  $61''.52$ , the quotient is 5822. Hence it appears that about 5800 years anterior to the year 1800, the perigee coincided with the autumnal equinox, and the apogee with the vernal equinox. It is perhaps worthy of remark, that this is about the period at which most chronologists fix the first residence of man upon the earth.

*Appearance, Dimensions, and Physical Constitution of the Sun.*

368. The sun presents the appearance of a luminous circular disc. But it does not necessarily follow from this that its surface is really flat; for, such is the appearance of all globular bodies, when viewed at a great distance. It is ascertained from observations with the telescope, that the sun has a rotatory motion: this being the fact, its surface must in reality be of a spherical form; for, otherwise, it would not, in presenting all its sides, always appear under the form of a circle.

369. The sun's real diameter is determined from his apparent diameter and horizontal parallax. Let A C B (Fig. 51) represent the sun or other heavenly body, and E the place of the earth; and let  $\delta = \angle AEB$  the sun's apparent diameter,  $d = 2 AS$  his real diameter,  $D = ES$  his distance from the earth, and  $R$  = the radius of the earth. We have from the triangle A E S,

$$AS = ES \tan \frac{1}{2} \angle AEB, \text{ or, } 2 AS = 2 ES \tan \frac{1}{2} \angle AEB;$$

and thus,  $d = 2 D \tan \frac{1}{2} \delta;$

but,  $D = \frac{R}{\sin H} \dots (\text{equa. 9});$

whence,  $d = 2 R \frac{\tan \frac{1}{2} \delta}{\sin H} = 2 R \frac{\frac{1}{2} \delta}{H} = 2 R \frac{\delta}{2 H}. (83).$

The mean apparent diameter of the sun is  $32' 1''.8$ , and his mean horizontal parallax  $8''.58$ . Accordingly we have, for the real diameter of the sun,

$$d = 2 R \frac{32' 1''.8}{2 (8''.58)} = 2 R \times 112 (\text{nearly}).$$

Thus the diameter of the sun is about 112 times the diameter of the earth. The volume of the sun then exceeds that of the earth nearly in the proportion  $112^3$  to  $1^3$ , or 1407168 to 1.

370. From equation (S3), we may derive the proportion

$$d : 2 R :: \delta : 2 H.$$

Thus, *the real diameter of a heavenly body is to the diameter of the earth, as the apparent diameter of the body is to double its horizontal parallax.*

371. When the sun is viewed with a telescope of considerable power and provided with colored glasses, black spots of an irregular form, surrounded by a dark border, or penumbra, are often seen on its disc. Their number, magnitude, and position on the disc, are extremely variable. In some years they are very frequent, and appear in large numbers; in others, none whatever are seen. In some instances, as many as fifty, of various forms and sizes, have been counted. Their absolute magnitude is often very great. Spots are not unfrequently seen that subtend an angle of  $1'$  or  $60''$ . Now, the apparent diameter of the earth as viewed at the distance of the sun, is equal to double the sun's horizontal parallax, or  $17''$ : the breadth of such spots must therefore exceed three times the diameter of the earth, or 24,000 miles. Spots two or three times as large as this have been seen.

372. The form and size of the spots are subject to rapid and almost incessant variations. When watched from day to day, or even from hour to hour, they are seen to enlarge or contract, and at the same time to change their forms. When a spot disappears, it always contracts into a point, and vanishes before the penumbra. Some spots disappear almost immediately after they become visible; others remain for weeks, or even months.

373. Spots and streaks more luminous than the general body of the sun are also frequently perceived upon parts of his disc, especially in the region of large spots, or of extensive groups of spots. These are called *Faculae*. The penumbra which surrounds each black spot is also abruptly terminated by a border of light more brilliant than the rest of the disc.

374. When the positions of the spots on the disc are observed from day to day, it is perceived that they all have a common motion in a direction from east to west. Some of the spots

close up and vanish before they reach the western limb ; others disappear at the western limb, and are never afterwards seen ; a few, after becoming visible at the eastern limb, have been seen to pass entirely across the disc, disappear from view at the western limb, and re-appear again at the eastern limb. The time employed by a spot in traversing the sun's disc, is about 14 days. The same time is occupied in passing from the western to the eastern limb, while it is invisible. The motions of the spots are accounted for, in all their circumstances, by supposing that the sun has a motion of rotation from west to east, around an axis nearly perpendicular to the plane of the ecliptic ; and that the spots are portions of the solid body of the sun. The truth of this explanation of the apparent motions of the sun's spots, is confirmed by the changes which are observed to take place in the magnitude and form of the more permanent spots during their passage across the disc. When they first come into view at the eastern limb, they appear as a narrow dark streak. As they advance towards the middle of the disc, they gradually open out, and increase in magnitude ; and after they have passed the middle of the disc, contract by the same degrees until they are again seen as a mere dark line upon the western limb.

375. A spot returns to the same position on the disc in about  $27\frac{1}{2}$  days. This is not, however, the precise period of the sun's rotation ; for, during this time the sun has apparently moved forward nearly a sign in the ecliptic ; the spot will therefore have accomplished so much more than a complete revolution, when it is again seen by an observer on the earth in the same position on the disc.

376. The apparent position of a spot with respect to the sun's centre may be accurately determined, from day to day, by observing, when the sun is crossing the meridian, the right ascensions and declinations both of the spot and centre. From three or more observations of this kind the period of the sun's rotation and the position of his equator may be ascertained.

377. The time of the sun's rotation on his axis is about  $25\frac{1}{2}$  days ; the inclination of his equator to the ecliptic  $7^{\circ} 30'$  ; and the heliocentric longitude of the ascending node of the equator  $80^{\circ} 7'$ .



378. The only theories relative to the physical constitution of the sun, which deserve notice, are those of Laplace and Herschel. Laplace supposed that the sun was an immense globe of solid matter in a state of ignition, and that the spots upon his disc were large cavities, where there was a temporary intermission in the evolution of luminous matter. Sir W. Herschel was of opinion that the sun was an opaque solid body, surrounded by two atmospheres, of which the first was opaque and non-luminous, and the second luminous. On this hypothesis the spots are accounted for by supposing that openings occasionally take place in the atmospheres, through which the dark body of the sun is seen. The penumbra is the portion of the obscure atmosphere, situated immediately around the opening made in it. This theory seems to account for all the circumstances of the aspect and variation of the form and magnitude of the spots, which the other does not do. Moreover, M. Fourier has remarked, that the light given out by inc<sup>a</sup>ndescent gases is not polarized, while that emitted by solids or liquids in combustion, is ; and M. Arago has ascertained that the light of the sun, in effect, does not possess the properties of polarized light in any sensible degree, which seems to prove that it emanates from a gas.

379. There has been observed, in connection with the sun, at certain periods of the year, a faint light that is visible before sunrise and after sunset, to which has been given the name of the *Zodiacal Light*, from the circumstance of its being mostly comprehended within the zodiac. Its colour is white, and its apparent figure that of a spindle, the base of which rests on the sun, and the axis of which lies in the plane of the sun's equator ; such as would be the appearance of an ellipsoid of revolution, having its centre coincident with that of the sun, and its transverse axis in the plane of the solar equator. Its extent varies with the season of the year and the state of the atmosphere ; being sometimes more than  $100^{\circ}$ , and at other times not more than  $40^{\circ}$  or  $50^{\circ}$ . No satisfactory explanation has yet been given of this singular phenomenon. It was long supposed to be the atmosphere of the sun, but Laplace has shown that this hypothesis is opposed to the theory of gravitation.

380. The zodiacal light is seen most distinctly in our north-

ern climates, early in the spring just after sunset, and early in the fall just before sunrise. During the month of March it may be seen directed towards the star Aldebaran. Towards the summer solstice, it is said to be discernible, in a very pure state of the atmosphere, both in the morning and evening. The reason of these variations in the distinctness of the zodiacal light, is found in the change of its inclination to the horizon at the time of sunset or sunrise. As its length lies in the plane of the sun's equator, its inclination to the horizon will be different, like that of this plane, according to the different positions of the sun in the ecliptic. Since the sun's equator makes but a small angle with the ecliptic, at sunset, the zodiacal light will be most inclined to the horizon, and therefore extend higher up in the heavens, towards the vernal equinox, when the inclination of the ecliptic to the horizon at sunset is at its maximum; and, at sunrise, it will be most inclined to the horizon towards the autumnal equinox, when the inclination of the ecliptic to the horizon at sunrise is the greatest.

---

## CHAPTER XV.

### OF THE MOON AND ITS PHENOMENA.

#### *Phases of the Moon.*

381. The most conspicuous of the phenomena exhibited by the moon, is the periodical change that is observed to take place in the form and size of its disc. The different appearances which the disc presents are called the *Phases* of the moon.

The phenomenon in question is a simple consequence of the revolution of the moon around the earth. Let E (Fig. 52) represent the position of the earth, A B C, &c. the orbit of the moon, which we will suppose for the present to lie in the plane of the

ecliptic, and E S the direction of the sun. As the distance of the sun from the earth is about 400 times the distance of the moon, lines drawn from the sun to the different parts of the moon's orbit, may be considered, without material error, as parallel to each other. If we regard the moon as an opaque non-luminous body, that hemisphere which is turned towards the sun, will continually be illuminated by him, and the other will be in the dark. Now, by virtue of the moon's motion, the enlightened hemisphere is presented to the earth under every variety of aspect in the course of a synodical revolution of the moon. Thus, when the moon is in conjunction, as at A, this hemisphere is turned entirely away from the earth, and she is invisible. Soon after conjunction, a portion of it on the right begins to be seen, and as this is comprised between the right half of the circle which limits the vision, and the right half of the circle which separates the enlightened and dark hemispheres of the moon, called the *Circle of Illumination*, it will obviously present the appearance of a crescent with the horns turned from the sun, as represented at B. As the moon advances, more and more of the enlightened half becomes visible, and thus the crescent enlarges, and the eastern limb becomes less concave. At the point C,  $90^\circ$  distant from the sun, one half of it is seen, and the disc is a semi-circle, the eastern limb being a right line. Beyond this point, more than half becomes visible; the nearer half of the circle of illumination falls to the left of the moon's centre, as seen from the earth, and thus becomes convex outwards. This phase of the moon is represented at D. When the moon appears under this shape, it is said to be *Gibbous*. In advancing towards opposition, the disc will enlarge, and the eastern limb become continually more convex; and finally at opposition, where the whole illuminated face is seen from the earth, it will become a full circle. From opposition to conjunction, the nearer half of the circle of illumination will form the right or western limb, and this limb will pass in the inverse order through the same variety of forms, as the eastern limb in the interval between conjunction and opposition. The different phases are delineated in the figure.

382. The moon's orbit is, in fact, somewhat inclined to the plane of the ecliptic, instead of lying in it, as we have sup-

posed; but, it is plain that its inclination cannot change the order, nor the period of the phases, and that it can have no other effect than to alter somewhat the size of the disc, at particular angular distances from the sun. In consequence of the smallness of the inclination, this alteration is too slight to be noticed.

383. When the moon is in conjunction, it is said to be *New Moon*; and when in opposition, *Full Moon*. At the time between new and full moon, when the difference of the longitudes of the moon and sun is  $90^\circ$ , it is said to be the *First Quarter*. And at the corresponding time between full and new moon, it is said to be the *Last Quarter*. In both these positions the moon appears as a semi-circle, and is said to be *dichotomized*. The two positions of conjunction and opposition, are called *Syzigies*; and those of the first and last quarter, *Quadratures*. The four points midway between the syzigies and quadratures, are called *Octants*.

384. The interval from new moon to new moon again, is called a *Lunar Month*, and sometimes a *Lunation*.

The mean daily motion of the sun in longitude is  $59' 8''.33$ , and that of the moon  $13^\circ 10' 35''.03$ ; wherefore the moon separates from the sun at the mean rate of  $12^\circ 11' 26''.70$  per day; and hence, to find the mean length of a lunar month, we have the proportion,

$$12^\circ 11' 26''.70 : 1\text{d.} :: 360^\circ : x = 29\text{d. } 12\text{h. } 44\text{m. } 2.7\text{s.}$$

385. *To determine the time of mean new or full moon in any given month.*

Let the mean longitude of the sun, and also the mean longitude of the moon, at the beginning of the year, be found, and let the former be subtracted from the latter (adding  $360^\circ$  if necessary); the remainder, which call R, will be the mean distance of the moon to the east of the sun, at the beginning of the year. As the moon separates from the sun at the mean rate of  $12^\circ 11' 26''.70$  per day,  $\frac{R}{12^\circ 11' 26''.70}$  will express the number of days and fractions of a day, which at this epoch have elapsed since the last new moon. This interval is called the *Astronomical Epact*. If we subtract it from 29d. 12h. 44m. 2.7s. we shall have the time

of mean new moon in January. This being known, the time of mean new moon in any other month of the year results very readily from the known length of a lunar month.

The time of mean new moon in any month being known, the time of mean full moon in the same month is obtained by the addition or subtraction, as the case may be, of half a lunar month.

This problem is in practice most easily resolved with the aid of tables. (See Problem XXVII.)

386. The time of true new moon differs from the time of mean new moon, for the same reasons that the true longitudes of the sun and moon differ from the mean. The same is true of the time of true full moon. For the mode of computing the time of true new or full moon from that of mean new or full moon, (see Problem XXVII.)

387. The earth, as viewed from the moon, goes through the same phases in the course of a lunar month that the moon does to an inhabitant of the earth. But, at any given time, the phase of the earth is just the opposite to the phase of the moon. About the time of new moon, the earth, then near its full, reflects so much light to the moon as to render the obscure part visible.

*Moon's Rising, Setting, and Passage over the Meridian.*

388. *To find the time of the meridian passage of the moon on a given day.*

Let  $S$  and  $M$  denote, respectively, the right ascension of the sun, and the right ascension of the moon, at noon on the given day, and  $M, S$  the hourly variations of the right ascension of the sun and moon : also let  $t$  = the required time of the meridian passage. At the time  $t$  the right ascensions will be,

For the moon, . . . .  $M + t m$ ,

For the sun, . . . .  $S + t s$ ;

and, as the moon is on the meridian, the difference of these arcs will be equal to the hour angle  $t$  ; whence,

$$t = M - S + t (m - s) ;$$

or, if all the quantities be expressed in seconds,

$$t = M - S + t \cdot \frac{m - s}{3600} \dots (84).$$



Thus, we find for the time of the meridian passage,

$$t = \frac{3600 (M - S)}{3600 - (m - s)} \dots (85).$$

The quantities  $M$ ,  $S$ ,  $m$ ,  $s$ , are, in practice, to be taken from ephemerides of the sun and moon.

Example. What was the time of the passage of the moon's centre over the meridian of New York, on the 1st of August, 1837?

When it is noon at New York, it is 4h. 56m. 4s. at Greenwich. Now, by the Nautical Almanac,

Aug. 1st, at 4h. ♀'s R. Ascen.	. . .	8 <sup>h</sup> . 58 <sup>m</sup> . 36.7 <sup>s</sup> .
“ at 5h. “ “ . . .		9 0 38.3

---

1h. : 56m. 4s. :: 2m. 1.6s. : 1m. 53.6s.

Aug. 1st, at 4h. ♀'s R. Ascen.	. . .	8 <sup>h</sup> . 58 <sup>m</sup> . 36.7 <sup>s</sup> .
Variation of R. Ascen. in 56m. 4s. .		1 53.6

---

♀'s R. Ascen. at M. Noon at N. York, 9 0 30.3

Aug. 1st, ☉'s hourly Variation of R. Ascen. . . . 9.704s.

1h. : 4h. 56m. 4s. :: 9.704s. : 47.8s.

Aug. 1st, M. Noon at Greenw., ☉'s R. Asc.	8 <sup>h</sup> . 45 <sup>m</sup> . 31.5 <sup>s</sup> .
Variation of R. Ascen. in 4h. 56m. 4s.	47.8

---

☉'s R. Ascen. at M. Noon at N. York,	8 46 19.3 (S)
♀'s “ “ “	9 0 30.3 (M)

---

$M - S = 14 \quad 11.0 = 851.0s.$

Aug. 1st, M. Noon at Greenw., ♀'s R. Asc. 8<sup>h</sup>. 50<sup>m</sup>. 27.7<sup>s</sup>.

Aug. 2d, “ “ “	9 38 18.7
----------------	-----------

---

24 ) 47 51.0

Aug. 1st, ♀'s mean hourly Varia. of R. Asc. 1 59.6 ( $m$ )

“ ☉'s “ “ “	9.7 ( $s$ )
-------------	-------------

---

$m - s = 1 \quad 49.9 = 109.9s.$

$$\begin{array}{rcl}
 3600 & . & . & . & . & \log. & 3.55630 \\
 M - S = 851.0 & . & . & . & . & \log. & 2.92993 \\
 3600 - (m - s) = 3490.1 & . & . & & \text{ar. co.} & \log. & 6.45716
 \end{array}$$


---

Appar. time of merid. passage, 14m. 37.8s. = 877.8s.  $\log.$  2.94339

Equa. of time by Almanac,  $\begin{array}{cc} 5 & 58 \end{array}$

---

Mean time of merid. pass., 0h. 20m. 36s.

The Nautical Almanac gives the time of the moon's passage over the meridian of Greenwich for every day of the year. From this, the time of the passage across the meridian of any other place may easily be determined, as follows: subtract the time of the meridian passage at Greenwich on the given day, from that on the following day, and say, as 24h.: the difference :: the longitude of the place : a fourth term. This fourth term added to the time of the meridian passage at Greenwich on the given day, will give the approximate time of the meridian passage on the same day at the given place. The fourth term of the preceding proportion may be corrected, and a more exact result obtained by stating the proportion, as 24h. : 24h. + difference above mentioned :: fourth term in question : the same corrected.

389. Since the moon has a motion with respect to the sun, the time of its rising and setting must vary from day to day. When first seen after conjunction, it will set soon after the sun. After this it will set (at a mean) about one hour later every succeeding night. At the first quarter, it will set about midnight; and at full moon, will set about sunrise and rise about sunset. During this interval it will rise in the day time, and all along from sunrise to sunset. From full to new moon, it will rise at night, and set during the day; and the time of the rising and setting will be about an hour later on every succeeding night and day; thus, at the last quarter it will rise about midnight and set about midday.

390. *To find the time of the moon's rising or setting on any given day.*

Compute the moon's semi-diurnal arc from equation (82), or (80), according as it is the time of the apparent rising or setting, or the time of the true rising or setting, that is desired. Correct

it for the moon's change of right ascension in the interval between the moon's passage over the meridian and setting, by the following proportion,  $24h. : 24 + m - S$  (Art. 388) :: semi-diurnal arc : corrected semi-diurnal arc ; and add it to the time of the moon's meridian passage, found as explained in Art. 388. The result will be the time of the moon's setting ; and if this be subtracted from 24 hours, the remainder will be the time of the moon's rising.

In consequence of the change of the moon's declination in the interval between its rising and setting, it would be more accurate to compute the semi-diurnal arc separately for the moon's rising. In computing the semi-diurnal arc by equation (80), the declination 6 hours before or after the meridian passage may be used at first ; and afterwards, if a more accurate result be desired, the calculation may be repeated with the declination found for the computed approximate time. In equation (82),  $R =$  refraction — parallax  $= 33' 51'' - 57' 1''$  (at a mean)  $= - 23' 10''$ .

### *Rotation and Librations of the Moon.*

391. The moon presents continually nearly the same face towards the earth ; for, the same spots are always seen in nearly the same position upon the disc. It follows, therefore, that it rotates on its axis in the same direction, and with the same angular velocity, or nearly so, that it revolves in its orbit.

392. The spots on the moon's disc, although they constantly preserve very nearly the same situations, are not, however, strictly stationary. When carefully observed, they are seen alternately to approach and recede from the edge. Those that are very near the edge successively disappear and again become visible. This vibratory motion of the moon's spots is called *Libration*.

393. There are three librations of the moon, that is, a vibratory motion of its spots from three distinct causes.

1. The moon's motion of rotation being uniform, small portions on its east and west sides alternately come into sight and disappear, in consequence of its *unequal motion in its orbit*. The periodical oscillation of the spots in an easterly and westerly direction from this cause, is called the *Libration in Longitude*.

2. The lunar spots have also a small alternate motion from north to south. This is called the *Libration in Latitude*, and is accounted for, by supposing that the moon's axis is not exactly

perpendicular to the plane of its orbit, and that it remains continually parallel to itself. On this supposition we ought sometimes to see beyond the north pole of the moon, and sometimes beyond the south pole.

3. Parallax is the cause of a third libration of the moon. The spectator upon the earth's surface being removed from its centre, the point towards which the moon continually presents the same hemisphere, he will see portions of the moon a little different according to its different positions above the horizon. The diurnal motion of the spots resulting from the parallax, is called the *Diurnal* or *Parallactic Libration*.

394. The exact position of the moon's equator, like that of the sun's, is derived from accurate observations of the situations of the spots upon the disc. From calculations founded upon such observations, it has been ascertained that the plane of the moon's equator is constantly inclined to the plane of the ecliptic under an angle of  $1^{\circ} 30'$ , and intersects it in a line which is always parallel to the line of the nodes. It follows from the last mentioned circumstance, that if a plane be supposed to pass through the centre of the moon, parallel to the ecliptic, it will intersect the plane of the moon's equator and that of its orbit, in the same line in which these planes intersect each other. The plane in question will lie between the plane of the equator and that of the orbit. It will make with the first an angle of  $1^{\circ} 30'$ , and with the second an angle of  $5^{\circ} 9'$ .

*Dimensions and Physical Constitution of the Moon.*

395. The phases of the moon prove it to be an opaque spherical body. Its diameter is found by means of equation (83), viz :

$$d = 2 R \frac{\delta}{2 H},$$

where  $d$  denotes the diameter sought,  $R$  the radius of the earth,  $\delta$  the apparent diameter of the moon at a given distance, and  $H$  its horizontal parallax at the same distance.

The greatest equatorial horizontal parallax of the moon is  $61' 24''$ , and the corresponding apparent diameter  $33' 31''$  : thus we have,

$$d = 2 R \frac{61' 24''}{33' 31''} = 2 R \frac{3}{11} \text{ (very nearly) } = 2161 \text{ miles.}$$

396. The diameter of the moon being to the diameter of the earth as 3 to 11, the surface of the moon is to the surface of the earth as  $3^2$  to  $11^2$ , or as 1 to 13; and the volume of the moon is to the volume of the earth as  $3^3$  to  $11^3$ , or as 1 to 49.

397. When the moon is viewed with a telescope, the edge of the disc, which borders upon the dark portion of the face, is seen to be very irregular and serrated. It is hence inferred that the surface of the moon is diversified with mountains and valleys. The truth of this inference is confirmed, by the fact, that bright insulated spots are frequently seen on the dark part of the face near the edge of the disc, which gradually enlarge until they become united to it. These bright spots are doubtless the tops of mountains illuminated by the sun, while the surrounding regions that are less elevated, are involved in darkness. The disc is also diversified with spots of different shapes and different degrees of brightness. The brighter parts are supposed to be elevated land and the dark to be valleys or cavities.

398. The number of the lunar mountains is very great. Their form and grouping is for the most part, similar to what obtains in volcanic districts of the earth: from which it is inferred that they are of volcanic origin.

399. From measurements made with the micrometer, of the lengths of their shadows, or of the distance of their summits, when first illuminated, from the adjacent boundary of the disc, the heights of a number of the lunar mountains have been computed. According to Herschel, the altitude of the highest is about  $1\frac{3}{4}$  English miles.\*

400. There seems to be no large bodies of water upon the surface of the moon, or, at least, upon the hemisphere which is turned towards the earth; for, the boundary of the illuminated hemisphere is in all its positions irregular throughout, whereas, if it ever fell upon any large body of water, it would, for the extent of it, be an unbroken and regular curve.

The moon also has no atmosphere, or, if it has any, it is so rare as not sensibly to diminish or refract the light of the stars, passing

---

\* Schroeter makes the elevation of some of the lunar mountains to exceed 5 miles.



through it; for, when a star experiences an occultation from the moon, it does not disappear until the body of the moon reaches it, and the duration of the occultation is as it is computed, without making any allowance for the refraction of the atmosphere.

---

## CHAPTER XVI.

### ECLIPSES OF THE SUN AND MOON.—OCCULTATIONS OF THE FIXED STARS.

401. An eclipse of a heavenly body is a privation of its light, occasioned by the interposition of some opaque body between it and the eye, or between it and the sun. Eclipses are divided, with respect to the objects eclipsed, into *eclipses of the sun, of the moon, and of the satellites* (Art. 304); and, with respect to circumstances, into *total, partial, annular, and central*. A *total* eclipse is one in which the whole disc of the luminary is darkened; a *partial* one is when only a part of the disc is darkened. In an *annular* eclipse the whole is darkened, except a ring or annulus, which appears round the dark part like an illuminated border; the definition of a central eclipse will be given in another place.

#### *Eclipses of the Moon.*

402. An eclipse of the moon is occasioned by an interposition of the body of the earth directly between the sun and moon, and thus intercepting the light of the sun; or the moon is eclipsed when it passes through part of the shadow of the earth, as projected from the sun. Hence it is obvious that lunar eclipses can happen only at the time of full moon, for it is then only that the earth can be between the moon and the sun.

403. Since the sun is much larger than the earth, the shadow of the earth must have the form of a cone, the length of which

will depend on the relative magnitudes of the two bodies and their distance from each other. Let the circles  $A G B, a g b$ , (Fig. 53), be sections of the sun and earth by a plane passing through their centres  $S$  and  $E$ ;  $A a, B b$  tangents to these circles on the same side, and  $A d, B c$  tangents on different sides. The triangular space  $a C b$  will be a section of the earth's shadow or *Umbra*, as it is sometimes called. The line  $E C$  is called the *Axis of the Shadow*. If we suppose the line  $c p$  to revolve about  $E C$ , and form the surface of the frustum of a cone, of which  $p c d g$  is a section, the space included within that surface and exterior to the umbra, is called the *Penumbra*. It is plain, that points situated within the umbra will receive no light from the sun; and that points situated within the penumbra will receive light from a portion of the sun's disc, and from a greater portion the more distant they are from the umbra.

404. *To find the length of the earth's shadow.*

Let  $L$  = the length of the shadow;  $R$  = the radius of the earth;  $\delta$  = sun's apparent semi-diameter; and  $p$  = sun's parallax. The right angled triangle  $E a C$  (Fig. 54) gives,

$$E C = \frac{E a}{\text{tang } E C a}.$$

$E a = R$ ; and  $E C a = S E A - E A C = \delta - p$ ; whence,

$$L = \frac{R}{\text{tang } (\delta - p)} \dots (86.)$$

As the angle  $(\delta - p)$  is only about  $16'$ , it will differ but little from its tangent, and therefore,

$$L = R \frac{1}{\delta - p} \text{ (nearly);}$$

or, if  $\delta$  and  $p$  be expressed in seconds,

$$L = R \frac{206264''.8}{\delta - p} \text{ (nearly)} \dots (87).$$

The shadow will obviously be the shortest when the sun is the nearest to the earth. We then have  $\delta = 16' 18''$ , and  $p = 9''$ , which gives  $L = 213 R$ . The greatest distance of the moon is a little less than  $64 R$ . It appears then, that *the earth's shadow always extends to more than three times the distance of the moon*.

405. Let  $k M h$  be a circular arc, described about  $E$  the cen-

tre of the earth, and with a radius equal to the distance between the centres of the earth and moon at the time of opposition. The angle  $ME m$ , the apparent semi-diameter of a section of the earth's shadow, made at the distance of the moon's centre, is called the *Semi-diameter of the Earth's Shadow*. And the angle  $ME h$ , the apparent semi-diameter of a section of the penumbra, at the same distance, is called the *Semi-diameter of the Penumbra*.

406. Were the plane of the moon's orbit coincident with the plane of the ecliptic, there would be a lunar eclipse at every full moon; but, as it is inclined to it, an eclipse can happen only when the full moon takes place either in one of the nodes of the moon's orbit, or so near it that the moon's latitude does not exceed the sum of the apparent semi-diameters of the moon and of the earth's shadow.

To determine the distance from the node, beyond which there can be no eclipse, we must ascertain the semi-diameter of the earth's shadow. Let this be denoted by  $A$ , and let  $P$  = the moon's parallax,

$$ME m = E m a - EC m \text{ (Fig. 54);}$$

but  $E m a = P$ ; and  $EC m = \delta - p$  (Art. 404); therefore,

$$ME m = A = P + p - \delta \dots (SS).$$

407. The semi-diameter of the shadow is the least when the moon is in its apogee and the sun is in its perigee, or when  $P$  has its maximum, and  $\delta$  its minimum value. In these positions of the moon and sun,  $P = 53' 48''$ ,  $\delta = 16' 18''$ , and  $p = 9''$ . Substituting, we obtain for the least semi-diameter of the earth's shadow  $37' 39''$ , and for its least diameter  $1^\circ 15' 18''$ . The greatest apparent diameter of the moon is  $33' 31''$ . Whence it appears, that *the diameter of the earth's shadow is always more than twice the diameter of the moon*.

The mean values of  $P$ ,  $p$ , and  $\delta$  are respectively  $57' 1''$ , and  $16' 1''$ ; which gives for the mean semi-diameter of the earth's shadow  $41' 9''$ .

408. If to  $P + p - \delta$ , the semi-diameter of the earth's shadow, we add  $d$ , the semi-diameter of the moon, the sum  $P + p + d - \delta$  will express the greatest latitude of the moon in opposition, at which an eclipse can happen.

It is easy for a given value of  $P + p + d - \delta$ , and for a given inclination of the moon's orbit, to determine within what distance from the node the moon must be, in order that an eclipse may take place. By taking the least and greatest inclinations of the orbit, the greatest and least values of  $P + p + d - \delta$ , and also taking into view the inequalities in the motions of the sun and moon, it has been found, that when at the time of mean full moon, the difference of the mean longitudes of the moon and node exceeds  $13^{\circ} 21'$ , there cannot be an eclipse; but when this difference is less than  $7^{\circ} 47'$  there must be one. Between  $7^{\circ} 47'$  and  $13^{\circ} 21'$  the happening of the eclipse is doubtful. These numbers are called the *Lunar Ecliptic Limits*.

409. To determine at what full moons in the course of any one year there will be an eclipse, find the time of each mean full moon (Art. 385); and for each of the times obtained find the mean longitude of the sun, and also of the moon's node, and compare the difference of these with the lunar ecliptic limits. Should, however, the difference, in any instance, fall between the two limits, farther calculation will be necessary.

This problem may be solved more expeditiously by means of tables of the sun's mean motion with respect to the moon's node. (See Prob. XXVIII.)

410. The magnitude and duration of an eclipse, depend upon the proximity of the moon to the node at the time of opposition. In order that the centre of the moon may be on the same right line with the centres of the sun and earth, or, in technical language, that a *central* eclipse may happen, the opposition must take place precisely in the node. A strictly central eclipse, therefore, seldom, if ever, occurs. As the mean semi-diameter of the earth's shadow is  $41' 9''$  (Art. 407), and the mean hourly motion of the moon with respect to the sun,  $30' 29''$ , the mean duration of a central eclipse would be about  $2\frac{1}{2}$  h.

411. Since the moon moves from west to east, an eclipse of the moon must commence on the eastern limb, and end on the western.

412. In the investigations in Arts. 404 and 406, we have supposed the cone of the earth's shadow to be formed by lines drawn from the edge of the sun, and touching the earth's surface. This, probably, is not the exact case of nature; for, the duration of the

eclipse, and, by consequence, the apparent diameter of the earth's shadow, is found, by observation, to be somewhat greater than would result from this supposition. This circumstance is accounted for, by supposing those solar rays, that, from their direction, would glance by and rase the earth's surface, to be stopped and absorbed by the lower strata of the atmosphere. In such a case the conical boundary of the earth's shadow would be formed by certain rays exterior to the former, and would be larger.

The moon in approaching and receding from the earth's total shadow, or umbra, passes through the penumbra, and thus its light, instead of being extinguished and recovered suddenly, experiences at the beginning of the eclipse a gradual diminution, and at the end a gradual increase. On this account the times of the beginning and end of the eclipse cannot be noted with precision, and in consequence astronomers differ as to the amount of the increase in the size of the earth's shadow from the cause above mentioned. It is the practice, however, in computing an eclipse of the moon, to increase the semi-diameter of the shadow by a  $\frac{1}{60}$  part; or, which amounts to the same, to add as many seconds as the semi-diameter contains minutes.

413. It is remarked in total eclipses of the sun, that the moon is not wholly invisible, but appears with a dull reddish light.

This phenomenon is doubtless another effect of the earth's atmosphere, though of a totally different nature from the preceding. Certain of the sun's rays, instead of being stopped and absorbed, are bent from their rectilinear course by the refracting power of the atmosphere, so as to form a cone of faint light, interior to that cone which has been mathematically described as the earth's shadow, which falling upon the moon renders it visible.

414. As an eclipse of the moon is occasioned by a real loss of its light, it must begin and end at the same instant, and present precisely the same appearance, to every spectator who sees the moon above his horizon during the eclipse. It will be shown that the case is different with eclipses of the sun.

#### *Calculation of an Eclipse of the Moon.*

415. The apparent distance of the centre of the moon from the axis of the earth's shadow, and the arcs passed over by the



centre of the moon and the axis of the shadow during an eclipse of the moon, being necessarily small, they may without material error be considered as right lines. We may also consider the apparent motion of the sun in longitude, and the motions of the moon in longitude and latitude, as uniform during the eclipse. These suppositions being made, the calculation of the circumstances of an eclipse of the moon is very simple.

Let  $NF$  (Fig. 55) be a part of the ecliptic,  $N$  the moon's ascending node,  $NL$  a part of the moon's orbit,  $C$  the centre of a section of the earth's shadow at the moon,  $CK$  perpendicular to  $NF$  a circle of latitude, and  $C'$  the centre of the moon at the instant of opposition: then  $CC'$ , which is the latitude of the moon, in opposition, is the distance of the centres of the shadow and moon at that time. The moon and shadow both have a motion, and in the same direction, as from  $N$  towards  $F$  and  $L$ . It is the practice, however, to regard the shadow as stationary, and to attribute to the moon a motion equal to the relative motion of the moon and shadow. The orbit that would be described by the moon's centre if it had such a motion, is called the *Relative Orbit* of the moon. Inasmuch as the circumstances of the eclipse depend altogether upon the relative motion of the moon and shadow, this mode of proceeding is obviously allowable.

As the shadow has no motion in latitude, the relative motion of the moon and shadow in latitude will be equal to the moon's actual motion in latitude. And since the centre of the earth's shadow moves in the plane of the ecliptic at the same rate as the sun, the relative motion of the moon and shadow in longitude will be equal to the difference between the motions of the sun and moon in longitude. We obtain, therefore, the relative position of the centres of the moon and shadow at any interval  $t$ , following opposition, by laying off  $Cm$  equal to the difference of the motions of the sun and moon in longitude in this interval, through  $m$  drawing  $mM$  perpendicular to  $NF$ , and cutting off  $mM$  equal to the latitude at opposition plus the motion in latitude in the interval  $t$ :  $M$  will be the position of the moon's centre in the relative orbit, the centre of the shadow being supposed to be stationary at  $C$ . As the motion of the sun in longitude, and of the moon in longitude and latitude, is considered uniform, the ratio of  $C'm' (= Cm$ , the difference between the motions of

the sun and moon in longitude,) to  $M m'$  the moon's motion in latitude, is the same, whatever may be the length of the interval considered. It follows, therefore, that the relative orbit of the moon  $N' C' M$  is a *right line*.

416. The relative orbit passes through  $C'$ , the place of the moon's centre at opposition: its position will therefore be known, if its inclination to the ecliptic be found. Now, we have,

$$\tan \text{inclina.} = \frac{M m'}{C' m'} = \frac{\text{moon's motion in latitude}}{\text{moon's mot. in long.} - \text{sun's mot. in long.}}$$

417. The following data are requisite in the calculation of the circumstances of a lunar eclipse :

$T$  = time of opposition.

$M$  = moon's hourly motion in longitude.

$n$  = moon's hourly motion in latitude.

$m$  = sun's hourly motion in longitude.

$\lambda$  = moon's latitude at opposition.

$d$  = moon's semi-diameter.

$\delta$  = sun's semi-diameter.

$P$  = moon's horizontal parallax.

$p$  = sun's horizontal parallax.

$s$  = semi-diameter of earth's shadow.

$I$  = inclination of relative orbit.

$h$  = moon's hourly motion on relative orbit.

$T, M, n, m, \lambda, d, \delta, P,$  and  $p$ , are derived from Tables of the sun and moon. (See Problems IX and XIV.)

The quantities  $s, I$  and  $h$ , may be determined from these,

$$s = P + p - \delta + \frac{1}{6} (P + p - \delta) \text{ (Arts. 406 and 412) } \dots (89);$$

$$\tan g I = \frac{n}{M - m} \text{ (Art. 416) } \dots (90).$$

The triangle  $C' M m'$  gives,

$$C' M = \frac{C' m'}{\cos I}, \text{ or, } h = \frac{M - m}{\cos I} \dots (91).$$

418. The above quantities being supposed to be known, let  $N' C F$  (Fig. 56) represent the ecliptic, and  $C$  the stationary centre of the earth's shadow. Let  $C C' = \lambda$ , and let  $N' C' L'$  represent the relative orbit of the moon. We here suppose the moon to be

north of the ecliptic at the time of opposition, and near its ascending node; when it is south of the ecliptic,  $\lambda$  is to be laid off below  $N' C F$ , and when it is approaching either node, the relative orbit is inclined to the right. Let the circle  $K F K' R$ , described about the centre  $C$ , represent the section of the earth's shadow at the moon; and let  $f, f'$  and  $g, g'$  be the respective places of the moon's centre, at the beginning and end of the eclipse, and at the beginning and end of the total eclipse.  $Cf = Cf' = s + d$ , and  $Cg = Cg' = s - d$ . Draw  $CM$  perpendicular to  $N' C' L'$ , and  $M$  will represent the place of the moon's centre when nearest the centre of the shadow: it will also be its place at the middle of the eclipse; for, since  $Cf = Cf'$ , and  $CM$  is perpendicular to  $N' C' f'$ ,  $C'f = C'f'$ .

#### 419. *Middle of the eclipse.*

The time of opposition being known, that of the middle of the eclipse will become known when we have found the interval ( $x$ ) employed by the moon in passing from  $M$  to  $C'$ . Now,

$$(\text{expressed in parts of an hour}) \ x = \frac{M C'}{h};$$

and in the right angled triangle  $CC'M$ , we have  $CC' = \lambda$ , and  $\angle C'CM = \angle C'N'C = I$ ; and therefore  $MC' = \lambda \sin I$ ; whence, by substitution,

$$x = \frac{\lambda \sin I}{h} = \frac{\lambda \sin I}{\frac{M - m}{\cos I}} \text{ (equa. 91)} = \frac{\lambda \sin I \cos I}{M - m};$$

$$\text{or, (expressed in seconds)} \ x = \frac{3600s. \cos I}{M - m} \cdot \lambda \sin I \dots (92).$$

Hence, if  $M$  = time of middle, we have,

$$M = T \pm x = T \mp \frac{3600s. \cos I}{M - m} \lambda \sin I \dots (93).$$

It is obvious that the *upper* sign is to be used when the latitude is *increasing*, and the *lower* sign when it is *decreasing*.

The distance of the centre of the moon from the centre of the shadow at the middle of the eclipse,

$$= CM = CC' \cos C'CM = \lambda \cos I \dots (94).$$

#### 420. *Beginning and end of the eclipse.*

Let any point  $l$  of the relative orbit be the place of the moon's centre at the time of any given phase of the eclipse. Let  $t$  = the

interval of time between the given phase and the middle ; and  $k = C l$ , the distance of the centres of the moon and shadow. In the interval  $t$  the moon's centre will pass over the distance  $M l$  ; hence,

$$t = \frac{M l}{h} = \frac{M l \cos I}{M - m} ;$$

but,  $M l = \sqrt{C l^2 - C M^2} = \sqrt{k^2 - \lambda^2 \cos^2 I}$  (equa. 94),

and therefore,  $t = \frac{\cos I}{M - m} \sqrt{k^2 - \lambda^2 \cos^2 I}$  ;

or, (in seconds)  $t = \frac{3600s. \cos I}{M - m} \sqrt{(k + \lambda \cos I)(k - \lambda \cos I)} \dots (95)$ .

Let  $T'$  denote the time of the supposed phase of the eclipse, and  $M$  the time of the middle ; and we shall have,

$$T' = M + t, \text{ or } T' = M - t,$$

according as the phase follows or precedes the middle.

Now at the beginning and end of the eclipse, we have,

$$k = C f \text{ or } C f' = s + d ;$$

substituting in equation (95) we obtain,

$$t' = \frac{3600s. \cos I}{M - m} \sqrt{(s + d + \lambda \cos I)(s + d - \lambda \cos I)} \dots (96) ;$$

$t'$  being found, the time of the beginning (B), and the time of the end (E), result from the equations,

$$B = M - t', E = M + t'.$$

#### 421. *Beginning and end of the total eclipse.*

At the beginning and end of the total eclipse,  $k = C g = C g' = s - d$  ; whence by equation (95),

$$t'' = \frac{3600s. \cos I}{M - m} \sqrt{(s - d + \lambda \cos I)(s - d - \lambda \cos I)} \dots (97) ;$$

and, denoting the time of the beginning by  $B'$  and the time of the end by  $E'$ , we have,

$$B' = M - t'', E' = M + t''.$$

#### 422. *Quantity of the eclipse.*

In a partial eclipse of the moon the magnitude or quantity of the eclipse is measured by the relative portion of that diameter of the moon, which, if produced, would pass through the centre of the earth's shadow, that is involved in the shadow. The

whole diameter is divided into twelve equal parts, called *Digits*, and the quantity is expressed by the number of digits and fractions of a digit in the part immersed. When the moon passes entirely within the shadow, as in a total eclipse, the quantity of the eclipse is expressed by the number of digits contained in the part of the same diameter prolonged outwards, which is comprised between the edge of the shadow and the inner edge of the moon. Thus the number of digits contained in  $SN$  (Fig. 56), expresses the quantity of the eclipse represented in the figure. Hence, if  $Q$  = the quantity of the eclipse, we shall have,

$$Q = \frac{NS}{\frac{1}{12} NV} = \frac{12 NS}{NV} = \frac{12 (NM + MS)}{NV} = \frac{12 (NM + CS - CM)}{NV} = \frac{12 (d + s - \lambda \cos I)}{2d};$$

or,

$$Q = \frac{6 (s + d - \lambda \cos I)}{d} \dots (98).$$

If  $\lambda \cos I$  exceeds  $(s + d)$  there will be no eclipse. If it is intermediate between  $(s + d)$  and  $(s - d)$  there will be a partial eclipse; and if it is less than  $(s - d)$  the eclipse will be total.

#### *Construction of an Eclipse of the Moon.*

423. The times of the different phases of an eclipse of the moon may easily be determined by a geometrical construction, within a minute or two of the truth. Draw a right line  $N'F$  (Fig. 57) to represent the ecliptic; and assume upon it any point  $C$ , for the position of the centre of the earth's shadow at the time of opposition. Then, having fixed upon a scale of equal parts, lay off  $CR = M - m$  the difference of the hourly motions of the sun and moon in longitude; and draw the perpendiculars  $CC' = \lambda$  the moon's latitude in opposition, and  $RL' = \lambda \pm n$  the moon's latitude an hour after opposition. The right line  $C'L'$  drawn through  $C'$  and  $L'$ , will represent the moon's relative orbit. It should be observed, that if the latitudes are south, they must be laid off below  $N'F$ , and that  $N'C'L'$  will be inclined to the right when the latitude is decreasing. With a radius  $CR' = s$  (equation 89) describe the circle  $EKF$   $K'$ , which will represent the section of the earth's shadow. With a radius  $= s + d$ , and another radius  $= s - d$ , describe about the centre  $C$  arcs intersecting  $N'L'$  in  $f, f'$ , and  $g, g'$ ;  $f$  and  $f'$  will



be the places of the moon's centre at the beginning and end of the eclipse, and  $g$  and  $g'$  the places at the beginning and end of the total eclipse. From the point  $C$  let fall upon  $N' C' L'$  the perpendicular  $C M$ ; and  $M$  will be the place of the moon's centre at the middle of the eclipse. To render the construction explicit, let us suppose the time of opposition to be 7h. 23m. 15s. At this time the moon's centre will be at  $C'$ . To find its place at 7h., state the proportion, 60m. : 23m. 15s. :: moon's hourly motion on the relative orbit : a fourth term. This fourth term will be the distance of the moon's centre from the point  $C'$  at 7 o'clock, and if it be taken in the dividers and laid off on the relative orbit from  $C'$  backwards to the point 7, it will give the moon's place at that hour. This being found, take in the dividers the moon's hourly motion on the relative orbit, and lay it off repeatedly, both forwards and backwards from the point 7, and the points marked off, 8, 9, 10, 6, 5, will be the moon's places at those hours respectively. Now, the object being to find the times at which the moon's centre is at the points  $f, f', g, g'$ , and  $M$ , let the hour spaces thus found be divided into quarters, and these subdivided into 5 minute or minute spaces, and the times answering to the points of division that fall nearest to these points, will be within a minute or so of the times in question. For example, the point  $f'$  falls between 9 and 10, and thus the end of the eclipse will occur somewhere between 9 and 10 o'clock. To find the number of minutes after 9 at which it takes place, we have only to divide the space from 9 to 10 into four equal parts or 15 minute spaces, subdivide the part which contains  $f$  into three equal parts or 5 minute spaces, and again that one of these smaller parts within which  $f'$  lies, into five equal parts or minute spaces.

### *Eclipses of the Sun.*

424. An eclipse of the sun is caused by the interposition of the moon between the sun and earth; whereby the whole, or part of the sun's light, is prevented from falling upon certain parts of the earth's surface.

Let  $A G B$  and  $a g b$  (Fig. 58) be sections of the sun and earth by a plane passing through their centres  $S$  and  $E$ ,  $A a$ ,  $B b$  tangents to the circles  $A G B$  and  $a g b$  on the same side, and  $A d$ ,  $B c$  tangents to the same on opposite sides. The figure  $A a b B$  will

be a section through the axis, of a frustum of a cone formed by rays tangent to the sun and earth on the same side, and the triangular space  $F'cd$  will be a section of a cone formed by rays tangent on opposite sides. An eclipse of the sun will take place somewhere upon the earth's surface, whenever the sun comes within the frustum  $AabB$ , and a total or an annular eclipse whenever the sun comes within the cone  $F'cd$ .

425. Let  $mm'M$  (Fig. 58) be a circular arc described about the centre  $E$ , and with a radius equal to the distance of the centres of the moon and earth at the time of conjunction. The angle  $mES$  is the apparent semi-diameter of a section of the frustum, and  $m'ES$  the apparent semi-diameter of a section of the cone at the distance of the moon. To find expressions for these semi-diameters in terms of determinate quantities, let the first be denoted by  $A$ , and the second by  $A'$ ; and let  $P$  = the parallax of the moon,  $p$  = the parallax of the sun, and  $\delta$  = the semi-diameter of the sun. Then we have,

$$mES = A = mEA + AES = Ema - EAm + AES;$$

$$\text{or,} \quad A = P - p + \delta \dots (99).$$

$$\text{and} \quad m'ES = m'EB - BES = Em'c - EBm' - BES;$$

$$\text{or,} \quad A' = P - p - \delta \dots (100).$$

Taking the mean values of  $P$ ,  $p$ , and  $\delta$  (Art. 407), we find for the mean value of  $A$   $1^\circ 12' 53''$ , and for the mean value of  $A'$   $40' 51''$ .

426. As the plane of the moon's orbit is not coincident with the plane of the ecliptic, an eclipse of the sun can happen only when conjunction or new moon takes place in one of the nodes of the moon's orbit, or so near it that the moon's latitude does not exceed the sum of the semi-diameters of the moon and of the luminous frustum (Art. 425) at the moon's orbit. Thus, denoting the moon's semi-diameter by  $d$ , and the greatest latitude of the moon in conjunction, at which an eclipse can take place, by  $L$ , we have,

$$L = P - p + \delta + d \dots (101).$$

For a total eclipse, the greatest latitude will be equal to the sum of the semi-diameters of the moon and the luminous cone. Hence, denoting it by  $L'$ ,

$$L' = P - p - \delta + d \dots (102).$$

In order that an annular eclipse may take place, the apparent

semi-diameter of the moon must be less than that of the sun, and the moon must come at conjunction entirely within the luminous frustum. Whence, if  $L''$  = the maximum latitude at which an annular eclipse is possible, we have,

$$L'' = P - p + \delta - d \dots (103).$$

427. In the same manner as in the case of an eclipse of the moon, it has been found that when at the time of mean new moon the difference of the mean longitudes of the sun or moon and of the node, exceeds  $19^{\circ} 44'$ , there cannot be an eclipse of the sun ; but when the difference is less than  $13^{\circ} 33'$ , there must be one. These numbers are called the *Solar Ecliptic Limits*.

428. In order to discover at what new moons in the course of a year an eclipse of the sun will happen, with its approximate time, we have only to find the mean longitudes of the sun and node at each mean new moon throughout the year (Art. 385), and take the difference of the longitudes, and compare it with the solar ecliptic limits. (For a more direct method of solving this problem, see Prob. XXVIII).

429. Eclipses both of the sun and moon recur in nearly the same order and at the same intervals at the expiration of a period of 223 lunations, or 18 years of 365 days, and 15 days,\* which for this reason is called the *Period of the Eclipses*. For, the time of a revolution of the sun with respect to the moon's node is 346.619851d., and the time of a synodic revolution of the moon is 29.5305887d. These numbers are very nearly in the ratio of 223 to 19. Thus, in a period of 223 lunations the sun will have returned 19 times to the same position with respect to the moon's node, and at the expiration of this period, will be in the same position with respect to the moon and node, as at its commencement. The eclipses which occur during one such period being noted, subsequent eclipses are easily predicted.

This period was known to the Chaldeans and Egyptians, by whom it was called *Saros*.

430. As the solar ecliptic limits are more extended than the lunar, eclipses of the sun must occur more frequently than eclipses of the moon.

\* More exactly 18 years (of 365 days) plus 15d. 7h. 42m. 29s.

As to the *number of eclipses* of both luminaries, there cannot be fewer than two, nor more than seven in one year. The most usual number is four, and it is rare to have more than six. When there are seven eclipses in a year, five are of the sun and two of the moon; and when but two, both are of the sun. The reason is obvious. The sun passes by both nodes of the moon's orbit but once in a year, unless he passes by one of them in the beginning of the year, in which case he will pass by the same again a little before the end of the year, as he returns to the same node in a period of 346 days. Now, if the sun be at a little less distance than  $13^{\circ} 33'$  from either node at the time of new moon, he will be eclipsed, and at the subsequent opposition the moon will be eclipsed near the other node, and come round to the next conjunction before the sun is  $13^{\circ} 33'$  from the former node. And when three eclipses happen about either node, the like number commonly happens about the opposite one; as the sun comes to it in 173 days afterward, and six lunations contain only four days more. Thus there may be two eclipses of the sun and one of the moon about each of the nodes; and the twelfth lunation from the eclipse in the beginning of the year, may give a new moon before the year is ended, which, in consequence of the retrogradation of the nodes, may be within the solar ecliptic limit; and hence there may be seven eclipses in a year, five of the sun and two of the moon. But when the moon changes in either of the nodes, she cannot be near enough the other node, at the next full moon, to be eclipsed, as in the interval the sun will move over an arc of  $14^{\circ} 32'$ , whereas the greatest lunar ecliptic limit is but  $13^{\circ} 21'$ , and in six lunar months afterwards she will change near the other node; in this case there cannot be more than two eclipses in a year, both of which will be of the sun. If the moon changes at the distance of a few degrees from either node, then an eclipse both of the sun and moon will probably occur in the passage of that node and also of the other.

431. Although solar eclipses are more frequent than lunar, when considered with respect to the whole earth, yet at any given place more lunar than solar eclipses are seen. The reason of this circumstance, is that an eclipse of the sun (unlike an eclipse of the moon) is visible only over a part of a hemisphere

of the earth. To show this, suppose two lines to be drawn from the centre of the moon tangent to the earth at opposite points : they will make an angle with each other equal to double the moon's horizontal parallax, or of  $1^{\circ} 54'$ . Therefore, should an observer situated at one of the points of tangency, refer the centre of the moon to the centre of the sun, an observer at the other would see the centres of these bodies distant from each other an angle of  $1^{\circ} 54'$ , and their nearest limbs separated by an arc of more than  $1^{\circ}$ .

432. Instead of regarding an eclipse of the sun as produced by an interposition of the moon between the sun and earth, as we have hitherto considered it, we may regard it as occasioned by the moon's shadow falling upon the earth. Fig. 59 represents the moon's shadow as projected from the sun and covering a portion of the earth's surface. Wherever the umbra falls, there is a total eclipse ; and wherever the penumbra falls, a partial eclipse.

433. In order to discover the extent of the portion of the earth's surface over which the eclipse is visible at any particular time, we have only to find the breadth of the portion of the earth covered by the penumbral shadow of the moon ; but we will first ascertain the length of the moon's shadow. As seen at the vertex of the moon's shadow, the apparent diameters of the moon and sun are equal. Now, as seen at the centre of the earth, they are nearly equal, sometimes the one being a little greater and sometimes the other. It follows, therefore, that *the length of the moon's shadow is about equal to the distance of the earth, being sometimes a little greater and at other times a little less.*

When the apparent diameter of the moon is the greater, the shadow will extend beyond the earth's centre ; and when the apparent diameter of the sun is the greater, it will fall short of it. If we increase the mean apparent diameter of the moon as seen from the earth's centre, viz :  $31' 7''$ , by  $\frac{1}{60}$ , the ratio of the radius of the earth to the distance of the moon, we shall have  $31' 38''$  for the mean apparent diameter of the moon as seen from the nearest point of the earth's surface. Comparing this with the mean apparent diameter of the sun as viewed from the same point, which is sensibly the same as at the centre of the earth, or  $32' 2''$ , we perceive that it is less ; from which we conclude, that



when the sun and moon are each at their mean distance from the earth, the shadow of the moon does not extend as far to the earth's surface.

434. To find a general expression for the length of the moon's shadow, let  $A G B$ ,  $a' g' b'$ , and  $a g b$  (Fig. 60), be sections of the sun, moon, and earth, by a plane passing through their centres  $S$ ,  $M$ , and  $E$ , supposed to be in the same right line, and  $A a'$ ,  $B b'$  tangents to the circles  $A G B$ ,  $a' g' b'$  : then  $a' K b'$  will represent the moon's shadow. Let  $L$  = the length of the shadow ;  $D$  = the distance of the moon ;  $D'$  = the distance of the sun ;  $d$  = apparent semi-diameter of the moon ; and  $\delta$  = apparent semi-diameter of the sun. At  $K$  the vertex of the shadow,  $M K a'$  the apparent semi-diameter of the moon, will be equal to  $S K A$  the apparent semi-diameter of the sun ; and as the distance of this point from the centre of the earth, even when it is the greatest, is small in comparison with the distance of the sun (Art. 433), the apparent semi-diameter of the sun will always be very nearly the same to an observer situated at  $K$  as to one situated at the centre of the earth. Now, since the apparent semi-diameter of the moon is inversely proportional to its distance,

$$\text{angle } M K a' : d :: M E : M K ;$$

and thus,  $\delta : d :: M E : M K :: D : L$  (nearly) ;

whence, 
$$L = D \frac{d}{\delta} \dots (104).$$

If a more accurate result be desired, we have only to repeat the calculations, after having diminished  $\delta$  in the ratio of  $D$  to  $(D + L)$ .

435. Now, to find the breadth of the portion of the earth's surface, covered by the penumbral shadow, let the lines  $A d'$ ,  $B c'$  (Fig. 60) be drawn tangent to the circles  $A G B$ ,  $a' g' b'$ , on opposite sides, and prolonged on to the earth. The space  $h c' d' k$  will represent the penumbra of the moon's shadow, and the arc  $g h$  one half the breadth of the portion of the earth's surface covered by it. Let this arc or the angle  $g E h = S$ , and denote the semi-diameter of the sun, and the semi-diameter and parallax of the moon, by the same letters as in previous Articles. The triangle  $M E h$  gives,

$$\text{angle } M E h = S = M h Z - h M E.$$

The angle  $h M E$  is the moon's parallax in altitude at the station  $h$ , and  $M h Z$  is its zenith distance at the same station. Denote the former by  $P'$  and the latter by  $Z$ . Thus,

$$S = Z - P' \dots (105).$$

The triangle  $h M S$  gives,

$$h M E = P' = M S h + M h S;$$

$M h S = d + \delta$ ; and  $M S h$  is the sun's parallax in altitude at the station  $h$ ; let it be denoted by  $p'$ . We have then,

$$P' = d + \delta + p' = d + \delta \text{ (nearly)} \dots (106);$$

and to find  $Z$  we have (equa. 11, p. 43),

$$P' = P \sin Z, \text{ or } \sin Z = \frac{P'}{P} \dots (107).$$

$P'$  and  $Z$  being found by these equations, equa. (105) will then make known the value of  $S$ .

If great accuracy is required, the calculation must be repeated, giving now to  $p'$  in equation (106) the value furnished by equation (11) which expresses the relation between the parallax in altitude of a body and its horizontal parallax, instead of neglecting it as before; and  $Z$  must be computed from the following equation:

$$\sin Z = \frac{\sin P'}{\sin P} \dots (108).$$

The penumbral shadow will obviously attain to its greatest breadth, when the sun is in its perigee and the moon is in its apogee. The values of  $d$ ,  $\delta$  and  $P$  under these circumstances are respectively  $14' 41''$ ,  $16' 18''$ , and  $53' 48''$ . Performing the calculations, we find that *the breadth of the greatest portion of the earth's surface ever covered by the penumbral shadow, is  $69^\circ 17'$ , or about 4800 miles.*

436. The breadth of the spot comprehended within the umbra, may be found in a similar manner. The arc  $g h'$  (Fig. 60) represents one half of it: denote this arc or the equal angle  $g E h'$  by  $S'$ ,

$$M E h' = S' = M h' Z' - h' M E;$$

or,

$$S' = Z - P' \dots (109).$$

$$h' M E = P' = M S h' + M h' S;$$

but  $M h' S = d - \delta$ , and  $M S h' = p'$ , sun's parallax in altitude at  $h'$ ; whence,

$$P' = d - \delta + p' = d - \delta \text{ (nearly) } \dots (110);$$

and we have, as before,

$$P' = P \sin Z, \text{ or } \sin Z = \frac{P'}{P} \dots (111).$$

The greatest breadth will obtain when the sun is in its apogee and the moon is in its perigee. We shall then have,

$$\delta = 15' 45'', d = 16' 45'', P = 61' 24''.$$

Making use of these numbers, we deduce for the *maximum breadth of the portion of the earth's surface covered by the moon's shadow*,  $1^\circ 50'$ , or 127 miles.

437. It should be observed that the deductions of the last two articles answer to the supposition that the moon is in the node, and that the axis of the shadow and penumbra passes through the centre of the earth. In every other case, both the shadow and penumbra will be cut obliquely by the earth's surface, and the sections will be ovals, and very nearly true ellipses, the lengths of which may materially exceed the above determinations.

438. Parallax not only causes the eclipse to be visible at some places and invisible at others, as shown in Art. 431; but, by making the distance of the centres of the sun and moon unequal, renders the circumstances of the eclipse at those places where it is visible, different at each place. This may also be inferred from the circumstance that the different places, covered at any time by the shadow of the moon, will be differently situated within this shadow. It will be seen, therefore, that an eclipse of the sun has to be considered in two points of view: 1st. *With respect to the whole earth*, or as a *general eclipse*; and 2d. *With respect to a particular place*.

439. The following are the principal facts relative to eclipses of the sun that remain to be noticed. 1st. The duration of a general eclipse of the sun cannot exceed about 6 hours. 2d. A solar eclipse does not happen at the same time at all places where it is seen; as the motion of the moon beyond the sun, and consequently of its shadow, is from west to east, the eclipse must begin *earlier* at the *western* parts and *later* at the *eastern*.

3d. The moon's shadow being tangent to the earth, at the commencement and end of the eclipse, the sun will be just rising at the place where the eclipse is first seen, and just setting at the place where it is last seen. At the intermediate places, the sun will at the time of the beginning and end of the eclipse have various altitudes. 4th. An eclipse of the sun begins on the *western* side and ends on the *eastern*. 5th. When the straight line passing through the centres of the sun and moon, passes also through the place of the spectator, the eclipse is said to be *central*: a central eclipse may be either annular or total, according as the apparent diameter of the sun is greater than that of the moon, or the reverse. 6th. A total eclipse of the sun cannot last at any one place more than *eight minutes*; and an annular eclipse more than *twelve and a half minutes*. 7th. In most solar eclipses the moon's disc is covered with a faint light, a phenomenon which is attributed to the reflection of the light from the illuminated part of the earth.

*Calculation of an Eclipse of the Sun.*

1. *Of the circumstances of the general eclipse.*

440. It is a simple inference from what has been established in Art. 426, that an eclipse of the sun will begin and end upon the earth, at the times before and after conjunction, when the distance of the centres of the moon and sun is equal to  $P - p + \delta + d$ ; that the total eclipse will begin and end when this distance is equal to  $P - p - \delta + d$ ; and the annular eclipse when the distance is equal to  $P - p + \delta - d$ .

441. The times of the various phases of the general eclipse of the sun, may be obtained by a process precisely analogous to that by which the times of those of an eclipse of the moon are found. Let C (Fig. 61) be the centre of the sun, and C' the centre of the moon, at the time of conjunction. We may suppose the sun to remain stationary at C, if we attribute to the moon a motion equal to its motion relative to the sun; for, on this supposition, the distance of the centres of the two bodies will, at any given period during the eclipse, be the same as that which obtains in the actual state of the case. Let N' C' L' represent the orbit that would be described by the moon if it had such a motion, which is called the *Relative Orbit*. Let C M be drawn perpendicular to it; and let  $Cf = C'f' = P - p + \delta + d$ ,

and  $Cg = Cg' = P - p - \delta + d$ , or  $P - p + \delta - d$ , according as the eclipse is total or annular. Then,  $M$  will be the place of the moon's centre at the middle of the eclipse;  $f$  and  $f'$  the places at the beginning and end of the eclipse; and  $g$  and  $g'$  the places at the beginning and end of the total, or of the annular eclipse. We shall thus have, as in eclipses of the moon,

$$\text{tang } I = \frac{n}{M - m}, \quad CM = \lambda \cos I, \quad C'M = \lambda \sin I \dots (112).$$

$$\text{Interval from con. to mid.} = \frac{3600s. \lambda \sin I \cos I}{M - m} \dots (113).$$

Interval from middle to beginning or end

$$= \frac{3600s \cos I}{M - m} \sqrt{(k' + \lambda \cos I)(k' - \lambda \cos I)} \dots (114).$$

Interval for total eclipse

$$= \frac{3600s. \cos I}{M - m} \sqrt{(k'' + \lambda \cos I)(k'' - \lambda \cos I)} \dots (115).$$

Interval for annular eclipse

$$= \frac{3600s. \cos I}{M - m} \sqrt{(k''' + \lambda \cos I)(k''' - \lambda \cos I)} \dots (116).$$

$$\text{Quantity} = \frac{6(k' - \lambda \cos I)}{d} \dots (117).$$

$$k' = P - p + \delta + d, \quad k'' = P - p - \delta + d, \quad k''' = P - p + \delta - d \quad (118).$$

The letters  $\lambda$ ,  $M$ ,  $m$ , &c. represent quantities of the same name as in the formulæ for a lunar eclipse; but they designate the values of these quantities at the time of *conjunction*, instead of opposition. These values are in practice obtained from tables of the sun and moon, as in a lunar eclipse.

442. The times of the different circumstances of a general eclipse of the sun, may also be found within a minute or two of the truth, by *construction*, in a precisely similar manner with those of an eclipse of the moon (Art. 423).

## 2. Of the phases of the eclipse at a particular place.

443. The phase of the eclipse, which obtains at any instant at a given place, is indicated by the relation between the apparent distance of the centres of the sun and moon and the sum of their apparent semi-diameters. And the calculation of the time



of any given phase of the eclipse, consists in the calculation of the time when the apparent distance of the centres has the value relative to the sum of the semi-diameters, answering to the given phase. Thus, if we wish to find the time of the beginning of the eclipse, we have to seek the time when the apparent distance of the centres of the sun and moon first becomes equal to the sum of their apparent semi-diameters.

444. The calculation of the different phases of an eclipse of the sun, for a particular place, involves, then, the determination of the apparent distance of the centres of the sun and moon, and of the apparent semi-diameters of the two bodies at certain stated periods.

The true semi-diameter of the sun, as given by the tables, may be taken for the apparent without material error. For the method of computing the apparent semi-diameter of the moon, for any given time and place, see Problem XXVII.

445. According to the celebrated astronomer Duséjour, in order to make the observations agree with theory, it is necessary to diminish the sun's semi-diameter, as it is given by the tables,  $3''.5$ . This circumstance is explained, by supposing that the apparent diameter of the sun is amplified, by reason of the very lively impression which its light makes upon the eye. This amplification is called *Irradiation*. He also thinks that the semi-diameter of the moon ought to be diminished  $2''$ , to make allowance for an *Inflexion* of the light which passes near the border of this luminary, supposed to be produced by its atmosphere. It must be observed, however, that the astronomers of the present day do not agree either as to the necessity or the amount of the diminutions just spoken of.

446. The determination of the apparent distance of the centres of the sun and moon may easily be accomplished, as will be shown in the sequel, when the apparent longitude and latitude of the two bodies have been found. Now, the true longitude of the sun, and the true longitude and latitude of the moon, may be found from the tables (Probs. IX and XIV); and from these the apparent longitudes and latitudes may be deduced, by correcting for the parallax. But the customary mode of proceeding is a little different from this: the true longitude and latitude of the sun are employed instead of the appa-

rent, and the parallax of the sun is referred to the moon; that is, the difference between the parallax of the moon and that of the sun is, by fiction, taken as the parallax of the moon. This supposititious parallax is called the moon's *Relative Parallax*. (See Prob. XVII.)

447. We will first show how to find the *approximate times* of the different phases of the eclipse. Put  $T$  = the time of new moon known to within 5 or 10 minutes. (Prob. XXVII.) For the time  $T$  calculate by the tables the sun's longitude, hourly motion, and semi-diameter, and the moon's longitude, latitude, horizontal parallax, semi-diameter, and hourly motions in longitude and latitude. Subtract the sun's horizontal parallax from the reduced horizontal parallax of the moon,\* and calculate the apparent longitude and latitude, and the apparent semi-diameter of the moon. From a comparison of the apparent longitude of the moon with the true longitude of the sun, we shall know whether apparent ecliptic conjunction occurs before or after the time  $T$ . Let  $T'$  denote the time an hour earlier or later than the time  $T$ , according as the apparent conjunction is earlier or later. With the sun and moon's longitudes, the moon's latitude, and the hourly motions in longitude and latitude at the time  $T$ , calculate the longitudes and the moon's latitude for the time  $T'$ ; and for this time also calculate the moon's apparent longitude and latitude. Take the difference between the apparent longitude of the moon and the true longitude of the sun at the time  $T$ , and it will be the apparent distance of the moon from the sun in longitude, at this time. Let it be denoted by  $n$ . Find, in like manner, the apparent distance of the moon from the sun in longitude at the time  $T'$ , and denote it by  $n'$ . In the same manner as at the time  $T$ , we find whether apparent conjunction occurs before or after the time  $T'$ . If it occurs between the times  $T$  and  $T'$ , the sum of  $n$  and  $n'$ , otherwise their difference, will be the apparent relative motion of the sun and moon in longitude in the interval  $T' - T$ , or  $T - T'$ ; from which the relative hourly motion will become known. The difference of the apparent latitudes of the moon, at the times  $T$  and

---

\* The reduced horizontal parallax of the moon is its horizontal parallax as reduced from the equator to the given place. (See Prob. XV.)

$T'$ , will make known the apparent relative hourly motion in latitude. With the relative hourly motion in longitude and the difference of the apparent longitudes at the time  $T$ , find by simple proportion, the interval between the time  $T$  and the time of apparent ecliptic conjunction; and then, with the apparent latitude of the moon at the time  $T$  and its hourly motion in latitude, find the apparent latitude at the time of apparent conjunction thus determined. Then, knowing the relative hourly motion of the sun and moon in longitude and latitude, together with the time of apparent conjunction, and the apparent latitude at that time, and regarding the apparent relative orbit of the moon as a right line, (which it is nearly,) it is plain that the time of beginning, greatest obscuration, and end, as well as the quantity of the eclipse, may be calculated after the same manner as in the general eclipse; the disc of the sun answering to the section of the luminous frustum mentioned in Art. 424, and the apparent elements answering to the true. Let  $C$  (Fig. 62) represent the centre of the sun supposed stationary,  $C C'$  the apparent latitude of the moon at apparent conjunction,  $N' C'$  the apparent relative orbit of the moon, determined by its passing through the point  $C'$ , and making a determinate angle with the ecliptic  $N' F$ , or by its passing through the situations of the moon at the times  $T$  and  $T'$ . Also, let  $R K F K'$  be the border of the sun's disc;  $f, f'$  the positions of the moon's centre at the beginning and end of the eclipse, determined by describing a circle around  $C$  as a centre, with a radius equal to the sum of the apparent semi-diameters of the sun and moon; and  $M$  (the foot of the perpendicular let fall from  $C$  upon  $N' C'$ ) its position at the time of greatest obscuration.

If the eclipse should be total or annular, then  $g, g'$  will be the positions of the moon's centre at the beginning and end of the total or annular eclipse; these points being determined by describing a circle around  $C$  as a centre, and with a radius equal to the difference of the apparent semi-diameters of the sun and moon.

The various circumstances of the eclipse may also be had by construction, after the same manner as in a lunar eclipse (Art. 423).

448. In order to be able to observe the beginning or end of a

solar eclipse, it is necessary to know the position of the point on the sun's limb where the first or last contact takes place. The situation of these points is designated by the distance on the limb, intercepted between them and the highest point of the limb, called the *Vertex*. The contacts will take place at the points  $t, t'$  (Fig. 62), on the lines  $Cf, Cf'$ . To find the position of the vertex, with the sun's longitude found for the beginning of the eclipse calculate the angle of position of the sun at that time,\* and lay it off to the right of the circle of latitude  $CK$  when the sun's longitude is between  $90^\circ$  and  $270^\circ$ , and to the left when the longitude is less than  $90^\circ$  or more than  $270^\circ$ . Suppose  $CP$  to be the circle of declination thus determined. Next, let  $Z$  (Fig. 18) be the zenith,  $P$  the elevated pole, and  $S$  the sun; then in the triangle  $ZPS$  we shall know  $ZP$  the co-latitude,  $\angle ZPS$  the hour angle of the sun, and we may deduce  $PS$  the co-declination of the sun, from the longitude of the sun as derived from the tables, (equa. 36). These three quantities being known,  $\angle ZSP$  the angle made by the vertical through the sun with its circle of declination, may be computed; and being laid off in the figure to the right or left of  $CP$  (Fig. 62), according as the time of beginning is before or after noon, the point  $Z$  or  $Z'$ , as the case may be, in which the vertical intersects the limb  $RKK'$ , will be the vertex, and the arc  $Zt$ , or  $Z't$  on the limb, will ascertain the situation of  $t$  the first point of contact, with respect to it.

The situation of the last point of contact may be found by the same mode of proceeding.

449. Let us now show how to find the exact times of the beginning, greatest obscuration, and end of the eclipse, the approximate times being known. Let  $B$  designate the approximate time of beginning, taken to the nearest minute. Calculate for the time  $B$  by means of the tables, the sun's longitude, hourly motion, and semi-diameter; also the moon's longitude, latitude, horizontal parallax, semi-diameter, and hourly motions in longitude and latitude. Then, making use of the relative parallax, calculate the apparent longitude, latitude, and semi-diameter of the moon.

---

\* A formula which makes known the angle of position of the sun, when the longitude of the sun and the obliquity of the ecliptic are given, is investigated in the Appendix. (See Prob. XIII).



Subtract the apparent longitude of the moon from the true longitude of the sun ; the difference will be the apparent distance of the moon from the sun in longitude ; let it be denoted by  $\alpha$ . Denote the apparent latitude of the moon by  $\lambda$ .

Now, let  $EC$  (Fig. 63) represent an arc of the ecliptic, and  $K$  its pole ; and let  $S$  be the situation of the sun, and  $M$  the apparent situation of the moon at the time  $B$ . Then  $MS$  is the apparent distance of the centres of the two bodies at this time. Denote it by  $\Delta$ .  $S m = \alpha$ , and  $M m = \lambda$ . The right angled triangle  $MSm$  being very small, may be considered as a plane triangle, and we therefore have, to determine  $\Delta$ , the equation,

$$\Delta^2 = \alpha^2 + \lambda^2 \dots (119).^*$$

450. Having computed the value of  $\Delta$ , we find, by comparing it with the sum of the apparent semi-diameters of the sun and moon, whether the beginning of the eclipse occurs before or after the approximate time  $B$ . Fix upon a time some 4 or 5 minutes before or after  $B$ , according as the beginning is before or after, and call it  $B'$ . With the sun and moon's longitudes, the moon's latitude, and the hourly motions in longitude and latitude, at the time  $B$ , find the longitudes and the moon's latitude at the time  $B'$ , and compute for this time the apparent longitude, latitude, and semi-diameter of the moon. Subtract the apparent longitude of the moon from the true longitude of the sun, and we shall have the apparent distance of the moon from the sun at the time  $B'$ . Subtract this from the same distance  $\alpha$  at the time  $B$ , and we shall have the apparent relative motion of the sun and moon in longitude during the interval of time between  $B$  and  $B'$ . Then find, by simple proportion, the apparent relative hourly motion in longitude, and denote it by  $k$ . Take the difference between the apparent latitudes of the moon at the times  $B$  and  $B'$ , and it will be the apparent relative motion of the sun and moon in latitude, in the interval ; from which deduce the apparent relative hourly motion in latitude, and call it  $n$ . Now, put  $t$  = the interval between the

\* In place of equation (119), the following equations may be employed in logarithmic computation :

$$\text{tang } \theta = \frac{\lambda}{\alpha}, \Delta = \frac{\alpha}{\cos \theta} ;$$

where  $\theta$  is an auxiliary arc.



approximate and true times of the beginning of the eclipse, and suppose  $S$  and  $M$  (Fig. 63) to be the situations of the sun and moon at the true time of beginning. In the time  $t$ , the apparent relative motions in longitude and latitude, will be, respectively, equal to  $k t$  and  $n t$ , and accordingly we shall have,

$$S m = \alpha - k t, \quad M m = \lambda + n t.$$

The small right-angled triangle  $S M m$  may be considered as a plane triangle; the hypotenuse  $S M = \downarrow$  = the sum of the apparent semi-diameters of the sun and moon, minus  $5''.5$ . We have then, to find  $t$ , the equation,

$$(\alpha - k t)^2 + (\lambda + n t)^2 = \downarrow^2,$$

or, developing and transposing,

$$(n^2 + k^2) t^2 - 2(\alpha k - \lambda n) t = \downarrow^2 - (\alpha^2 + \lambda^2) = \downarrow^2 - \Delta^2;$$

making  $A = \downarrow^2 - \Delta^2$ , and  $B = \alpha k - \lambda n$ ,

$$(n^2 + k^2) t^2 - 2 B t = A,$$

and, 
$$t = \frac{B - \sqrt{B^2 + A(n^2 + k^2)}}{n^2 + k^2} \dots (120).$$

The negative sign must be prefixed to the radical, for, if we suppose  $A$  to be equal to zero,  $t$  must be equal to zero. Multiplying the numerator and denominator by  $B + \sqrt{B^2 + A(n^2 + k^2)}$ , and restoring the value of  $A$ , we obtain,

$$(\text{in seconds}) t = \frac{3600s. (\downarrow^2 - \Delta^2)}{B + \sqrt{B^2 + (\downarrow^2 - \Delta^2)(n^2 + k^2)}} \dots (121).$$

Although this equation has been investigated for the beginning of the eclipse, it is plain that it will answer equally well for the determination of the other phases, if we give the proper values and signs to  $\downarrow$ ,  $\alpha$ ,  $\lambda$ ,  $n$ , and  $k$ .  $k$  is positive before conjunction and negative after it;  $n$  is negative, when the moon appears to recede from the north pole of the ecliptic;  $\lambda$  is negative, when it is south;  $\alpha$  is always positive.\*

\* Developing the radical in equation (120), and neglecting all the terms after the second, as being very small, we obtain for the beginning and end of the eclipse the more convenient formula,

$$t = \frac{1800s. (\psi^2 - \Delta^2)}{B}.$$

451. The values of the quantities  $\alpha$ ,  $\lambda$ ,  $n$ , and  $k$ , are found for the other phases after the same manner as for the beginning.

To obtain the value of  $\downarrow$  at the time of greatest obscuration, find the relative motions in longitude and latitude ( $k$  and  $n$ ), during some short interval near the middle of the eclipse, which is the approximate time of greatest obscuration; then compute the inclination of the relative orbit by the equation

$$\text{tang } I = \frac{n}{k} \dots (122). \text{ (See equa. 90):}$$

after which  $\downarrow$  will result from the equation

$$\downarrow = \lambda \cos I \dots (123). \text{ (See equa. 94).}$$

For the beginning and end of the total eclipse, we have,  $\downarrow =$  appar. semi-diam. of moon — appar. semi-diam. of sun  $+ 1''.5$ ; and for the beginning and end of the annular eclipse,  $\downarrow =$  appar. semi-diam. of sun, — appar. semi-diam. of moon, —  $1''.5$ .

452. If the value of  $\downarrow$ , given by equation (123), be substituted in equation (121), this equation will make known the time of greatest obscuration; but this may be found more conveniently by a different process. Let N C F (Fig. 64) represent a portion of the ecliptic, E M L a portion of the relative orbit passed over about the time of greatest obscuration, C the stationary position of the sun's centre, and M the place of the moon's centre at the instant of its nearest approach to C. Also, let  $\alpha = C R$  the apparent distance of the moon from the sun in longitude at the time of the nearest approach of the centres,  $\lambda' = R M$  the moon's apparent latitude at the same time,  $k = M k$  the apparent relative motion in longitude in some short interval about this time, and  $n = k n$  the moon's apparent motion in latitude during the same interval. The right angled triangles M  $n k$  and C M R are similar, for their sides are respectively perpendicular to each other; whence,

$$M k : M R :: k n : C R;$$

$$\text{and } C R = M R \frac{k n}{M k}, \text{ or, } \alpha = \lambda' \frac{n}{k} \dots (124).$$

If the moon's apparent latitude be found for the approximate time of greatest obscuration, and substituted for  $\lambda'$  in equation (124), this equation will give very nearly the apparent distance

( $\alpha$ ) of the two bodies in longitude at the true time of greatest obscuration. With this, and the apparent distance at the approximate time of greatest obscuration, together with the relative apparent motion in longitude, the true time of greatest obscuration may be found nearly by simple proportion. A more accurate result may then be had by finding the moon's apparent latitude for the time obtained, substituting it for  $\lambda'$  in equation (124) and then repeating the calculations.

453. A simpler, though less accurate method than that already given, of finding the times of beginning and end of the total or annular eclipse, is to compute the half duration of the total or annular eclipse, and add it to, and subtract it from, the time of greatest obscuration. This interval may easily be determined, if we can find the rate of motion on the relative orbit, and the distance passed over by the moon's centre during the interval. Let  $g, g'$  (Fig. 64) be the places of the moon's centre at the instants of the two interior contacts, and  $M m$  the distance passed over in some short interval ( $L$ ). Let  $\theta = \angle M n k$  the complement of the inclination of the relative orbit,  $k = M k$ ,  $k' = M n$ , and  $t$  = half duration of total or annular eclipse. The triangles  $M n k, C R M$  give,

$$M n = \frac{M k}{\sin M n k}, \text{ or } k' = \frac{k}{\sin \theta} \dots (125),$$

$$\text{and, } \text{tang } R C M = \text{tang } M n k = \frac{R M}{C R}, \text{ or, } \text{tang } \theta = \frac{\lambda'}{\alpha} \dots (126).$$

Finding the value of  $\theta$  by the last equation and, substituting it in equation (125), we obtain the value of  $k'$ ; and then, to find  $t$ , we have,

$$k' : L :: M g : t, \text{ or } t = \frac{L \times M g}{k'}.$$

$$M g = \sqrt{C g^2 - C M^2} = \sqrt{\downarrow^2 - \Delta^2} \text{ (Art. 451);}$$

$$\text{whence, } t = \frac{L \sqrt{\downarrow^2 - \Delta^2}}{k'} = \frac{L (\downarrow + \Delta)(\downarrow - \Delta)}{k'} \dots (127).$$

454. The apparent distance of the centres of the two bodies at the time of greatest obscuration being known, the quantity of the eclipse may be readily found. We have but to subtract the apparent distance from the sum of the apparent semi-diameters,

and state the proportion, as the sun's apparent diameter : the remainder : : 12 digits : the digits eclipsed. (For a more particular description of the method of calculating a solar eclipse, see Prob. XXX).

### *Occultations.*

455. At all places upon the earth's surface, which at a given time have the moon in the horizon, its apparent place will differ from its true place, by the amount of its horizontal parallax. It follows, therefore, that a star will be eclipsed by the moon somewhere upon the earth, in case its true distance from the moon's centre is less than the sum of the moon's semi-diameter and horizontal parallax.

The greatest value of the moon's semi-diameter is  $16' 45''$ , and that of its horizontal parallax  $61' 24''$ . If we add the sum of these numbers to  $5^{\circ} 17' 34''$ , the maximum latitude of the moon, we obtain as the result  $6^{\circ} 35' 43''$ . It is then only the stars which have a latitude less than  $6^{\circ} 35' 43''$ , that can experience an occultation from the moon.

456. By considering the various situations of the stars liable to an occultation, taking the greatest and least values of the sum of the moon's semi-diameter and horizontal parallax, and allowing for the inequalities of the motions of the moon, it has been found, that, if at the time of the mean conjunction of the moon and a star, (that is, when the moon's mean longitude is the same with the longitude of the star,) their difference of latitude exceed  $1^{\circ} 37'$ , there cannot be an occultation ; if the difference be less than  $51'$ , there must be an occultation somewhere on the earth ; and that between these limits there is a doubt, which can only be removed by the calculation of the moon's true place.

457. The calculation of an occultation is very nearly the same as that of a solar eclipse. The only difference is in the data. The star has no diameter, parallax, or motion in longitude ; and, as it is situated without the ecliptic, we have, in place of the latitude of the moon, employed in solar eclipses, the difference between the latitude of the moon and that of the star, and in place of the difference between the longitudes of the two bodies and their relative hourly motion in longitude, these quantities referred to an arc passing through the star and

parallel to the ecliptic. Thus, if  $E C$  (Fig. 63) represent the ecliptic,  $K$  its pole,  $s$  the situation of the star,  $M$  that of the moon, and  $s m'$  an arc passing through  $s$  and parallel to the arc  $E C$ , we have in place of  $m M$ ,  $m' M = m M - m m'$ , and in place of  $S m$ ,  $s m'$ . The hourly variation of  $S' m$  must also be reduced to the arc  $s m'$ .

458. The reduction of the difference of longitude of the moon and star, to the parallel to the ecliptic, passing through the star, is effected by multiplying this difference by the cosine of the latitude of the star. For, let  $A B$  (Fig. 65) be an arc of the ecliptic, and  $A' B'$  the corresponding arc of a circle parallel to it; then, since similar arcs of circles are proportional to their radii, we have,

$$B C : B' C' :: A B : A' B' = \frac{A B \cdot B' C'}{B C};$$

but,  $B' C' = C a = B' C \cos B C B' = B C \cos B B'$ .

Hence,  $A' B' = \frac{A B \cdot B C \cos B B'}{B C} = A B \cos B B'$ .

The reduction of the relative hourly motion in longitude to the parallel in question, is obviously effected in the same manner.

## CHAPTER XVII.

OF THE PLANETS AND THE PHENOMENA OCCASIONED BY  
THEIR MOTIONS IN SPACE.

### *Apparent Motions of the Planets with respect to the Sun.*

459. The apparent motion of an inferior planet, with reference to the sun, is materially different from that of a superior planet. The inferior planets always accompany the sun, being seen alternately on the east and west side of him, and never receding from him beyond a certain distance, while the superior



planets are seen at every variety of angular distance. This difference of apparent motion arises from the difference of situation of the orbits of an inferior and superior planet, with respect to the orbit of the earth, the one lying within and the other without the earth's orbit.

Let  $CAC'B$  (Fig. 66) represent the orbit of either one of the inferior planets, Venus for example, and  $PKT$  the orbit of the earth; which we will suppose to be circles and to lie in the same plane; and let  $MLN$  represent the sphere of the heavens to which all bodies are referred. Suppose, for the present, that the earth is stationary in the position  $P$ , and through  $P$  draw the lines  $PA$ ,  $PB$ , tangent to the orbit of Venus, and prolong them on till they intersect the heavens at  $a$  and  $b$ . When Venus is at  $C$ , (the earth being at  $P$ ), she will be in superior conjunction, and when at  $C'$  in inferior conjunction. Now, by inspecting the figure, it will be seen that in passing from  $C$  to  $C'$ , she will be seen in the heavens on the east side of the sun, and in passing from  $C'$  to  $C$  on the west side of the sun; also, that in passing from  $C$  to  $A$  she will recede from the sun in the heavens, from  $A$  to  $C'$  approach him, from  $C'$  to  $B$  recede from him again, and from  $B$  to  $C$  approach him again.  $a$  and  $b$  will be her positions in the heavens at the times of her greatest eastern and western elongations.

When Venus is to the east of the sun, she is seen in the evening, and called the *Evening Star*; and when to the west, she is seen in the morning, and called the *Morning Star*.

460. We have in the foregoing investigation supposed the earth to be stationary, a supposition which is contrary to the fact; but it is plain that the only effect of the earth's motion in the case under consideration, as it is slower than that of the planet, is to cause the points  $A$ ,  $C'$ ,  $B$  to advance in the orbit, without altering the nature of the apparent motion of the planet with respect to the sun. The orbits of the earth and planet are also ellipses of small eccentricity, and are slightly inclined to each other, instead of being circles and lying in the same plane: on this account, as the greatest elongations will occur in various parts of the orbits, they will differ in value. The greatest elongation of Venus varies from  $45^\circ$  to  $47^\circ 12'$ . Its mean value is about  $46^\circ$ .

461. Owing to the circumstance of the orbit of Mercury being

within the orbit of Venus, the greatest elongation of this planet is less than that of Venus. It varies between the limits  $16^{\circ} 12'$ , and  $28^{\circ} 48'$ ; and is, at a mean,  $22^{\circ} 30'$ .

462. Next, suppose  $P K T$  (Fig. 66) to be the orbit of a superior planet, and  $C A C' B$  that of the earth; and as the velocity of the earth is much greater than that of the planet, let us, for the present, regard the planet as stationary in the position  $P$ , while the earth describes the circle  $C A C'$ . When the earth is at  $C$ , the planet, being at  $P$ , is in conjunction with the sun. When the earth is at  $A$ ,  $S A P$  the elongation of the planet, is  $90^{\circ}$ . When it arrives at  $C'$  the planet is in opposition, or  $180^{\circ}$  distant from the sun. And when it reaches  $B$ , the elongation is again  $90^{\circ}$ . At intermediate points the elongation will have intermediate values. If, now, we restore to the planet its orbital motion, we shall manifestly be conducted to the same results relative to the change of elongation, as the only effect of such motion will be to throw the points  $A, C', B$  forward in the orbit. It appears, then, that in the course of a synodic revolution a superior planet will be seen at all angular distances from the sun, both on the east and west side of him. From conjunction to opposition, that is, while the earth is passing from  $C$  to  $C'$ , the planet will be to the right, or to the west of the sun; and will therefore be below the horizon at sunset, and rise some time in the course of the night. But, from opposition to conjunction, or while the earth is moving from  $C'$  to  $C$ , it will be to the east of the sun, and therefore above the horizon at sunset.

463. *To find the length of the synodic revolution of a planet.*

Let us first take an inferior planet, Venus for instance. Suppose we assume, at a given instant, the sun, Venus, and the earth to be in the same right line; then, after any elapsed time (a day for instance), Venus will have described an angle  $m$ , and the earth an angle  $M$  around the sun. Now,  $m$  is greater than  $M$ ; therefore at the end of a day, the separation of Venus from the earth, (measuring the separation by an angle formed by two lines drawn from Venus and the earth to the sun) will be  $m - M$ ; at the end of two days (the mean daily motions continuing the same), the angle of separation will be  $2(m - M)$ ; at the end of three days,  $3(m - M)$ ; at the end of  $s$  days,  $s(m - M)$ . When the angle of separation then amounts to  $360^{\circ}$ , that is, when

$s(m - M) = 360^\circ$ , the sun, Venus, and the earth must be again in the same right line, and in that case,

$$s = \frac{360^\circ}{m - M} \dots (128).$$

In which expression  $s$  denotes the mean duration of a synodic revolution,  $m$  and  $M$  being taken to denote the mean daily motions.

We may obtain from equation (128) another equation, in which the synodic revolution is expressed in terms of the sidereal periods of the earth and planet.

Let  $P$  and  $p$  denote the sidereal periods in question ; then since

$$1 \text{ d.} : M^\circ :: P : 360^\circ,$$

and

$$1 : m :: p : 360 ;$$

$M = \frac{360^\circ}{P}$  and  $m = \frac{360^\circ}{p}$  ; substituting

$$s = \frac{360^\circ}{360^\circ \left( \frac{1}{p} - \frac{1}{P} \right)} = \frac{P p}{P - p} \dots (129).$$

Equations (128), (129), although investigated for an inferior planet, will answer equally well for a superior planet, provided we regard  $m$  as standing for the mean daily motion of the earth,  $M$  for that of the planet,  $p$  for the sidereal period of the earth, and  $P$  for that of the planet. For, the earth holds towards a superior planet the place of an inferior planet, and a synodic revolution of the earth to an observer on the planet, will obviously be a synodic revolution of the planet to an observer on the earth.

464. Equation (128) shows that the length of a mean synodic revolution depends altogether upon the amount of the difference of the mean daily motions of the earth and planet, and is the greater the less is this difference.

It follows therefore that the synodic revolution is the longest for the planets nearest the earth.

It appears by equation (129), that the length of a synodic revolution is, for an inferior planet, greater than the sidereal period of the planet, and for a superior planet, greater than the sidereal period of the earth. The actual lengths of the synodic revolutions of the different planets are given in Table V.

465. The mean synodic revolution of a planet being known, and also the time of one conjunction or opposition, we may easily ascertain its mean elongation at any given time, and thus approximately the time of its rising, setting, and meridian passage.

*Stations and Retrogradations of the Planets.*

466. The apparent motions of the planets in the heavens, as has already been stated (Art. 8), are not, like those of the sun and moon, continually from west to east, or direct, but are sometimes also from east to west, or retrograde. The retrograde motion takes place over arcs of but a small number of degrees; and in changing the direction of their motions, the planets are for several days stationary in the heavens. These phenomena are called the *Stations* and *Retrogradations* of the planets. We now propose to inquire theoretically into the particulars of the motions in question, and to show how the phenomena just mentioned result from the motions of the planets in connection with the motion of the earth.

Let C A C' B (Fig. 66), represent the orbit of an inferior planet, and P K T the orbit of the earth; both considered as circles, and as situated in the same plane. If the earth were continually stationary in some point P of its orbit, it is plain that while the planet was moving from B the position of greatest western elongation, to A the position of greatest eastern elongation, it would advance in the heavens from *b* to *a*; that, while it was moving from A to B, that is, from greatest eastern to greatest western elongation, it would retrograde in the heavens from *a* to *b*; and that, in passing the points A and B, as it would be moving directly towards or from the earth, it would for a time appear stationary in the heavens in the positions *a* and *b*.

But the earth is in fact in motion, and the actual apparent motion of the planet is in consequence materially different from this. Let A, A' (Fig. 67), be the positions of the planet and earth at the time of the greatest eastern elongation, C', P their positions at inferior conjunction, and B, B' their positions at the greatest western elongation. At the time of the greatest eastern elongation while the planet describes a certain distance A D on the line of the centres of the earth and planet, the earth moves forward in its or-

bit a certain distance  $A'D'$ ; so that, instead of appearing stationary at  $a$  in the interval, the planet will advance in the heavens from  $a$  to  $d$ . From the same cause, it will have a direct motion about the time of the greatest western elongation. As it advances from  $A$  towards  $C'$ , the direct motion will continue; but, as the daily arc described by the planet will make a less and less angle with the daily arc described by the earth, the rate of motion will continually decrease, and finally when the planet has come into a position with respect to the earth, such that the lines of direction of the planet,  $mp, m'p'$ , at the beginning and end of the day are parallel, it will be stationary in the heavens. As the daily arc of the planet is greater than that of the earth, and becomes parallel to it in inferior conjunction, the planet will be in the position in question before it comes into inferior conjunction.

Subsequent to this, the inclination of the daily arcs still diminishing, the lines of direction of the planet at the beginning and end of the day will diverge, and therefore the motion will be retrograde. After inferior conjunction, the inclination of the arcs, will, at corresponding positions of the earth and planet, obviously, be the same as before. It follows therefore, that the planet will be at its western station, when it is at the same angular distance from the sun as at its eastern station; that its motion will be retrograde, until it has passed inferior conjunction and arrived at its western station; and that after this it will be direct.  $q$  and  $n$  represent the positions of the planet and the earth at the time of the western station;  $C'q = C'p$ , and  $Pn = Pm$ .

The diminution of the elongation of the planet at its two stations is not the only effect of the earth's motion in the case under consideration; it also accelerates the direct, and retards the retrograde motion of the planet, and gives to the planet along with the sun an apparent motion of revolution around the earth.

467. Let us now pass to the case of a superior planet. Suppose  $AC'B$  (Fig. 67), to be the orbit of the earth, and  $A'P'B'$  that of the planet. Since the earth is an inferior planet to an observer stationed upon a superior planet, it appears by the foregoing article, that it will, to an observer so situated, have a retrograde motion while it is passing over a certain arc  $pC'q$  in the inferior part of its orbit, and a direct motion during the remainder of the synodic revolution. Now, it is plain, that the direction of



the planet's motion, as seen from the earth, will always be the same as the direction of the earth's motion as seen from the planet. When the earth is at  $C'$  the middle of the arc  $p C' q$ , the planet is in opposition. It follows, therefore, that a superior planet has a retrograde motion during a small portion of its synodic revolution, about the time of opposition.

### *Phases of the Inferior Planets.*

468. To the naked sight the disc of the planet Venus appears circular, like that of each of the other planets, but the telescope shows this to be an optical illusion. When Venus is repeatedly observed with a telescope, it is seen to present in its various positions with respect to the sun the same variety of phases as the moon; being a full circle at superior conjunction, a half circle at the greatest eastern and western elongations, and a crescent with the horns turned from the sun, before and after inferior conjunction.

469. Mercury exhibits precisely similar phases, but being smaller, at a greater distance from the earth, and much nearer the sun, its phases are not so easily observed as those of Venus.

470. The phases of Venus are easily accounted for, by supposing it to be an opaque spherical body, and to shine by reflecting the sun's light, and by taking into consideration its motion with respect to the sun and earth. The hemisphere turned towards the sun is illuminated by him, and the other is in dark, and as the planet revolves around the sun, various portions of the enlightened half are turned towards the earth; in superior conjunction, the whole of it; at the greatest elongations, one half; and near inferior conjunction, but a small part. This will be abundantly evident on inspecting Fig. 68. The phases corresponding to the positions represented, are delineated in the figure.

The phases of Mercury are obviously susceptible of a similar explanation.

471. The disc of the planet Mars also undergoes changes of form, but they are of comparatively moderate extent. It is sometimes gibbous, but never has the form of a crescent. Indeed, on the supposition that Mars is an opaque body illuminated by the sun, we would not see the whole of the enlightened hemisphere, except

in conjunction and opposition, but there would always be more than half of it turned towards the earth, and therefore the disc should always be larger than a half circle.

472. The discs of the other superior planets do not experience any perceptible variation of form, for the reason, doubtless, that their orbits are so large with respect to the orbit of the earth, that all, or very nearly all of their illuminated hemispheres, is constantly visible from the earth.

### *Transits of the Inferior Planets.*

473. The two inferior planets Venus and Mercury, at inferior conjunction, sometimes, though rarely, pass between the sun and earth, and are seen as a dark spot crossing the sun's disc. This phenomenon is called a *Transit*. It will take place, in the case of either planet, whenever, at the time of inferior conjunction, it is so near either node that its geocentric latitude is less than the apparent semi-diameter of the sun.

474. The transits of Venus take place alternately at intervals of 8 and  $105\frac{1}{2}$  or  $121\frac{1}{2}$  years. The last were in the years 1761 and 1769. The next will be in 1874 and 1882.

475. In consequence of the greater distance of Mercury from the earth, a greater portion of its orbit is directly interposed between the sun and earth, than of the orbit of Venus; moreover, the synodic revolution of Mercury is shorter than that of Venus. On these accounts, it happens that the transits of Mercury are much more frequent than those of Venus. The last transit of Mercury was in the year 1835. The next two will take place in 1845 and 1848.

476. A transit is calculated in a precisely similar manner with a solar eclipse; the planet in the one calculation answering to the moon in the other.

477. A transit is an important phenomenon in a practical point of view, as it furnishes the most exact means we possess of ascertaining the sun's parallax. In order to understand how this phenomenon can be used for this purpose, we have only to consider that, in consequence of the difference of the parallaxes of the sun and Venus, observers at different stations upon the earth will refer the planet to different points upon the sun's disc, and that therefore, to such observers, the transit will take place along dif-

ferent chords, and be accomplished in unequal portions of time. It is then to be expected, that, if the durations of the transit at two different places should be noted, the *difference* of the parallaxes of the sun and Venus, upon which alone the difference of the duration depends, could be computed. This computation is in fact possible. Also, the *ratio* of the parallaxes being inversely as that of the distances, could be found by the elliptical theory, and thus the parallax both of the sun and Venus would become known.

478. The parallax of the sun, as it is now known, was deduced from observations upon the transits of Venus in 1769 and 1761. Expeditions were fitted out on the most efficient scale, by the British, French, Russian, and other governments, and sent to various parts of the earth, remote from each other, to observe the transit of 1769, that the parallax of the sun might be computed from the results of the observations. The sun's parallax as determined by Professor Encke from the observations made upon the transit in question, and that of 1761, is  $8''.5776$ .

*Appearances, Dimensions, Rotation, and Physical Constitution of the Planets.*

479. It appears from admeasurement with the telescope and micrometer, that the apparent diameter of a planet is subject to sensible variations. The apparent diameter of Venus, as well as of Mercury, is greatest in inferior conjunction, and least in superior conjunction; while the apparent diameter of each of the other planets is greatest in opposition and least in conjunction. These variations of the apparent diameters of the planets, are necessary consequences of the changes that take place in the distances of the planets from the earth.

480. The real diameter of a planet is deduced from its apparent diameter and horizontal parallax. (See Art. 395.) When the diameters of the planets have been found, their relative surfaces and volumes are easily obtained; for the surfaces are as the squares of the diameters, and the volumes as the cubes.

481. The order of magnitude of the planets is as follows: 1 Jupiter, 2 Saturn, 3 Uranus, 4 the Earth, 5 Venus, 6 Mars, 7 Mercury, 8 Pallas, 9 Ceres, 10 Juno, 11 Vesta. The range of magnitude, for the principal planets, is from 1 to 21000.

482. Spots more or less dark have been seen upon the discs of most of the principal planets; and by passing across them from east to west and re-appearing at the eastern limbs, have established that the planets upon which they are observed, rotate upon axes from west to east. From repeated careful observations upon the situations of these spots, the periods of rotation, and the positions of the axes, have been determined.

The periods of rotation of Mercury, Venus, the Earth, and Mars, are all about 24 hours, and of Jupiter and Saturn about 10 hours. Those of the other planets are not known. The axes of rotation remain continually parallel to themselves, as the planets revolve in their orbits.

483. The amount of light and heat, which the sun bestows upon the planets, decreases as we recede from the sun, in the same ratio that the square of the distance increases. (See Table IV.)

#### *Mercury.*

484. In consequence of its proximity to the sun, Mercury is rarely visible to the naked eye. When seen, it presents the appearance of a star of the 3d or 4th magnitude. Its phases show that it is opaque, and illuminated by the sun. Its apparent diameter varies with its distance from 5" to 12". Its real diameter is about 3140 miles, or  $\frac{2}{5}$  of that of the earth, and its volume is about  $\frac{1}{16}$  of the earth's volume.\*

Mercury performs a rotation on its axis in 24h. 5 $\frac{1}{2}$ m., and its axis is inclined to the ecliptic under a small angle. It is believed to be surrounded by a dense atmosphere.

#### *Venus.*

485. Venus is the most brilliant of all the planets, and generally appears larger and brighter than any of the fixed stars. At times, it emits so much light as to be visible in the open day, if the eye be protected from the sun. It is found by calculation, that the epochs in the course of a synodic revolution, at which Venus gives most light to the earth, are those at which, being in the inferior part of its orbit, it has an elongation of about 40°. They are about 36 days before and after inferior conjunction. The disc is then considerably less than a semi-circle, but the

---

\* The exact diameters, volumes, times of rotation, &c., of the different planets, as far as known, may be found in Table IV.



increased proximity to the earth more than compensates for the diminished size of the disc. Venus will besides attain to greater splendour in some revolutions than others, in consequence of being nearer the earth, when in the most favorable position.

486. As seen through a telescope, Venus presents a disc of nearly uniform brightness, and spots have very rarely been seen upon it. Its phases prove it to be an opaque spherical body, shining by reflecting the sun's light. Its apparent diameter varies with its distance from 10" to 61". Its real diameter is about 7700 miles, and its volume about  $\frac{1}{14}$  less than that of the earth. The period of its rotation is 23h. 21m. The inclination of its axis to the plane of its orbit is not exactly known, but is not far from 18°. Schroëter inferred, from a gradual diminution of its light observed at the edge of its disc, that it was surrounded by an atmosphere analogous to our own.

### *Mars.*

487. Mars is of the apparent size of a star of the first or second magnitude, and is distinguished from the other planets by its red and fiery appearance. The observed variation in the form of its disc (Art. 471), shows that it derives its light from the sun. Its greatest and least apparent diameters are respectively 4" and 18". Its real diameter is about 4100 miles, or rather more than  $\frac{1}{2}$  of the diameter of the earth, and its bulk is about  $\frac{1}{7}$  of that of the earth.

Mars revolves on its axis in 24h. 39m.; and its axis is inclined to the ecliptic in an angle of about 60°. It appears, from measurements made with the micrometer, that its polar diameter is less than the equatorial, and thus, that, like the earth, it is flattened at its poles. According to Sir W. Herschel, its oblateness (Art. 145) is  $\frac{1}{16}$ .

488. Spots of different shades are generally visible upon the surface of Mars, most of which always present the same appearance whenever they are distinctly seen.

The ruddy colour of the light of Mars has been generally attributed to a dense atmosphere surrounding the planet. Sir J. F. W. Herschel, however, has observed, in examining this planet with a good telescope, that some of its spots are of a reddish colour, and thence concludes that the ruddy colour of its light is owing to a red tinge in its soil.



*Jupiter and its Satellites.*

489. Jupiter is the most brilliant of the planets, except Venus, and sometimes even surpasses Venus in brightness. The eclipses of its satellites prove that it is an opaque body, and that it shines by reflecting the light of the sun. Its apparent diameter when greatest, is 46", and when least, 30".

Jupiter is the largest of all the planets. Its diameter is nearly 11 times the diameter of the earth, or about 86,000 miles, and its bulk is nearly 1300 times that of the earth. It turns on an axis nearly perpendicular to the ecliptic, and completes a rotation in 9h. 56m. The polar diameter is about  $\frac{1}{14}$  less than the equatorial.

490. When Jupiter is examined with a good telescope, its disc is always observed to be crossed by several obscure spaces, which are nearly parallel to each other, and to the plane of the equator. These are called the *Belts* of Jupiter. They are generally confined to the immediate vicinity of the equator, but they sometimes also extend to considerable distances from it, and have even been seen distributed over the whole face of the planet.

491. The satellites of Jupiter, as it has been already remarked, are visible with telescopes of moderate power. With the exception of the second, which is a little smaller, they are somewhat larger than the moon. The orbits of the satellites lie very nearly in the plane of Jupiter's equator.

492. Sir W. Herschel, in examining the satellites of Jupiter with a telescope, perceived that they underwent periodical variations of brightness. These variations he supposed to proceed from a rotation of the satellites upon axes, which caused them to turn different faces towards the earth; and from repeated and careful observations made upon them, he discovered that each satellite made one turn upon its axis in the same time that it accomplished a revolution around the primary; and, therefore, like the moon, presented continually the same face to the primary.

*Saturn, with its Satellites and Ring.*

493. Saturn shines with a pale dull light. Its apparent diameter varies only 3" or 4" by reason of the change of distance,

and is at the mean distance about  $16''$ . The eclipses of its satellites prove that it is opaque, and illuminated by the sun.

Saturn is the largest of the planets, next to Jupiter. Its diameter is nearly 10 times the diameter of the earth, or 79,000 miles; and its volume is nearly 1000 times that of the earth. The rotation on its axis is performed in 10h. 29m. The inclination of its axis to the ecliptic is about  $60^\circ$ . Its oblateness is  $\frac{1}{16}$ .

494. The disc of Saturn, like that of Jupiter, is frequently crossed with dark bands or belts, in a direction parallel to its equator. Extensive dusky spots are also occasionally seen upon its surface.

495. The planet Saturn is distinguished from all the other planets in being surrounded by a broad, thin, luminous ring, situated in the plane of its equator, and entirely detached from the body of the planet. This ring sometimes casts a shadow upon the planet, and is in turn, at times, partially obscured by the shadow of the planet; from which we conclude that it is opaque, and receives its light from the sun.

It is inclined to the plane of the ecliptic in an angle of about  $30^\circ$ , and during the motion of Saturn in its orbit, it remains continually parallel to itself. The face of the ring is, therefore, never viewed perpendicularly from the earth, and for this reason never appears circular, although such is its actual form. Its apparent form is that of an ellipse, more or less eccentric, according to the obliquity under which it is viewed, which varies with the position of Saturn in its orbit. When it is seen under the larger angles of obliquity, it appears as a luminous band, nearly encircling the planet, and is visible in telescopes of small power. Stars can also be seen between it and the planet in these positions. At other times, when viewed very obliquely, it can be seen only with telescopes of high power. When it is approaching the latter state, it has the appearance of two handles or *ansæ*, one on each side of the planet.

It is also at times invisible. This is the case whenever the earth and sun are on different sides of the plane of the ring, for the reason that the illuminated face is then turned from the earth. When the plane of the ring passes through the centre of the sun, the illuminated edge can be seen only in telescopes of extraordi-

nary power, and appears as a thread of light cutting the disc of the planet.

496. Since the orbit of Saturn is very large in comparison with the orbit of the earth, the plane of the ring will, during the greater part of the revolution of Saturn, pass without the orbit of the earth; and when this is the case, the ring will be visible, as the earth and sun will be on the same side of its plane. During the period, which is about a year, that the plane of the ring is passing by the orbit of the earth, the earth will sometimes be on the same side of it as the sun, and sometimes on opposite sides. In the former case the ring will be invisible, and in the latter will be seen so obliquely as to be visible only in telescopes of considerable or great power. All this will perhaps be better understood on consulting Fig. 69, where *ABC* represents the orbit of Saturn, *EFG* that of the earth, and *P, Q, N, R, S*, different positions of the ring.

The plane of the ring will pass through the sun every semi-revolution of Saturn, or, at a mean, about every 15 years, and at the epochs at which the longitude of the planet is respectively  $170^\circ$  and  $350^\circ$ , these being the longitudes of the nodes of the ring. The ring will then disappear once in about 15 years; but, owing to the different situations of the earth in its orbit, under circumstances oftentimes quite different. And the disappearance will occur when the longitude of the planet is about  $170^\circ$ , or  $350^\circ$ . The ring will be seen to the greatest advantage when the longitude of the planet is not far from  $80^\circ$  or  $260^\circ$ . The last disappearance took place in 1833; the next will be in 1848. At the present time (1838) the north face of the ring is visible.

497. From observations made upon bright spots seen on the face of the ring, Herschel discovered that it revolved from west to east about an axis perpendicular to its plane, and passing through the centre of the planet, (or very nearly). The period of its rotation is 10h. 29m. It is remarkable that this is the period in which a satellite assumed to be at a mean distance equal to the mean distance of the particles of the ring, would revolve around the primary according to the third law of Kepler.

The breadth of the ring is about one half greater than its distance from the surface of the planet, and is about equal to one third the diameter of the planet, or 29,000 miles.

498. What we have called Saturn's ring, consists in fact of two concentric rings, which turn together, although entirely detached from each other. The void space between them is perceived in telescopes of high power, under the form of a black circular line. According to the calculations of Sir John Herschel, from the micrometric measures of Prof. Struve, the breadth of the interior ring is about 17200 miles, and of the exterior about 10600 miles; the interval between the rings is about 1800 miles, and the distance from the planet to the inside of the interior ring is about 19000 miles. The thickness of the rings is not well known, the edge subtends an angle less than  $1''$ , which, at the distance of the planet, answers to about 4000 miles. Herschel makes it less than 100 miles.

499. The satellites of Saturn, with the exception of the 7th, revolve very nearly in the plane of the ring. The orbit of the 7th is inclined about  $30^\circ$  to this plane.

The 7th satellite is by far the largest and most conspicuous. The 1st and 2d, which just skirt the edge of the ring, have only been seen by Sir William Herschel in his large telescope.

The 7th satellite is subject to periodical variations of lustre, which prove its rotation on an axis in the period of a sidereal revolution of Saturn.

#### *Uranus and its Satellites.*

500. Uranus is scarcely visible to the naked eye. In a telescope it appears as a small round uniformly illuminated disc. Its apparent diameter is about  $4''$ , from which it never varies much, owing to the smallness of the earth's orbit in comparison with its own. Its real diameter is about 34000 miles, and its bulk 80 times that of the earth. Analogy leads us to believe that this planet is opaque and turns on an axis, but there is no direct proof that this is the case.

501. The satellites of Uranus were discovered by Sir W. Herschel. They are discernible only with telescopes of the highest power.

#### *Vesta—Juno—Ceres—Pallas.*

502. These four planets, although less distant than several of the others, are so extremely small, that they can only be seen with telescopes of considerable power.

503. The magnitudes of these planets are not well known.

Vesta is the smallest and also the most brilliant. Ceres and Pallas are said to have a nebulous or hazy appearance, indicative of an extensive vaporous atmosphere.

---

## CHAPTER XVIII.

### OF COMETS.—THEIR APPEARANCE, MAGNITUDE, AND PHYSICAL CONSTITUTION.

504. A comet usually consists of a mass of some luminous vapoury substance, called the *Coma*, condensed towards its centre around a brilliant *Nucleus* that is in general not very distinctly defined, from which diverges in a direction opposite to the sun, a stream of luminous transparent vapour, called the *Tail*. The coma and nucleus together form what is called the *Head* of the comet.

505. The length and form of the tail are very various. In some instances it is only a few degrees in length, and in others it is more than a quadrant. The tail of the great comet, which appeared in 1680, extended to a distance of  $70^{\circ}$  from the head, and that of the comet of 1618, to  $104^{\circ}$ .

506. When a comet first appears, no tail is perceptible, and its light is very faint. As it approaches the sun, it becomes brighter; the tail also, after a time, shoots out from the coma, and increases from day to day in extent and distinctness. As the comet recedes from the sun, the tail precedes the head, being still on the opposite side from the sun, and grows less and less, at the same time that, along with the head, it decreases in brightness, till at length the comet resumes nearly its first appearance.

The tail of a comet is the longest, and the whole comet is intrinsically the most luminous, soon after it has passed its perihelion. Its apparent size and lustre will not, however, generally be



the greatest at this time, as they will depend upon the distance and position of the earth, as well as the actual size and intrinsic brightness of the comet.

507. Individual comets offer considerable varieties of aspect. Some comets have been seen which were wholly destitute of a tail: such, among others, was the comet of 1682. Others have had more than one of these appendages. The comet of 1744 had six, which were spread over an extent of  $117^{\circ}$ ; and that of 1824 two, the one directed towards the sun, the other from him. Others still are without a nucleus, as the comet of 1795.

The comets that are visible only in telescopes, which are very numerous, have generally no distinct nucleus, and are unprovided with a tail. They have the appearance of round masses of luminous vapour, somewhat more dense towards the centre. Such are Encke's and Biela's comets.

508. Comets are the most voluminous bodies in the solar system. The tail of the great comet of 1680, was found by Newton to have been, when longest, no less than 123,000,000 miles in length. Some other comets have had tails of nearly as great length. The heads of comets are usually many thousand miles in diameter. That of the comet of 1811 had a diameter of 540000 miles.

509. The quantity of matter which enters into the constitution of a comet is exceedingly small. This is proved by the fact that the comets have had no influence upon the motions of the planets or satellites, although they have in many instances passed near these bodies. The comet of 1770, which was quite large and bright, passed through the midst of Jupiter's satellites without deranging their motions in the least perceptible degree. It also appears that the cometic matter is very rare and subtle, from the circumstance of stars of small magnitude being visible through all parts of the comet, with perhaps the single exception of the nucleus.

510. Of the physical constitution of the comets, little is known. It is not yet fully ascertained whether the vapoury substance of the coma and tail is self-luminous, or is illuminated by the sun. The nucleus is by some astronomers supposed to be, in some instances, a solid, and by others to be, in all cases, a highly condensed vapour. It has, in the case of a few comets, presented a

well-defined disc, like a solid, but it has, in no one instance, exhibited phases, which it is to be expected it would, if it were a solid shining by reflecting the light of the sun.

---

## CHAPTER XIX.

OF THE FIXED STARS.—THEIR NUMBER, AND DISTRIBUTION  
OVER THE HEAVENS—ANNUAL PARALLAX, AND DISTANCE  
—VARIABLE STARS—DOUBLE STARS—CLUSTERS OF STARS,  
AND NEBULÆ.

511. The number of stars that can be seen with the naked eye, does not much exceed 3000, and it is generally stated that not many more than 1000 are ever visible at any one time to the naked eye. But the telescope brings into view many millions, and every improvement made in it greatly increases the number.

512. As to the number of stars belonging to each different magnitude, astronomers assign from 15 to 20 to the first magnitude, from 30 to 60 to the second, about 200 to the third, and so on; the numbers increasing very rapidly as we descend in the scale of brightness; the whole number of stars already registered down to the seventh magnitude, inclusive, amounting to 15000 or 20000.

513. It is not to be understood that the classification of the stars into different magnitudes is made according to any fixed definite proportion subsisting between the degrees of apparent brightness of the stars belonging to different classes. Stars of almost every gradation of brightness, between the highest and the lowest, are met with. Those which offer marked differences of lustre, form the basis of the classification; others, which do not differ very widely from these, are united to them. As a necessary consequence, there are some stars of intermediate lustre, which cannot

be assigned with certainty to either magnitude. Thus, in the catalogue of the Astronomical Society of London, 3 stars are marked as intermediate between the first and second magnitudes, and 29 between the second and third.

514. As to the proportions of light emitted from the average stars of the different magnitudes, according to the experimental comparisons of Sir Wm. Herschel, they are, from the first to the sixth magnitude, in the ratio of the numbers, 100, 25, 12, 6, 2, 1.

515. With the exception of the three or four brightest classes, the stars are not distributed indiscriminately over the sphere of the heavens, but are accumulated in far greater numbers on the borders of the milky way, and in the milky way itself, which the telescope shows to consist of an immense number of stars of small magnitude in close proximity.

### *Annual Parallax and Distance of the Stars.*

516. The *Annual Parallax* of a fixed star is the angle made by two lines conceived to be drawn, the one from the sun and the other from the earth, and meeting at the star, at the time the earth is in such part of its orbit that its radius vector is perpendicular to the latter line; or, in other words, it is the greatest angle that can be subtended at the star by the radius of the earth's orbit. Thus, let S (Fig. 70) be the sun,  $s$  a fixed star, and E the earth, in such a position that the radius vector S E is perpendicular to the line of direction of the star, E  $s$ ; then the angle S  $s$  E is the annual parallax of the star  $s$ .

517. If the annual parallax of a star was known, we might easily find its distance from the earth; for, in the right angled triangle S E  $s$  we would know the angle S  $s$  E and the side S E, and we should only have to compute the side E  $s$ . Now, if any of the fixed stars have a sensible parallax, it could be detected by a comparison of the places of the star, as observed from two positions of the earth in its orbit, diametrically to each other; and accordingly, the attention of astronomers furnished with the most perfect instruments, has long been directed to such observations upon the places of some of the fixed stars which were supposed to be the nearest, in order to determine their annual parallax. But, after exhausting every refinement of observation, they have not been able to establish that any of them have a measura-

ble parallax. Now, such is the nicety to which the observations have been carried, that, did the angle in question amount to as much as  $1''$ , it could not possibly have escaped detection and universal recognition. We may then conclude, that *the annual parallax of the nearest fixed star is less than  $1''$* .

518. Taking the parallax at  $1''$ , the distance of the star comes out 206265 times the distance of the sun from the earth, or about 20 billions of miles. The distance of the nearest fixed star must therefore be greater than this. A juster notion of the immense distance of the fixed stars, than can be conveyed by figures, may be gained from the consideration that light, which traverses the distance between the sun and earth in 8m. 13s., and would perform the circuit of our globe in  $\frac{1}{8}$  of a second, employs more than three years in coming from the nearest fixed star to the earth.

519. The amount of light received from the same body at different distances, varies inversely as the square of the distance. Hence, if we admit the light of a star of each magnitude to be half that of one of the next higher magnitude, a star of the first magnitude would have to be removed to 181 times its distance, to appear no brighter than one of the sixteenth. Accordingly, if the difference in the apparent magnitude of the stars arises for the most part from a difference of distance, (which is the more probable supposition), there must be a multitude of stars visible in telescopes, the light of which has taken at least five hundred years to reach the earth.

### *Variable Stars.*

520. Several of the fixed stars are subject to periodical changes of brightness, and are hence called *Variable Stars*, or *Periodical Stars*. One of the most remarkable of the variable stars is the star *Omicron*, in the constellation Cetus. From being as bright as a star of the second magnitude, it gradually decreases, until it entirely disappears; and, after remaining for a time invisible, re-appears, and gradually increasing in lustre, finally recovers its original appearance. The period of these changes is 334 days. It remains at its greatest brightness about two weeks, employs about three months in waning to its disappearance, continues invisible for about five

months, and during the remaining three months of its period increases to its original lustre. Such is the general course of its phases. It does not, however, always recover the same degree of brightness, nor increase and diminish by the same gradations. And it is related by Hevelius, that in one instance it remained invisible for a period of four years, viz: from October, 1672, to December, 1676.

521. The greater number of variable stars undergo a regular increase and diminution of lustre, without ever, like the star just noticed, becoming entirely invisible. The star Algol, or  $\beta$  Perseii, is a remarkable variable star of this description. For a period of 2d. 14h. it appears as a star of the second magnitude, after which it suddenly begins to diminish in splendour, and in about  $3\frac{1}{2}$  hours is reduced to a star of the fourth magnitude. It then begins again to increase, and in  $3\frac{1}{2}$  hours more is restored to its usual brightness, going through all its changes in 2d. 20h. 48m.

522. There are also some instances on record of temporary stars having made their appearance in the heavens; breaking forth suddenly in great splendour, and without changing their positions among the other stars, after a time entirely disappearing. One of the most noted of these is the star which suddenly shone forth with great brilliancy on the 11th of November, 1572, in the constellation Cassiopeia, and was attentively observed by Tycho Brahé, a celebrated Danish astronomer. It was then as bright as any of the permanent stars, and continued to increase in splendour till it surpassed Jupiter when brightest, and was visible at mid-day. It began to diminish in December of the same year, and in March 1574 it entirely disappeared, after having remained visible for sixteen months, and has not since been seen.

In the years 945 and 1264, brilliant stars appeared in the same regions of the heavens. It is conjectured from the tolerably near agreement of the intervals of the appearance of these stars and that of 1572, that the three may be one and the same star, with a period of about 300 years. The places of the stars of 945 and 1264 are, however, too imperfectly known to establish this with any degree of certainty.

523. What is no less remarkable than the changes we have



noticed, several stars, which are mentioned by the ancient astronomers, have now ceased to be visible, and some are now visible to the naked eye which are not in the ancient catalogues.

### *Double Stars.*

524. Many of the stars which to the naked eye appear single, when examined with telescopes are found to consist of two (in some instances three) stars in close proximity to each other. These are called *Double Stars*. This class of bodies were first attentively observed by Sir William Herschel, who, in the years 1782 and 1785, published catalogues of a large number of them which he had observed. The list has since been greatly increased by Professor Struve, of Dorpat, Sir J. F. W. Herschel, and other observers.

525. Double stars are of various degrees of proximity. In a great number of instances, the angular distance of the individual stars is less than  $1''$ , and the two can only be separated by the most powerful telescopes. In other instances, the distance is from  $1'$  to  $2'$ , and the separation can be effected with telescopes of very moderate power. They are divided into six different classes, according to their distances, those in which the proximity is the closest forming the first class.

526. Sir William Herschel instituted a series of observations upon several of the double stars, with the view of ascertaining whether the apparent relative situation of the individual stars experienced any change, in consequence of the annual variation of the parallax of the star. With micrometers adapted to the purpose, he measured from time to time the apparent distance of the two stars, and the angle formed by their line of junction with the meridian at the time of the meridian passage, called the *Angle of Position*. Instead, however, of finding that annual variation of these angles, which the parallax of the earth's annual motion would produce, he observed that, in many instances, they were subject to regular progressive changes, which seemed to indicate a real motion of the stars with respect to each other. After continuing his observations for a period of twenty-five years, he satisfactorily ascertained that the changes in question were in reality produced by a motion of revolution of one star around the other, or of both

around their common centre of gravity ; and in two papers, published in the Philosophical Transactions for the years 1803 and 1804, he announced the important discovery that there exist sidereal systems, composed of two stars revolving about each other in regular orbits. These stars have received the appellation of *Binary Stars*, to distinguish them from other double stars which are not thus physically connected, and whose apparent proximity may be occasioned by the circumstance of their being situated on nearly the same line of direction from the earth, though at very different distances from it.

527. Since the time of Sir W. Herschel, the observations upon the binary stars have been continued by several distinguished astronomers. From the observations made upon some of them, astronomers have been enabled to deduce the form of their orbits, and approximately the lengths of their periods. The orbits are ellipses of considerable eccentricity. The periods are of various lengths, as will be seen from the following enumeration of those which are considered as the best ascertained :  $\gamma$  *Leonis* 1200 years ;  $\gamma$  *Virginis* 629 years ; 61 *Cygni* 452 years ;  $\sigma$  *Coronæ* 287 years ; Castor 253 years ; 70 *Opiuchi* 80 years ;  $\xi$  *Ursæ* 58 years ;  $\zeta$  *Canceri* 55 years ; and  $\eta$  *Coronæ* 43 years.

### *Clusters of Stars—Nebulæ.*

528. Many spaces are discovered in the heavens which are faintly luminous, and shine with a pale white light. These are called *Nebulæ*. Some are visible to the naked eye, but the greater number cannot be seen without the aid of a good telescope. On applying to them telescopes of great power, they are found for the most part to consist of a multitude of small stars, distinctly separate, but very near each other, and more or less condensed towards the centre.

529. There are also clusters of stars in close proximity, dispersed here and there over the sphere of the heavens, which are seen to be such with the naked eye, or with telescopes of only moderate power. One of the most conspicuous of these clusters, is that called the *Pleiades*.

To the unaided sight it appears to consist of six or seven stars, but a telescope even of moderate power exhibits within the space they occupy fifty or sixty conspicuous stars. The constellation

called *Coma Berenices*, is another group, more diffused, and composed of larger stars.

In the constellation *Cancer* there is a luminous spot, or nebula, called *Præsepe*, or the bee-hive, which a telescope of moderate power resolves entirely into stars. In *Perseus* is another spot crowded with stars, which become separately visible with a good telescope.

530. A considerable number of nebulae are met with in different parts of the heavens, which offer no appearance of stars, even when examined with telescopes of the highest power. A very great diversity of form and aspect obtains among them. One of the most prominent is that near the star  $\nu$  in *Andromeda*. It is visible to the naked eye, and has often been mistaken for a comet.

## PART III.

### OF THE THEORY OF UNIVERSAL GRAVITATION.

---

#### CHAPTER XX.

##### OF THE PRINCIPLE OF UNIVERSAL GRAVITATION.

531. It is demonstrated in treatises on Mechanics, that if a body move in a curve in such a manner that the areas traced by the radius vector about a fixed point, increase proportionally to the times, it is solicited by an incessant force constantly directed towards this point. Now, by Kepler's first law, the areas described by the radii vectores of the planets about the sun, are proportional to the times. It follows therefore from this law, that each planet is acted upon by a force which urges it continually towards the sun.

This fact is technically expressed, by saying that the planets *gravitate* towards the sun, and the force which urges each planet towards the sun is called its *Gravity*, or Force of Gravity towards the sun.

532. It is also proved by the principles of Mechanics, that if a body, continually urged by a force directed to some point, describe an ellipse of which that point is a focus, the force by which it is urged must vary inversely as the square of the distance. It therefore follows from Kepler's second law, viz : that the planets describe ellipses having the centre of the sun at one of their foci ; that the force of gravity of each planet towards the sun, varies inversely as the square of the distance from the sun's centre.

533. By taking into view Kepler's third law, it is proved that it is one and the same force, modified only by distance from the

sun, which causes all the planets to gravitate towards him, and retains them in their orbits. This force is conceived to be an attraction of the matter of the sun for the matter of the planets, and is called the *Solar Attraction*.

534. The motions of the satellites are in conformity with Kepler's laws; hence, the planets which have satellites are endowed with an attractive force of the same nature with that of the sun.

535. The existence of a similar attractive power in each of the planets that are devoid of satellites, is proved by the fact that the observed inequalities of their motions, and of those of the other planets, may be shown upon this supposition to be necessary consequences of the attractions of the planets for each other.

536. In like manner the inequalities in the motions of the satellites and their primaries, show that the satellites possess the same property of attraction as the sun.

537. We learn from the motions produced by the action of the sun and planets upon each other, that the intensities of their attractive forces are, at the same distance, proportional to their masses, and that the whole attraction of the same body for different bodies, is, at the same distance, proportional to the masses of these bodies. From which we may infer that a mutual attraction exists between the particles of bodies, and that the whole force of attraction of one body for another, is the result of the attractions of its individual particles. Moreover, analysis shows, that in order that the law of attraction of the whole body may be that of the inverse ratio of the square of the distance, this must also be the law of attraction of the particles. The fact, as well as the law of the mutual attraction of particles, is also revealed by the tides, and other phenomena referable to such attraction.

538. The celestial phenomena compared with the general laws of motion, conduct us therefore to this great principle of nature; namely, *that all particles of matter mutually attract each other in the direct ratio of their masses, and in the inverse ratio of the squares of their distances*. This is called the principle of *Universal Gravitation*. The theory of its existence was first promulgated by Sir Isaac Newton, and is hence often called *Newton's Theory of Universal Gravitation*. The force which urges the particles of matter towards each other, is called the *Force of Gravitation*, or the *Attraction of Gravitation*.



539. In the following chapters our object will be to develop the most important effects of the principle of gravitation thus arrived at by induction. The perfect accordance that will be observed to obtain between the deductions from the theory of universal gravitation and the results of observation, will afford additional confirmation of the truth of the theory.

---

## CHAPTER XXI.

### THEORY OF THE ELLIPTIC MOTION OF THE PLANETS.

540. Let the attraction of the unit of mass of the sun for the unit mass of a planet, at the unit of distance, be designated by 1. The whole attraction exerted by the sun upon the unit of mass, at the same distance, will then be expressed by the mass of the sun ( $M$ ); or, in other words, by the number of units which its mass contains. And the attraction  $F$ , at any distance  $r$ , will result from the proportion  $M : F :: r^2 : 1^2$ , which gives  $F = \frac{M}{r^2}$ . This, in the language of Dynamics, is the *Accelerating Force* soliciting the planet.

As  $\frac{M}{r^2}$  expresses the attraction of the sun for a unit of mass of the planet, its attraction for the entire mass  $m$  of the planet will be expressed by  $m \frac{M}{r^2}$ . This is the *moving force* of the planet, and since it is, at the same distance, proportional to the mass of the planet, the velocity due to its action is the same, whatever may be the mass.

541. The planet has also an attraction for the sun, as well as the sun for the planet, and the expression for its attractive force, or for the accelerating force animating the sun, will obviously

be  $\frac{m}{r^2}$ . The sun will then tend towards the planet, as the planet towards the sun. But, if the two bodies were to set out from a state of rest, the velocity of the planet would be as many times greater than the velocity of the sun, as the mass of the sun is greater than that of the planet. For, the velocity of the planet would be to that of the sun as the attractive force of the sun is to the attractive force of the planet, that is, as  $\frac{M}{r^2} : \frac{m}{r^2}$ , or as  $M : m$ .

As the attraction of the particles of the sun and planet are mutual and equal, the attraction of the planet for the entire mass of the sun must be equal to the attraction of the sun for the entire mass of the planet.

542. *The sun and any planet revolve about their common centre of gravity.*

To show this, we would remark, in the first place, that it is a principle of Mechanics that the mutual actions of the different members of a system of bodies cannot affect the state of the centre of gravity of the system. This is called the *Principle of the Preservation of the Centre of Gravity*. It follows from it that the common centre of gravity of the sun and any planet is at rest, unless it has a motion of translation in common with the two bodies, imparted by a force extraneous to the system. As we are concerned at present only with the relative motion of the sun and planet, such motion of translation, if it does exist, may be left out of account. Now, let S (Fig. 71) be the sun, and P any planet, supposed for the moment to be at rest. If neither of the two bodies should receive a velocity in a direction oblique to PS the line of their centres, they would move towards each other by virtue of their mutual attraction, and meet at C their common centre of gravity.\* But, if the body P have a projectile velocity given to it in any direction Pt, inclined to the line PS, it is susceptible of proof that its motion relative to the sun may be in an ellipse, as is observed to be the case with the planets.

Now, while the planet moves in space, the line of the centres

---

\* The common centre of gravity of two bodies lies on the line joining their centres, and divides this line into parts inversely proportional to the masses of the bodies.

of the planet and sun must continually pass through the stationary position of the centre of gravity, and therefore, when the planet has advanced to any point  $p$ , the sun will have shifted its position to some point  $s$  on the line  $pC$  prolonged. Moreover, as the two bodies mutually gravitate towards each other, the paths of each in space will be continually ~~conclude~~ towards the other body, and, therefore, also towards the centre of gravity  $C$ , which is constantly in the same direction as the other body. Since the planet performs a revolution around the sun, the sun and planet must each continue to move about the point  $C$  until they have accomplished a revolution and returned to the line  $PCS$ . Also as the distance  $PS$  of the two bodies will be the same at the end as at the beginning of the revolution, as well as the ratio of their distances  $PC$  and  $SC$  from the centre of gravity, they will return to the positions  $PS$ , from which they set out, and will therefore move in continuous curves.

As the distances of the sun and planet from their common centre of gravity are constantly reciprocally proportional to their masses, the orbit of the sun will be exceedingly small in comparison with the orbit of the planet.

543. If to both the sun and planet there should be applied a force equal to the accelerating force of the sun,  $\frac{m}{r^2}$  (Art. 541), but in an opposite direction, the sun would be solicited by two forces that would destroy each other, but the planet would now be urged towards the sun remaining stationary, with the accelerating force  $\frac{M+m}{r^2}$ , or a force the intensity of which was equal to the sum of the intensities of the attractive forces of the sun and planet, at the distance of the planet. Now, the application of a common force will not alter the relative motion of the two bodies. Hence, in investigating this motion we are at liberty to conceive the sun to be stationary, if we suppose the planet to be solicited by the accelerating force  $\frac{M+m}{r^2}$ . As the mass of the sun is very much greater than that of any planet, but little error will be committed in neglecting the attraction of the planet, and taking into account only the sun's action  $\frac{M}{r^2}$ .

544. Analysis makes known the general laws of the motion of a body, when impelled by a projectile force, and afterwards continually attracted towards the sun's centre by a force varying inversely as the square of the distance. We learn by it that the body will necessarily describe some one of the conic sections, around the sun situated at one of its foci. We learn, also, that the nature of the orbit, as well as the length of the major axis, is wholly dependent, for any given distance of the planet, upon the *intensity* of the projectile force, but that the position of the axis, and the eccentricity of the orbit, depend also upon the *angle* of projection, (that is, the angle included, at the commencement of the motion, between the line of direction of the projectile force and the radius vector.) As to the relative intensity of the projectile force necessary to the production of each one of the conic sections, a certain intensity of force will produce a parabola; any less intensity, an ellipse or circle; and any greater, a hyperbola.

545. If the velocity that would at a given distance be imparted by the sun's attraction in a second of time, which is the measure of its intensity at the given distance, be found, and also the distance of a planet at any time, as well as its velocity and the angle made by the direction of its motion with the radius vector, the form, dimensions, and position of the planet's orbit can be computed. This is to determine the orbit *à priori*. The practice has been, however, to determine the various elements of a planet's orbit by observation, (as already described, Chap. VIII).

The elements being known, the equations of the elliptic motion, investigated on the principles of Mechanics, serve to make known the position and velocity of the planet at any time. (The investigation of these equations may be found in the *Encyclopædia Metropolitana*, Article *Physical Astronomy*, page 653, in the *Mécanique Elementaire* de Francœur, and in many other similar works.)\*

546. The physical theory of the motion of a satellite around

---

\* The equations are the same with those deduced directly from Kepler's laws of the planetary motions. (See App., Solution of Kepler's Problem.)

its primary is obviously the same as that of the motion of a planet around the sun.

547. According to the principle of the preservation of the centre of gravity (Art. 542), the centre of gravity of the whole solar system must either be at rest, or have a motion of translation in space in common with the system, resulting from the action of a foreign force. If any general motion of the solar system subsists, it has not yet been recognized from observation.

548. The sun and planets revolve around their common centre of gravity. The path of the sun's centre results from the joint action of all the planets, and is a complicated curve. As the quantity of matter in all the planets taken together is very small, compared with that in the sun, the extent of the curve described by the centre of the sun cannot be very great. It is found by computation, that the distance between the sun's centre and the centre of gravity of the system can never be equal to the sun's diameter.

549. It is demonstrated in treatises on Mechanics, that if foreign forces act upon a system of bodies, the centre of gravity of the system will move just as the whole mass of the system concentrated at the centre of gravity would move, under the action of the same forces. It follows from this principle, that from the attraction of the sun for a primary planet and its satellites, their common centre of gravity will revolve around the sun, just as the whole quantity of matter in the planet and its satellites concentrated at this point would, under the influence of the same attraction. Moreover, the same considerations which show that the sun and planets revolve about their common centre of gravity, will also show that a primary planet and its satellites revolve about their common centre of gravity. It appears, therefore, that in the case of a planet which has satellites, it is not, strictly speaking, the centre of the planet that moves agreeably to the first and second laws of Kepler, but the common centre of gravity of the planet and its satellites; the planet and satellites revolving around the centre of gravity, as it describes its orbit about the sun.

550. It may be worth while here to remark, that the revolution of the earth around the common centre of gravity of the



earth and moon, occasions an inequality, both of longitude and latitude, in the apparent motion of the sun. It is, however, exceedingly small, for the reason that the distance of the earth's centre from the centre of gravity is very short, in comparison with the distance of the sun. The mass of the earth is to that of the moon as 80 to 1, while the distance of the moon is to the radius of the earth as 60 to 1. It follows, therefore, that the common centre of gravity of the earth and moon lies within the body of the earth.

551. It appears also from the physical investigation of the elliptic motion of the planets, that Kepler's third law is not rigorously true. In consequence of the action of the planets upon the sun, the ratio of the periodic times of the different planets depends upon the masses of the planets, as well as their distances from the sun. If  $p$  and  $p'$  be the periodic times of any two of the planets,  $a$  and  $a'$  their mean distances from the sun's centre, and  $m$  and  $m'$  their quantities of matter, that of the sun being denoted by 1, then, disregarding the actions of the other planets,

$$p^2 : p'^2 :: \frac{a^3}{1+m} : \frac{a'^3}{1+m'}.$$

As  $m$  and  $m'$  are very small fractions, the error resulting from their omission will be very small. If we omit them, we shall have,

$$p^2 : p'^2 :: a^3 : a'^3 ;$$

which is Kepler's third law.

## CHAPTER XXII.

### THEORY OF THE PERTURBATIONS OF THE ELLIPTIC MOTION OF THE PLANETS AND OF THE MOON.

552. We have, in a previous chapter, given a general idea of the mode of determining from theory and observation combined, the law and amount of the perturbations or inequalities of the

lunar and planetary motions. We propose now to give some insight into the nature and manner of operation of the disturbing forces, and will commence with the perturbations of the moon produced by the action of the sun.

553. We have already (Art. 259) showed how the intensity and direction of the disturbing force of the sun, in any given position of the moon in its orbit, may be determined. Let us now derive the disturbing forces that take effect in the three directions in which the motion of the moon can be changed; namely, in the direction of the radius vector, of the tangent to the orbit, and of the perpendicular to its plane. Let E (Fig. 72) be the earth, M the moon, and S the sun. Let the force exerted by the sun upon the moon be decomposed into two forces, one acting along the line  $MS'$  parallel to  $ES$ , and the other from M towards E. If the component along  $MS'$  were equal to the force exerted by the sun upon the earth, the motion of the moon about the earth would not be changed by the action of these two forces. Hence, the difference between them will be the disturbing force in the direction  $MS'$ . The component along  $ME$  is another disturbing force. It is called the *Additious Force*, because it tends to increase the gravity of the moon towards the earth. The disturbing force along  $MS'$  will generally be inclined to the plane of the orbit, and may be decomposed into three forces, one in the direction of the tangent, another in the direction of the radius vector, and a third in the direction of the perpendicular to the plane. The first mentioned component is called the *Tangential Force*; the second is called the *Ablatitious Force*; and the third we shall call the *Perpendicular Force*.

The actual disturbing force in the direction of the radius vector is equal to the difference between the additious and ablatitious forces, and is called the *Radial Force*. This and the tangential and perpendicular forces constitute the disturbing forces, the direct operation of which is to be considered.

554. To obtain general analytical expressions for these forces, let the distance of the sun from the earth (which for the present we shall suppose to be constant) be denoted by  $a$ , and the distances of the moon from the earth and sun, respectively, by  $y$ , and  $z$ . Also let  $F$  = the force exerted by the earth upon the moon,  $P$  = the force exerted by the sun upon the earth, and  $Q$  = the force

exerted by the sun upon the moon. Then, if we denote the mass of the earth by 1, and take  $m$  to stand for the mass of the sun, we shall have, (Art. 539),

$$F = \frac{1}{y^2}, P = \frac{m}{a^2}, Q = \frac{m}{z^2}.$$

Let the force  $Q$  be represented by the line  $MS$  (Fig. 72); and let its component parallel to  $ES$ , or  $MS' = R$ , and its component along the radius vector, or  $ME = T$ .

$$Q : T :: MS : ME; \text{ or, } \frac{m}{z^2} : T :: z : y.$$

Whence, addititious force  $T = \frac{m y}{z^3} \dots (130).$

In a similar manner we obtain,

$$R = \frac{m a}{z^3} \dots (131).$$

The disturbing force in the direction of the sun

$$= R - P = \frac{m a}{z^3} - \frac{m}{a^2} = m a \left( \frac{1}{z^3} - \frac{1}{a^3} \right).$$

Now, let  $\alpha, \beta, \gamma$ , denote the angles made by the line  $MS'$ , respectively, with the tangent, the radius vector, and the perpendicular to the plane of the orbit, and we shall have for the components of the disturbing force  $R - P$ , along these lines;

$$\text{tangential force} = m a \left( \frac{1}{z^3} - \frac{1}{a^3} \right) \cos \alpha \dots (132);$$

$$\text{ablative force} = m a \left( \frac{1}{z^3} - \frac{1}{a^3} \right) \cos \beta \dots (133);$$

$$\text{perpendicular force} = m a \left( \frac{1}{z^3} - \frac{1}{a^3} \right) \cos \gamma \dots (134).$$

Combining equation (133) with equation (130) we obtain for the radial force,

$$\text{radial force} = m y \frac{1}{z^3} - m a \left( \frac{1}{z^3} - \frac{1}{a^3} \right) \cos \gamma.$$

555. The obliquity of the orbit of the moon to the plane of the ecliptic, affects but very slightly the value of the tangential and radial forces. If we leave it out of account, or suppose the moon's orbit to lie in the plane of the ecliptic, we shall have

(Fig. 73)  $\beta = S' M L = S E M$  the elongation of the moon  $= \varphi$ , and  $\alpha =$  complement of  $\varphi$ , which gives,

$$\text{tangential force} = m a \left( \frac{1}{z^3} - \frac{1}{a^3} \right) \sin \varphi \dots (135).$$

$$\text{radial force} = m y \frac{1}{z^3} - m a \left( \frac{1}{z^3} - \frac{1}{a^3} \right) \cos \varphi \dots (136).$$

556. Equation (134) may be transformed into another, which is better adapted to the purposes we have in view. Let  $M K$  (Fig. 72) represent the perpendicular to the plane of the moon's orbit,  $M F$  the intersection of the plane  $S M K$  with the plane of the moon's orbit, and  $S I$ ,  $I F$  the intersections of a plane passing through  $S$  and perpendicular to  $E N$ , the line of nodes, with the plane of the ecliptic and the plane of the orbit.  $S F$  will be perpendicular to both  $I F$  and  $M F$ . Denote  $S I F$  the inclination of the orbit to the ecliptic, by  $I$ ,  $S E N$  the angular distance of the sun from the node by  $N$ , and  $S E$  and  $S M$  by  $a$ , and  $z$ , as before.

Now, in equation (134)  $\gamma$  stands for the angle  $S' M K$ , but  $S' M K = S M K$  (nearly), and,

$$\cos S M K = \sin S M F = \frac{S F}{S M};$$

$$S F = S I \sin S I F, \text{ and } S I = S E \sin S E I;$$

$$\text{whence, } S F = S E \sin S E I \sin S I F = a \sin N \sin I;$$

substituting,

$$\cos \gamma = \cos S M K = \frac{a \sin N \sin I}{S M} = \frac{a \sin N \sin I}{z}.$$

Thus we have,

$$\text{perpen. force} = m a \left( \frac{1}{z^3} - \frac{1}{a^3} \right) \frac{a \sin N \sin I}{z} \dots (137).$$

557. The variable  $z$  may be eliminated from equations (135), (136) and (137), and other equations obtained, involving only the variables  $y$  and  $\varphi$ . Let  $M L$  (Fig. 72) be drawn through the place of the moon perpendicular to  $E S$ . Then, using the same notation as in the preceding articles,

$$L S = z \text{ (nearly), } E L = E M \cos L E M = y \cos \varphi;$$

$$\text{but, } L S = S E - E L;$$

$$\text{whence, } z = a - y \cos \varphi, \text{ and } z^3 = a^3 - 3 a^2 y \cos \varphi:$$

neglecting the terms containing the higher powers of  $y$  than the first, as they are very minute,  $y$  being only about  $\frac{1}{4} \frac{1}{60} a$ .

$$\frac{1}{z^3} = \frac{1}{a^3 - 3 a^2 y \cos \varphi} = \frac{1}{a^3} + \frac{3 y \cos \varphi}{a^4};$$

neglecting all the terms of the quotient that involve higher powers of  $y$  than the first. Substituting this value of  $\frac{1}{z^3}$  in equation (135), we obtain,

$$\text{tangential force} = \frac{3 m y \cos \varphi \sin \varphi}{a^3};$$

or, (App. For. 13),

$$\text{tangential force} = \frac{3 m y \sin 2 \varphi}{2 a^3} \dots (138).$$

Making the same substitution in equation (136), and neglecting the term containing  $y^2$ , there results,

$$\text{radial force} = \frac{m y (1 - 3 \cos^2 \varphi)}{a^3};$$

or, (App. For. 9),

$$\text{radial force} = - \frac{m y (1 + 3 \cos 2 \varphi)}{2 a^3} \dots (139).$$

In equation (137) we have to substitute, besides, the value of  $z$ , viz:  $a - y \cos \varphi$ ; then dividing and neglecting as before, we have,

$$\text{perpen. force} = \frac{3 m y \cos \varphi}{a^3} \sin N \sin I \dots (140.)$$

558. If the disturbing forces retained constantly the same intensity and direction, the result would be a continual progressive departure from the elliptic place; but, in point of fact, these forces are subject to periodical changes of intensity and direction from several causes, from which results a compensation of effects, and an eventual return to the elliptic place. The causes of the variation of the disturbing forces are:

1. The revolution of the moon around the earth.
2. The elliptic form of the apparent orbit of the sun.
3. The elliptic form of the orbit of the moon.
4. The inclination of the two orbits.



As the variations of the radial and tangential forces, resulting from the inclination of the orbits, are very minute, we shall leave them out of account, and in the consideration of the effects of these forces, shall, for the sake of simplicity, regard the orbits as lying in the same plane.

The first mentioned circumstance is the most prominent cause of variation, and gives rise to the more conspicuous perturbations. The other two serve to modify the variations of the forces resulting from the first, and occasion each a distinct set of periodical perturbations.

559. Let us now investigate, in succession, the effects of each of the disturbing forces, commencing with the tangential force. The tangential force takes effect directly upon the velocity of the moon in its orbit; and as its line of direction does not pass through the earth, it disturbs the equable description of areas. It also affects the radius vector indirectly, by changing the centrifugal force. To understand the detail of its action, we must inquire into the variations which it undergoes.

If we regard  $y$  as constant in the expression for the tangential force, (equa. 138), which amounts to considering the moon's orbit as circular, the expression will become equal to zero when  $\sin 2\varphi = 0$ , and will have its maximum value when  $\sin 2\varphi = 1$ . It will also change its sign with  $\sin 2\varphi$ . It appears, therefore, that the tangential force is zero in the syzgies and quadratures, where it also changes its direction. and that it attains its maximum value in the octants. It will be seen, on inspecting Fig. 74, that it will be a retarding force in the first quadrant, (A B). Accordingly, it will be an accelerating force in the second, a retarding force again in the third, and an accelerating force again in the fourth.

This will also appear upon considering the direction of the disturbing force parallel to the line of the centres of the sun and earth, in the various quadrants. In the nearer half of the orbit the sun tends to draw the moon away from the earth, and the force in question is directed towards the sun. In the more remote half the sun tends to draw the earth away from the moon, but we may regard it, instead, as urging the moon from the earth by the same force; for, the relative motion will be the same on this supposition. In the part of the orbit supposed,

then, the disturbing force under consideration will be directed from the sun, as represented in Fig. 74.

560. It appears, then, that the tangential force will alternately retard and accelerate the motion of the moon during its passage through the different quadrants, and that the maximum of velocity will occur in the syzigies A, C, where the accelerating force becomes zero, and the minimum of velocity in the quadratures B, D, where the retarding force becomes zero. On the supposition that the orbit is a circle, the arcs A B, B C, C D, and D A, would be equal, and the retardation of the velocity in one quadrant would be compensated for by an equal acceleration in the next, and at the close of a synodic revolution, the velocity of the moon would be the same as at its commencement. As the velocity is greatest in the syzigies, and least in the quadratures, and as the degree of retardation is the same as that of acceleration, the mean motion\* must have place in the octants. Now, as the moon moves from the syzigy A with a motion greater than the mean motion, her true place will be in advance of her mean place, and will become more and more so till she reaches the octant, where the true motion is equal to the mean. The difference between the true and mean place will then be the greatest; for after that, the true motion becoming less than the mean, the mean place will approach nearer to the true, till at the quadrature they coincide. Beyond B, the true motion still continuing less than the mean, the mean place will be in the advance of the true, and the separation will increase till at the octant the true motion has attained to an equality with the mean motion, after which, the mean motion being the slowest, the true place will approach the mean till at the syzigy C they again coincide. Corresponding effects will take place in the two remaining quadrants. We perceive, therefore, that the tangential force produces an inequality of longitude, which attains to its maximum positive and negative value in the octants, and is zero in the syzigies. This is the inequality known in Plane Astronomy by the name of *Variation*, (Art. 272).

561. Let us now inquire into the modifications of the effects of the tangential force, that result from the elliptic form of the sun's

---

\* The expressions, mean motion, true motion, mean place, true place, are here to be understood only in relation to the perturbation under consideration.

orbit. Suppose that at the moment when the moon sets out from conjunction, the sun is in the apogee of its orbit: then it is plain, that, during the whole revolution of the moon, the sun's disturbing force would be on the increase by reason of the diminution of the sun's distance, and *that*, in consequence, the retardation in the first quadrant would be less than the acceleration in the second, and the retardation in the third less than the acceleration in the fourth. So, that, when the moon had again come around into conjunction, the acceleration would have over-compensated the retardation. This kind of action would go on so long as the sun approached the earth; but when it had passed the perigee of its orbit, and began to recede from the earth, the reverse effect would take place, and a retardation of the moon's orbital motion would happen each revolution. If the anomalistic revolution of the sun was an exact multiple of the synodic revolution of the moon, the acceleration in each revolution of the moon during the passage of the sun from the apogee to the perigee of its orbit, would be compensated for by an equivalent retardation in the revolution of the moon answering to the same distance of the sun in its passage from the ~~apogee to the perigee~~ and the velocity of the moon would be the same at the close of an anomalistic revolution of the sun as at its commencement. But as this relation does not, in fact, subsist between the anomalistic revolution of the sun and the synodic revolution of the moon, a compensation between the accelerations and retardations answering to the different revolutions of the moon, will not be effected until conjunctions shall have occurred at every variety of distance of the sun in each half of its orbit. Since the anomalistic and synodic revolutions are incommensurable, the sun will be, in reality, in every variety of position in its orbit, at the time of conjunction, in process of time; so that eventually the original velocity in conjunction will be regained. It appears, therefore, that the variation of the moon's motion from one revolution to another, occasioned by the elliptic form of the sun's orbit, is periodic. Its period will be the interval of time in which the moon will perform a certain number of synodic revolutions, while the sun performs a certain number of anomalistic revolutions. Avoiding unnecessary precision, we find it to consist of but a moderate number of years.

*+ perigee to the Apogee*

562. We have next to consider the consequences of the elliptic form of the moon's orbit. We remark in the first place that the orbit being an ellipse, the areas A E B, B E C, C E D, and D E A (Fig. 74) will be unequal, and therefore, by the laws of elliptic motion, the arcs A B, B C, C D, and D A will be described in unequal times. It follows from this, that the retardation in the first quadrant will not be exactly compensated by the acceleration in the second, and that the retardation in the third will not be exactly compensated by the acceleration in the fourth. Therefore, at the end of the synodic revolution the moon will have an excess or deficiency of velocity. Its mean motion will then vary from one revolution to another, by reason of the ellipticity of its orbit. This variation will be periodic, like that just considered, and for similar reasons. The excess or deficiency of velocity at the close of any one revolution, will in time be compensated by an equal deficiency or excess occurring at the close of another revolution, when the sun has a certain different position with respect to the perigee of the moon's orbit.

563. We pass now to the consideration of the action of the radial force. The direct general effect of the radial force, is an alteration in the intensity of the moon's gravity towards the earth, and in its law of variation. Its specific effects are periodical variations in the magnitude, eccentricity and position of the orbit. As it is directed towards the earth, it will not disturb the equable description of areas. To discover the variations of this force we have only to discuss the general analytical expression for it, already investigated. It is,

$$\text{radial force} = \frac{m y (1 - 3 \cos^2 \varphi)}{a^3}.$$

We shall have, radial force = 0, when  $1 - 3 \cos^2 \varphi = 0$ , or when  $\cos \varphi = \pm \sqrt{\frac{1}{3}}$ . This value of  $\cos \varphi$  answers to four points lying on either side of the quadratures, and about  $35^\circ$  distant from them. When  $\cos \varphi$  is numerically greater than  $\sqrt{\frac{1}{3}}$ , the result will be negative, and when it is less than  $\sqrt{\frac{1}{3}}$ , the result will be positive. It follows, therefore, that the radial force increases the gravity of the moon in the quadratures, and for about  $35^\circ$  on each side of them, and that during the remainder of a synodic revolution it diminishes it.



When the moon is in quadratures  $\cos \varphi = 0$ , and

$$\text{radial force} = \frac{m y}{a^3} \dots (141).$$

In the syzgies, we have  $\cos \varphi = \pm 1$ , which gives

$$\text{radial force} = \frac{2 m y}{a^3} \dots (142).$$

It appears then that the diminution of the moon's gravity in the syzgies, is double of its increase in the quadratures.

We learn also from equations (141) and (142), that the radial force in the quadratures and syzgies varies directly as the distance; from which we conclude that the gravity of the moon varies at these points by a different law from that of the inverse squares. In the quadratures the gravity will be increased most at the greatest distance, when it is the least; and thus it will vary in a less rapid ratio than the square of the distance. In the syzgies it will be diminished most at the greatest distance, or when it is the least; and accordingly, at these points it will vary in a more rapid ratio than the square of the distance.

564. An easy investigation, with the aid of the differential calculus, proves that the mean diminution of the moon's gravity from the sun's action is  $\frac{m r}{2 a^3}$ ;  $r$  representing in this case the mean distance of the moon from the earth. The value of this expression is readily found to be equal to about the 368th part of the whole gravity of the moon to the earth.

In consequence of this diminution, the moon must describe her orbit at a greater distance from the earth, with a less angular velocity, and in a longer time, than if she were acted on only by the attraction of the earth.

565. The radial force of the sun alters the eccentricity of the moon's orbit, and differently in different revolutions of the moon, according to the position of the line of syzgies with respect to the line of apsides. When these lines are coincident, the eccentricity is increased. For, suppose P M A N (Fig. 75) to be the elliptic orbit of the moon that would be described under the influence of a force varying inversely as the square of the distance. In going from the apogee to the perigee, the gravity will increase in a greater ratio than that of the inverse square



of the distance; the true orbit will therefore fall within the ellipse, and the perigean distance ( $E P'$ ) will be less than for the ellipse. Consequently, the eccentricity will increase so much the more as the major axis diminishes. On the other hand, in going from the perigee to the apogee, the gravity will decrease in a greater ratio than the inverse square of the distance, and the moon will consequently recede farther from the earth, than if it were not disturbed by the sun. Therefore, in this half of the orbit the eccentricity will also be increased. When the apsides are in quadratures, the eccentricity will be diminished; for, the gravity will then vary from the apogee to the perigee, and from the perigee to the apogee, in a less ratio than that of the inverse squares; and therefore the results will be contrary to those just obtained. The eccentricity will have its maximum value when the apsides are in syzgies, and its minimum when they are in quadratures; for, in every other position of the line of apsides with respect to the line of syzgies, the radial force in the apogee and perigee will be less than in these positions (equa. 137), and therefore alter less the proportional gravity of the moon in the apogee and perigee. It is evident, from the gradual decrease of the radial force, as we recede from the syzgies and quadratures, that the eccentricity will continually diminish in the progress of the apsides from the syzgies to the quadratures, and that it will continually increase from the quadratures to the syzgies.

The change in the eccentricity of the moon's orbit, thus produced, will be attended with a corresponding change in the equation of the centre, and thus of the longitude. And this change is the conspicuous inequality of the moon known by the name of *Evection*, (Art. 272).

566. The radial force also produces a motion of the line of apsides. If the moon was only acted upon by the attraction of the earth, its orbit would be an ellipse, and the motion from one apsis to another, or, in other words, from one point where the orbit cuts the radius vector at right angles to the other, would be  $180^\circ$ . In point of fact, however, the gravity due to the earth's attraction is constantly either diminished or increased by the radial disturbing force of the sun, and therefore its true orbit must continually deviate from the ellipse that would be described under the sole

action of the earth's attraction. When from the action of this force there is a diminution of the moon's gravity, she will continually recede from the ellipse in question, her path will be less bent, and she must therefore move through a greater angular distance, before the central force will have deflected her course into a direction at right angles to the radius vector. Accordingly she will move through a greater angular distance than  $180^\circ$  in going from one apsis to another, and thus the apsides will advance. On the other hand, when the same force increases the moon's gravity, her path will fall within the ellipse, its curvature will be increased, and therefore it will be brought to intersect the radius vector at right angles, at a less angular distance. In this case, therefore, the apsides will move backwards. Now, we have shown, (Art. 563), that the radial disturbing force of the sun alternately diminishes and increases the moon's gravity to the earth. It follows, therefore, that the motion of the apsides will be alternately direct and retrograde; but since, as has been shown, (Art. 563), the diminution subsists during a longer part of the moon's revolution, and is moreover greater than the increase, the direct motion will exceed the retrograde, and therefore in an entire revolution the apsides will advance.

567. The observed motion of the apsides of the moon's orbit is not, however, wholly produced by the radial disturbing force. It is in part due to the action of the tangential force. This force alters the centrifugal force of the moon, and thus changes its gravity towards the earth, at the same time with the radial force.

568. The elliptic form of the sun's orbit is the occasion of a change in the radial force, from which results a perturbation of longitude, called the *Annual Equation* (Art. 272). The mean diminution of the moon's gravity, arising from the action of the sun, or the mean radial force, is equal to  $\frac{m r}{2 a^3}$  (Art. 563).

Hence this diminution is inversely proportional to the cube of the sun's distance from the earth. Therefore, as the sun approaches the perigee of its orbit, its distance from the earth diminishing, the mean diminution of the moon's gravity to the earth will increase, and consequently

the moon's distance from the earth will become greater, and its motion slower, than it otherwise would be. The contrary will take place while the sun is moving from the perigee to the apogee.

569. The disturbing force perpendicular to the plane of the moon's orbit, produces a tendency in the moon to quit that plane, from which there results a change in the position of the line of the nodes, and a change in the inclination of the plane of the orbit to that of the ecliptic. If we examine the general expression for this force, viz :

$$\text{perpen. force} = \frac{3 m y \cos \varphi}{a^3} \sin N \sin I,$$

we see that for any given values of  $S$  and  $I$ , it will be zero in the quadratures, and have its greatest value in the syzgies; and that it will change its direction in the quadratures, lying, in the nearer half of the orbit, on the same of its plane as the sun, and in the more remote half, on the opposite side. We perceive also that it will be zero for every value of  $\varphi$ , or for every elongation of the moon, when the angle  $S$  is zero, that is, when the sun is in the plane of the orbit, and will attain its maximum, for any given elongation, when the line of direction of the sun is perpendicular to the line of nodes. It will also be the less, other things being the same, the smaller is the inclination  $I$ .

570. Now, let  $N M' R$  (Fig. 76) represent the orbit of the moon, and  $S$  the sun, supposed stationary, the line of the nodes being in quadratures; and let  $L, L'$  be the points of the orbit  $90^\circ$  distant from the nodes. The direction of the force, in the various points of the orbit, is indicated by the arrows drawn in the figure. When the moon is at any point  $M'$  between  $L$  and the descending node  $N'$ , she will be drawn out of the plane in which she is moving by the disturbing force  $M' K'$ , and compelled to move in a line such as  $M' t'$ . The node  $N'$  will, therefore, retrograde to some point  $n'$ . When she is at any point  $M$ , moving from the ascending node  $N$  towards  $L$ , her course will be changed to the line  $M t$ , lying, like the line  $M' t'$ , below the orbit, which being produced backwards, meets the plane of the ecliptic in some point  $n$ , behind  $N$ . The nodes, therefore, retrograde in this position of the moon, as well as in the former. When the moon is in the half  $N' L' N$  of the orbit, lying below the ecliptic, the absolute direction of the dis-

turbing force will be reversed, and thus its tendency will be the same as before, namely, to draw the moon towards the ecliptic. It follows, therefore, that throughout this half of the orbit, as in the other, the motion of the nodes will be retrograde. Accordingly, when the nodes are in quadratures, or  $90^\circ$  distant from the sun, they will retrograde during every part of the revolution of the moon.

571. Suppose the sun now to be fixed on the line of nodes, or the nodes to be in syzgies. In this case the perpendicular force will be zero, (Art. 569), and, therefore, there will be no disturbance of the plane of the moon's orbit.

572. Next, let the situation of the sun be intermediate between the two just considered, as represented in Figs. 76 and 77. The effect of the disturbing force will be the same as in the first situation from the quadrature  $q$  (Fig. 76) to the node  $N'$ , and from the quadrature  $q'$  to the node  $N$ . But throughout the arcs  $Nq$ ,  $N'q'$  the direction of the force, and therefore the effects, will be reversed. The node will then retrograde, as before, while the moon moves over the arcs  $qN'$  and  $q'N$ , and advance while she is in the arcs  $Nq$ ,  $N'q'$ . But as the force is greatest over the arcs  $qN'$ ,  $q'N$ , which contain the syzgies, (Art. 569), and as these arcs are also longer than the arcs  $Nq$ ,  $N'q'$ , the node will, on the whole, retrograde each revolution. The velocity of retrogradation will, however, be less than when the nodes are in quadratures, and proportionably less, as the distance of the sun from this position is greater.

In the position represented in Fig. 77, a direct motion will take place over the arcs  $q'N'$  and  $qN$ ; but as  $Nq'$ , and  $N'q$ , the arcs of retrograde motion, are of greater extent than  $q'N'$ , and  $qN$ , and moreover contain the syzgies, the retrograde motion in each revolution must exceed the direct, as before.

If we suppose the sun to be situated on the other side of the line of nodes, the effect of the disturbing force will obviously be the same in any one position of the sun, as in the position diametrically opposite to it. It appears, then, that the line of the nodes has a retrograde motion in every possible position of the sun.

573. We have thus far supposed the sun to remain stationary in the various positions in which we have supposed it, during the revolution of the moon. It remains, then, to consider the effect of the



sun's motion in this interval. And, first, it is plain, that, as the sun advances from S towards N', (Fig. 76), the arcs N q, N' q' will increase, and the arcs q N' and q' N diminish; from which it appears, that, during the advance of the sun from the point 90° behind the descending node to this node, its motion in the course of each revolution of the moon will cause the retrograde motion of the node to be slower than it otherwise would be. While the sun moves from the ascending node to the 90° from it, the effect of its motion will obviously be just the reverse of this. During its passage from the descending to the ascending node, the effect will be the same in either quadrant, as in that diametrically opposite.

The variation in the intensity of the perpendicular force, conspires with the difference of situation of the sun and its motion during a revolution of the moon, in diminishing or increasing, as the case may be, the velocity of retrogradation of the nodes.

574. Let us now treat of the change of the inclination of the orbit, resulting from the disturbing action of the sun. And, first, if we refer to Fig. 76, we shall see that when the nodes are in quadrature, the inclination will diminish while the moon is moving from the ascending node N to the point L 90° distant from it, and increase while she is moving from L to the other node N'. In the other half of the orbit, the tendency of the disturbing force is the same, (Art. 570); and, therefore, while the moon is moving from N' to L', the inclination will diminish, and while she is moving from L' to N, it will increase. The diminutions and increments will compensate each other, and the original inclination will be regained at the close of the revolution.

575. When the nodes are in syzgies there will be no change of inclination (Art. 569).

576. In the situations of the sun, represented in Figs. 76 and 77, the inclination will decrease from q to L and from q' to L', and increase from L to q' and from L' to q, the effects being the same as when the nodes are in quadratures over the arcs q L and L N' in Fig. 76, and N L and L q' in Fig. 77, and being reversed over the arcs N q and N' q' in Fig. 76, and q N and q' N' in Fig. 77. When the sun has the position represented in Fig. 76, the arcs of increase L q' and L' q will be greater than the arcs of diminution q L and q' L'. The disturbing force will also be greater in the former arcs than in the latter. In the position



supposed, therefore, there will be, on the whole, an increase of inclination every revolution. When the sun is in the position represented in Fig. 77, the arcs of diminution  $qL$  and  $q'L'$  will be the greater; and the force in them will also be the greater. In this case, therefore, there will be a diminution of the inclination each revolution of the moon.

When the sun is on the other side of the line of nodes, the results will be the same as in the positions diametrically opposite.

577. To inquire now into the consequences of the sun's motion during the revolution of the moon. As the sun moves from  $S$ , towards  $N'$  (Fig. 76) the arcs  $Lq'$ ,  $L'q$ , over which there is an increase of the inclination, will increase; and the arcs  $qL$ ,  $q'L'$ , over which there is a diminution, will diminish. The motion of the sun will, therefore, in approaching the descending node, render the increase of the inclination each revolution of the moon greater than it otherwise would be. When the sun is receding from the ascending node, the corresponding arcs will experience corresponding changes, and therefore the diminution will now be less than if the sun were stationary.

The results will be similar for the opposite quadrants on the other side of the line of nodes.

578. Since the inclination diminishes as the sun recedes from either node, and increases as it approaches either node, it will be the least when the nodes are in quadratures, and the greatest when they are in syzgies.

579. The perturbations of the elliptic motion of the moon, comprising inequalities of orbit longitude, and variations in the form and position of the orbit, which have now been under consideration, depend upon the configurations of the sun and moon, with respect to each other, the perigee of each orbit, and the node of the moon's orbit. Their effects will disappear when the configurations upon which they depend become the same. They are therefore *periodical*.

580. The perturbations of the motions of a planet, produced by the action of another planet, are precisely analogous to the perturbations of the motions of the moon, produced by the action of the sun. The disturbing forces are obviously of the

same kind, and they are subject to variations from precisely similar causes. But, owing to the smallness of the masses of the planets and their great distances, their disturbing forces are much more minute than the disturbing force of the sun. From this cause, together with the slow relative motion of the disturbing and disturbed body, the motion of the apsides and nodes, and the accompanying variations of eccentricity and inclination, are very much more gradual in the case of the planets than in the case of the moon. Their periods comprise many thousands of years, and on this account they are called *Secular Motions* or *Variations*. In consequence of the greater feebleness of the disturbing forces, the periodical inequalities are also much less in amount. Moreover, as the motion of a planet is much slower than that of the moon, and as the variations of its orbit are more gradual than those of the lunar orbit, the compensations produced by a change of configurations are much more slowly effected, and thus the periods of the inequalities are much longer.

581. The motions of the moon would be subject to no secular variations, if the apparent orbit of the sun were unchangeable; but the secular variation of the eccentricity of the sun's orbit, which answers to an equal variation of the eccentricity of the earth's orbit, that is produced by the action of the planets, gives rise to a secular inequality in the motion of the moon, called the *Acceleration of the Moon*. This inequality was discovered from observation. Its physical cause was first made known by Laplace.

## CHAPTER XXIII.

OF THE RELATIVE MASSES AND DENSITIES OF THE SUN,  
MOON, AND PLANETS; AND OF THE RELATIVE INTENSITY  
OF THE GRAVITY AT THEIR SURFACE.

582. The perturbations which a planet produces in the motions of the other planets, depend for their amount chiefly upon the ratio of the mass of the planet to the mass of the sun, and the ratio of the distance of the planet from the sun to the distance of the planet disturbed from the same body. Now, the ratio of the distances is known by the methods of Plane Astronomy; consequently, the observed amount of the perturbations ought to make known the ratio of the masses, the only unknown element upon which it depends.

This is one method of determining the masses of the planets. The masses of those planets which have satellites may be found by another and simpler method. The theory of gravitation has furnished the following equation, from which they may easily be derived :

$$\frac{M + m}{1 + M} = \frac{d^3 P^2}{D^3 p^2},$$

in which 1 denotes the mass of the sun,  $M$  the mass of the planet,  $m$  the mass of one of its satellites,  $D$  the mean distance of the planet from the sun,  $d$  the mean distance of the satellite from the planet, and  $P$  and  $p$  the periodic times of the planet and satellites respectively. As the mass of the satellite is small compared with that of the planet, and the mass of the planet is small compared with that of the sun, they may be neglected in the above equation with but little error; and thus we have,

$$M = \frac{d^3 P^2}{D^3 p^2} \text{ (very nearly).}$$

583. The second column of Table IV exhibits the relative masses of the sun, moon, and planets, according to the most received determinations, that of the earth being denoted by 1.

584. The quantities of matter of the sun, moon, and planets, as

well as their bulks, being known, their densities may be easily computed; for, the densities of bodies are proportional to their quantities of matter divided by their bulks. The third column of Table IV contains the densities of the sun, moon, and planets, that of the earth being denoted by 1. It will be seen on inspecting it, that, for the most part, the densities of the planets decrease as we recede from the sun.

585. The relative intensity of the gravity at the surface of the sun, moon, and planets, may also readily be found, when the masses and bulks of these bodies are known. For, supposing them to be spherical, and not to rotate on their axes, the gravity at their surface will be directly as their masses, and inversely as the squares of their radii, or, in other words, proportional to their masses divided by the squares of their radii. The centrifugal force at the surface of a planet, generated by its rotation on its axis, diminishes the gravity due to the attraction of the matter of the planet. The diminution thus produced on any of the planets is not, however, very considerable. The method of determining the centrifugal force at the surface of a body in rotation, is given in treatises on Mechanics. (See Courtenay's Mechanics, pages 250 and 251).

The fourth column of Table IV exhibits the relative intensity of the gravity at the surface of the sun, moon, and planets, that at the surface of the earth being denoted by 1.

---

## CHAPTER XXIV.

OF THE FIGURE AND ROTATION OF THE EARTH; AND OF THE PRECESSION OF THE EQUINOXES AND NUTATION.

586. We have already seen (Art. 145) that measurements made upon the earth's surface, establish that the figure of the earth is that of an oblate spheroid, and that the oblateness at the poles is about  $\frac{1}{305}$ .

587. From the amount and law of the variation of the force of gravity upon the earth's surface, ascertained by observations upon the length of the seconds' pendulum, it is proved that the matter of the earth is not homogeneous, but denser towards the centre, and that it is arranged in concentric strata of nearly an elliptical form and uniform density.

The fact of the greater density of the earth towards its centre, has also been established by observations upon the deviation of a plumb line from the vertical, produced by the attraction of a mountain. Such observations were made for the purpose of determining the mean density of the earth by Dr. Maskelyne, on the sides of the mountain Schehallien in Scotland. The observed deviation of the plumb line made known the ratio of the attraction of the mountain to that of the whole earth, and thus the relative quantities of matter in the mountain and earth. These being ascertained, and the figure and bulk of the mountain having been determined by a survey, the relative density of the earth and mountain became known by the principle mentioned in Art. 584, and thence the actual density of the earth, the density of the mountain having been found by experiment. The result was, that the mean density of the earth is 4.95, the density of water being 1.

588. The spheroidal form of the surface of the earth and of its internal strata is easily accounted for, if we suppose the earth to have been originally in a fluid state. The tendency of the mutual attraction of its particles would be to give it a spherical form; but, by virtue of its rotation, all its particles, except those lying immediately on the axis, would be animated by a centrifugal force increasing with their distance from the axis. If, therefore, we conceive of two columns of fluid extending to the earth's centre, one from near the equator, and the other from near either pole, the weight of the former would by reason of the centrifugal force be less than that of the latter. In order, then, that they may sustain each other in equilibrio, that near the equator must increase in length, and that near the pole diminish. As this would be true at the same time for every pair of columns situated as we have supposed, the surface of the whole body of fluid about the poles must fall, and that of the fluid about the equator rise. In this manner the earth would become flattened at the poles and protuberant at the equator.



589. Upon a strict investigation it appears that a homogeneous fluid of the same mean density with the earth, and rotating on its axis at the same rate that the earth does, would be in equilibrium, if it had the figure of an oblate spheroid, of which the axis was to the equatorial diameter as 229 to 230, or of which the oblateness was  $\frac{1}{229}$ . If the fluid mass supposed to rotate on its axis be not homogeneous, but be composed of strata that increase in density from the surface to the centre, the solid of equilibrium will still be an elliptic spheroid, but the oblateness will be less than when the fluid is homogeneous.

590. The time of the earth's rotation, as well as the position of its axis, would change if any variation should take place in the distribution of the matter of the earth, or in case of the impact of a foreign body.

If any portion of matter be, from any cause, made to approach the axis, its velocity will be diminished, and the velocity lost being imparted to the mass, will tend to accelerate the rotation. If any portion of matter be made to recede from the axis, the opposite effect will be produced, or the rotation will be retarded. In point of fact, the changes that take place in the position of the matter of the earth, whether from the washing of rains upon the sides of mountains, or evaporation, or any other known cause, are not sufficient ever to produce any sensible alteration in the circumstances of the earth's rotation on its axis.

591. It is ascertained from direct observation, that there has in reality been no perceptible change in the period of the earth's rotation since the time of Hipparchus, 120 years before the beginning of the present era. We may therefore conclude, *à posteriori*, that there has been no material change in the form and dimensions of the earth in this interval.

592. Were the axis of the earth to experience any change of position with respect to the matter of the earth, the latitudes of places would be altered. A motion of 200 feet might increase or diminish the latitude of a place to the amount of 2", an angle which can be measured by modern instruments. Now, in point of fact, the latitudes of places have not sensibly varied since their first determination with accurate instruments; therefore, in this interval the axis of the earth cannot have changed. Indeed, since the earth's surface and its internal strata are arranged symmetri-

cally with respect to the present axis of rotation, it is to be inferred that this axis is the same as that which obtained at the epoch when the matter of the earth changed from a fluid to a solid state.

593. The motions of the earth's axis, along with the whole body of the earth, which give rise to the Precession of the Equinoxes and Nutation, are consequences of the spheroidal form of the earth, inasmuch as they are produced by the actions of the sun and moon upon that portion of the matter of the earth which lies on the outside of a sphere conceived to be described about the earth's axis. The physical theory of the phenomena in question is ana'ogous to that of the retrogradation of the moon's nodes. The sun produces a retrograde movement of the points in which the circle described by each particle of the protuberant mass cuts the plane of the ecliptic, as it does of the moon's nodes; the effect produced is, however, exceedingly small, by reason of the inertia of the interior spherical mass connected with the extenal mass upon which the action takes place. The moon, in like manner, occasions a retrograde movement of the nodes of the same particles on the plane of its orbit. The actions of the sun and moon will not be the same each revolution of a particle. That of the sun will vary during the year with the angular distance of the sun from the node (Art. 569); and that of the moon will vary during each month with the distance of the moon from the node, and also during a revolution of the nodes of the moon's orbit by reason of the change in the inclination of the orbit to the equator. The mean effect of both bodies is the *precession*; the inequality resulting from the change in the sun's action during the year is the *solar nutation*; and the inequality consequent upon the retrogradation of the moon's nodes is the *lunar nutation*; or the chief part of it: the change in the position of the equinox occasioned by the moon's revolution, never exceeds  $\frac{1}{4}$  of a second of an arc; and the change of the obliquity of the ecliptic from this cause is still less.

## CHAPTER XXV.

## OF THE TIDES.

594. The alternate rise and fall of the surface of the ocean twice in the course of a lunar day, or about 25 hours, is the phenomenon known by the name of the *Tides*. The rise of the water is called the *Flood Tide*, and the fall the *Ebb Tide*.

595. The interval between one high water and the next, is, at a mean, half a mean lunar day, or 12h. 25m. 14s. Low water has place nearly, but not exactly, at the middle of this interval; the tide, in general, employing nine or ten minutes more in ebbing than in flowing. As the interval between one period of high water and the second following one is a lunar day, or 1d. 0h. 50m. 28s., the *retardation* in the time of high water from one day to another is 50m. 28s., in its mean state.

596. The time of high water is mainly dependent upon the position of the moon, being always at any given place about the same length of time after the moon's passage over the superior or inferior meridian. As to the length of the interval between the two periods, at different places, in the open sea it is only from two to three hours; but on the shores of continents, and in rivers, where the water meets with obstructions, it is very different at different places, and, in some instances, is of such length that the time of high water seems to precede the moon's passage.

597. The height of the tide at high water is not always the same, but varies from day to day; and these variations have an evident relation to the phases of the moon. It is greatest at the syzgies; after which it diminishes and becomes the least at the quadratures.

598. The tides, about the time of the syzgies, are called the *Spring Tides*; and those about the time of quadratures are called the *Neap Tides*.

The highest of the spring tides is not that which has place nearest the syzgy, but is in general the third, and in some in-

stances, the fourth following tide. In like manner the lowest of the neap tides is the third or fourth tide after the quadrature.

The spring tides are, in general, about twice the height of the neap tides. At Brest, in France, the former rise to the height of 19.3 feet, and the latter only to 9.2 feet. In the Pacific Ocean the highest of the tides of the syzigies is 5 feet, and the lowest of the tides of the quadratures is between 2 and 2.5 feet.

599. The tides are also affected by the declinations of the sun and moon; for they diminish the tides of the syzigies, which occur at the equinoxes; and augment the tides of the quadratures, which occur at the solstices. Also, when the moon or the sun are out of the equator, the evening and morning tides differ somewhat in height. At Brest, in the syzigies of the summer solstice, the tides of the morning of the first and second day after the syzgy are smaller than those of the evening by 6.6 inches. They are greater by the same quantity, in the syzigies of the summer solstice.

600. The distance of the moon from the earth has also a sensible influence upon the tides. In general, they increase and diminish as the distance increases or diminishes, but in a more rapid ratio.

601. The daily retardation of the time of high water varies with the phases of the moon. It is at its minimum towards the syzigies, when the tides are at their maximum; and it is then about 40m. But, towards the quadratures, when the tides are at their minimum, the retardation is the greatest possible; and amounts to about 1h. 15m.

The variation in the distance of the sun and moon from the earth, (and particularly the moon), has an influence also on this retardation.

The daily retardation of the tides varies likewise with the declination of the sun and moon.

602. The facts which have been enumerated clearly indicate that the tides are produced by the actions of the sun and moon upon the waters of the ocean; but in a greater degree by the action of the moon. To explain them, let us suppose at first that the whole surface of the earth is covered with water. It has been shown (Art. 563) that the sun's action increases or dimin-



ishes the moon's gravity to the earth, according to her position with respect to the line of syzgies or quadratures, or, in other words, according to her elongation from the sun. In a similar manner will the moon's action increase or diminish the gravity of a particle of matter at the earth's surface, according to its elongation from the moon, as seen from the earth's centre; for, any particle of matter upon the earth's surface is attracted towards the earth's centre, just as if the whole mass of the earth were concentrated at its centre; and the whole earth is attracted by the moon, just as it would be if its matter were concentrated at its centre. If we conceive a plane to pass through the centre of the earth, at right angles to the line joining the centres of the earth and moon; within about  $35^{\circ}$  of this plane on each side, the gravity at the surface will be increased (Art. 563); and at the remaining parts, that is, for about  $55^{\circ}$  around the points in which the line of the centres intersects the surface, the gravity will be diminished.

The equilibrium of the waters upon the surface of the earth will in consequence be disturbed. Around the points just mentioned, where the gravity is diminished, they will rise, and in the middle parts between these points, where the gravity is increased, they will fall. In consequence of the earth's diurnal rotation, the parts of the surface, at which the rise and fall of the water will take place, will be continually changing. Were the entire rise and fall produced instantaneously, the points of highest water would constantly be the precise points in which the line of the centres of the moon and earth intersects the surface, and it would always be high water on the meridian passing through these points, both in the hemisphere where the moon is, and in the opposite one. On the west side of this meridian, the tide would be flowing; on the east side of it, it would be ebbing; and on the meridian at right angles to the same, it would be low water. But it is plain that the effects of the moon's action will not be instantaneously produced, and therefore that the points of highest water will fall behind the moon. It appears from observation, that in the open sea the meridian of high water is about  $30^{\circ}$  to the east of the moon.

The great tide wave thus raised by the moon, and which follows it in its diurnal motion, will be a mere undulation, or alternate rise and fall of the water, without any progressive motion,



if, as we have supposed, it is no where obstructed by shallows, islands, or the shores of continents.

603. It is evident that the sun will produce precisely similar effects with the moon, and will raise a tide wave similar to the lunar tide wave, which will follow it in its diurnal motion.

604. To show that the effects of the sun are less in degree than those of the moon, let us take the general expression for the change of the moon's gravity arising from the action of the sun, namely,

$$\frac{m}{a^3} \times y (1 - 3 \cos^2 \varphi) \dots (a),$$

in which  $m$  denotes the mass of the sun,  $a$  its distance (the mean distance of the moon being taken as 1),  $y$  the distance of the moon in its given position, and  $\varphi$  its elongation from the sun as seen from the earth's centre. This formula will serve to express the change in the gravity of a particle of matter upon the earth's surface, produced by the sun's action, if we take  $m$  = the mass of the sun, as before,  $a$  = its distance expressed in terms of the radius of the earth as unity,  $y$  = the distance of the particle from the centre of the earth, and  $\varphi$  = its elongation from the sun as seen from the earth's centre. If we designate the corresponding quantities for the moon by  $m'$ ,  $a'$ ,  $y$ ,  $\varphi$ , we shall have for the change of the gravity of a particle, produced by the moon's action,

$$\frac{m'}{a'^3} \times y (1 - 3 \cos^2 \varphi) \dots (b).$$

For particles at equal elongations from the sun and moon, we shall have  $\varphi$  the same in expressions (a), and (b), and  $y$  may be regarded as the same without material error. For such particles, then, the alterations of the gravity, produced by the sun and moon, will bear the same ratio to each other as the quantities

$\frac{m}{a^3}$  and  $\frac{m'}{a'^3}$ . Now, if we give to  $m$ ,  $m'$ ,  $a$ ,  $a'$ , their values, we

shall find that the latter quantity is nearly three times greater than the former. Accordingly, the effect of the moon's action, at corresponding elongations of the particles, and therefore generally, is nearly three times greater than that of the sun.

605. The actual tide will be produced by the joint action of the sun and moon, or it may be regarded as the result of the combination of the lunar and solar tide waves.

At the time of the syzgies, the action of the sun and moon will be combined in producing the tides, both bodies tending to produce high as well as low water at the same places. But at the quadratures they will be in opposition to each other, the one tending to raise the surface of the water where the other tends to depress it, and *vice versa*. The tides should, therefore, be much higher at the syzgies than at the quadratures.

Between the syzgies and the quadratures the two bodies will neither directly conspire with each other, nor directly oppose each other, and tides of intermediate height will have place. The points of highest water will also, in the configuration supposed, neither be the vertices of the lunar nor of the solar tide wave, but certain points between them. This circumstance will occasion a variation in the length of the interval between the time of the moon's passage and the time of high water.

606. The effect of the moon's action being to that of the sun's nearly as 3 to 1, (Art. 604), the spring tides will be to the neap tides nearly as 2 to 1. For, let  $x$  = the effect of the moon, and  $y$  = the effect of the sun: then the ratio of  $x + y$  to  $x - y$  will be the ratio of the heights of the spring and neap tides. Now,

$$x = 3y, \text{ and thus } \frac{x + y}{x - y} = \frac{3y + y}{3y - y} = 2.$$

This result is conformable to observation.

607. The height of the tide, as well as the interval between the time of high water and that of the moon's meridian passage, will vary not only with the elongation of the moon from the sun, but also with the distance and declination of the moon and sun. For, expressions (a) and (b) show that the intensities of the moon's and sun's actions vary inversely as the cube of their distance; and the changes of the declinations of the two bodies must be attended with a change both in the absolute and relative situation of the vertices of the lunar and solar tide waves.

608. The laws of the tides, which would obtain on the hypothesis of the earth being covered entirely with water, are found to correspond only partially with those of the actual tides. The continents have a material influence upon the formation and propagation of the tide wave.

609. Professor Whewell infers, from a careful discussion of a great number of observations upon the tides, that the tide of the

Atlantic Ocean is, for the most part, produced by a derivative tide wave, sent off from the great wave which in the Southern Ocean follows the moon in its diurnal motion around the earth. This wave advances more rapidly in the open sea than along the coasts, where it meets with obstructions.

Where portions of the tide wave, extending from one point of the coast to another, become detached, and advance into a narrow space, particularly high tides will occur. In this way (as it is supposed) it happens that the tide rises at certain places in the bay of Fundy, to the height of 60 or 70 feet.

610. In channels peculiar tides occur in consequence of the meeting of the waves which enter the channels at their two extremities. Where the two waves meet in the same state, unusually high tides occur. This is observed to be the case at some points in the Irish Channel. In the port of Batsha, in Tonquin, the tides arrive by two channels, of such lengths that the two waves meet in opposite states, or that the flood tide arrives by one channel just as the ebb tide begins to leave by the other, and the consequence is, that there is neither high nor low water.

This is the case when the moon is in the equator. When she has a northern or southern declination, there is a small rise and fall of the water once in a lunar day, owing to the inequality of the morning and evening tides of the open sea.

611. Lakes and inland seas have no perceptible tides, for the reason that their extent is not sufficient to admit of any sensible inequality of gravity, as the result of the action of the moon.

612. The tides experienced in rivers and seas communicating with the ocean, are not produced by the direct actions of the sun and moon, but are waves propagated from the great wave of the open sea.

In rivers of considerable length, the ascending tides are encountered by those which are returning, so that a great variety of tides occur along their shores.

613. The mean interval between noon and the time of high water at any port, on the day of new or full moon, is called the *Establishment* of that port. It will be, approximately, the interval between the time of the meridian passage of the moon and the time of high water on any day of the month. To ob-

tain this interval for a given day more nearly, it is necessary to correct the establishment for the effects of the change of the distance and declination of the sun and moon, and of the change in the elongation of the moon from the sun. When it has been determined, by adding it to the time of the meridian passage of the moon, we have the time of the next high water.

# PART IV.

## ASTRONOMICAL PROBLEMS.

---

### EXPLANATIONS OF THE TABLES.

THE Tables which form a part of this work, and which are employed in the resolution of the following Problems, consist of Tables of the Sun, Tables of the Moon, Tables of the Mean Places of some of the Fixed Stars, Tables of Corrections for Refraction, Aberration and Nutation, and Auxiliary Tables.

The Tables of the Sun, which are from XVII to XXXIV inclusive, are, for the most part, abridged from Delambre's Solar Tables. The mean longitudes of the sun and of his perigee for the beginning of each year, found in Table XVIII, have been computed from the formulæ of Prof. Bessel, given in the Nautical Almanac of 1837. The Table of the Equation of Time was reduced from the table in the *Connaissance des Temps* of 1810, which is more accurate than Delambre's Table, this being in some instances liable to an error of 2 seconds. The Table of Nutation (Table XXVII) was extracted from Francœur's *Practical Astronomy*. The maximum of nutation of obliquity is taken at  $9''.25$ . The Tables of the Sun will give the sun's longitude from the mean equinox within a fraction of a second of the result obtained immediately from Delambre's Tables, as corrected by Bessel. The Tables of the Moon, which are from XXXV to LXXXV inclusive, are abridged and computed from Burckhardt's Tables of the Moon. To facilitate the determination of the hourly motions in longitude and latitude, the equations of the hourly motions have all been rendered positive, like those of the longitude. Some few new tables have been computed for the same purpose. The longitude and hourly motion



in longitude will very rarely differ from the results of Burckhardt's Tables more than  $0''.5$ , and never as much as  $1''$ . The error of the latitude and hourly motion in latitude will be still less. The other tables have been taken from some of the most approved modern Astronomical Works. (For the principles of the construction of the Tables, see Chap. X.)

Before entering upon the explanation of each of the tables, it will be proper to define a few terms that will be made use of in the sequel.

The given quantity with which a quantity is taken from a table, is called the *Argument*.

The angular arguments are expressed in some of the tables according to the sexagesimal division of the circle. In others, they are given in parts of the circle supposed to be divided into a 100, a 1000, or 10000 &c. parts.

Tables are of *Single* or *Double Entry*, according as they contain one or two arguments. The *Epoch* of a table, is the instant of time for which the quantities given by the table are computed. By the *Epoch* of a quantity, is meant the value of the quantity found for some chosen epoch, from which its value at other epochs is to be computed by means of its known rate of variation.

Table I, contains the latitudes, and longitudes from the meridian of Greenwich, of various conspicuous places in different parts of the earth. The longitudes serve to make known the time at any one of the places in the table, when that at any of the others is given. The latitude of a place is an important element in various astronomical calculations.

Table II, is a table of the Elements of the Orbits of the Planets with their secular variations, and serves to make known the elements at any given epoch different from that of the table. From these the elliptic place of the planets at the given epoch may be computed.

Table III, is a similar table for the Moon.

Tables IV, V, VI, VII, require no explanation.

Table VIII, gives the mean Astronomical Refractions; that is, the refractions which have place when the barometer stands at 30 inches, and the thermometer of Fahrenheit at  $50^{\circ}$ .

Table IX, contains the corrections of the Mean Refractions for

+ 1 inch in the barometer, and  $-1^{\circ}$  in the thermometer, from which the corrections to be applied, at any observed height of the barometer and thermometer, are easily derived.

Table X, gives the Parallax of the Sun for any given altitude on a given day of the year ; for reducing a solar observation made at the surface of the earth to what it would have been, if made at the centre.

Table XI, is designed to make known the Sun's Semi-diurnal Arc, answering to any given latitude, and to any given declination of the sun ; and thus the time of the sun's rising and setting, and the length of the day.

Table XII, serves to make known the value of the Equation of Time, with its essential sign, which is to be applied to the apparent time to convert it into the mean. If the sign of the equation taken from the table be changed, it will serve for the conversion of mean time into apparent. This table is constructed for the year 1840.

Table XIII, is to be used in connection with Table XII, when the given date is in any other year than 1840. It furnishes the Secular Variation of the Equation of Time, from which the proportional part of its variation in the interval between the given date and the epoch of Table XII is easily derived.

Table XIV, contains certain other Corrections to be applied to the equation of time taken from Table XII, when its exact value, to within a small fraction of a second, is desired.

Table XV, gives the Fraction of the Year, corresponding to each date. This table is useful, when quantities vary by known and uniform degrees, in deducing their values at any assumed time from their values at any other time.

Table XVI, is for converting Hours, Minutes, and Seconds into decimal parts of a Day.

Table XVII, is for converting Minutes and Seconds of a degree into the decimal division of the same. It will also serve for the conversion of minutes and seconds of time into decimal parts of an hour.

The last two tables will be found frequently useful in arithmetical operations.

Table XVIII, is a table of Epochs of the Sun's Mean Longitude, of the Longitude of the Perigee, and of the Arguments for

finding the small equations of the Sun's place. They are all calculated for the first of January of each year, at mean noon on the meridian of Greenwich. Argument I, is the mean longitude of the Moon minus that of the Sun ; Argument II, is the heliocentric longitude of the Earth ; Argument III, is the heliocentric longitude of Venus ; Argument IV, is the heliocentric longitude of Mars ; Argument V, is the heliocentric longitude of Jupiter ; Argument VI, is the mean anomaly of the Moon ; Argument VII, is the heliocentric longitude of Saturn ; and Argument N, is the supplement of the longitude of the Moon's Ascending Node. Argument I, is for the first part of the equation depending on the action of the Moon. Arguments I and VI, are the arguments for the remaining part of the lunar equation. Arguments II and III, are for the equation depending on the action of Venus ; Arguments II and IV, for the equation depending on the action of Mars ; Arguments II and V, for the equation depending on the action of Jupiter ; and Arguments II and VII, for the equation depending on the action of Saturn. Argument N, is the argument for the Nutation in longitude : it is also the argument for the Nutation in right ascension, and of the obliquity of the ecliptic.

Table XIX, shows the Motions of the Sun and Perigee, and the variations of the arguments, in the interval between the beginning of the year and the first of each month.

Table XX, shows the Motions of the Sun and Perigee, and the variations of the arguments, for Days and Hours.

Table XXI, gives the Sun's Motions for Minutes and Seconds. Tables XVIII to XXI, make known the mean longitude of the Sun from the mean equinox at any moment of time.

Table XXII. Mean Obliquity of the Ecliptic for the beginning of each year contained in the table. It is found for any intermediate time by a simple proportion.

Tables XXIII and XXIV, furnish the Sun's Hourly Motion and Semi-diameter.

Table XXV, is designed to make known the Equation of the Sun's Centre. When the equation has the negative sign, its supplement to 12s. is taken. This is to be added along with the other equations of longitude, and 12s. are to be subtracted from the sum. The signs of the argument are given both at the head

and foot of the columns. The numbers in the table are the values of the equation of the centre, or of its supplement, diminished by  $46''.1$ . This constant is subtracted from each value, to balance the different quantities added to the other equations of the longitude, in order to render them affirmative. The epoch of this table is the year 1840.

Table XXVI, gives the Secular Variation of the Equation of the Sun's Centre, from which the proportional part of the variation in the interval between the given date and the year 1840, may be derived.

Table XXVII, is for the Nutation in Longitude and Right Ascension, and of the Obliquity of the Ecliptic. The nutation in longitude and in right ascension, serve to transfer the origin of the longitude and right ascension from the mean to the true equinox. And the nutation of obliquity serves to change the mean into the true obliquity.

Tables XXVIII to XXXIII, give the Equations of the Sun's Longitude, due respectively to the attractions of the Moon, Venus, Jupiter, Mars, and Saturn.

Table XXXIV is for the variable part of the Sun's Aberration. The numbers have all been rendered positive by the addition of the constant  $0''.3$ .

Table XXXV, contains the Epochs of the Moon's Mean Longitude, and of the Arguments for finding the equations which are necessary in determining the True Longitude and Latitude of the Moon. They are all calculated for the first of January of each year, at mean noon on the meridian of Greenwich. The Argument for the Evection is diminished by  $30'$ ; the Anomaly by  $2^\circ$ ; the Argument for the Variation by  $9^\circ$ ; and the Supplement of the Node is increased by  $7'$ . This is done to balance the quantities which are added to the different equations in order to render them affirmative.

Tables XXXVI to XL, inclusive, give the Motions of the Moon, and the variations of the arguments for Months, Days, Hours, Minutes, and Seconds; and, together with Table XXXV, are for finding the Moon's Mean Longitude and the Arguments at any assumed moment of time.

Tables XLI to LIII, inclusive, give the various Equations of the Moon's Longitude. It is to be observed, with respect to Table



XLI, that the right hand figure of the argument is supposed to be dropped. But when the greatest attainable accuracy is desired, it can be retained, and a cypher conceived to be written after the numbers in the columns of Arguments in the table. In Tables L, LI, LII, and LV, the degrees will be found by referring to the head or foot of the column. (See Problem II, Note 2).

Table LIV, is for the Nutation of the Moon's Longitude.

Tables LV to LIX, inclusive, are for finding the Latitude of the Moon.

Tables LX to LXIII, inclusive, are for the Equatorial Parallax of the Moon.

Table LXIV, furnishes the Reductions of Parallax and of the Latitude of a Place. The reduction of parallax is for obtaining the parallax at any given place from the equatorial parallax. The reduction of latitude is for reducing the true latitude of a place, as determined by observation, to the corresponding latitude, on the supposition of the earth being a sphere. The ellipticity to which the numbers in the table correspond is  $\frac{1}{300}$ .

Tables LXV and LXVI. Moon's Semi-diameter, and the Augmentation of the Semi-diameter depending on the altitude.

Tables LVII to LXXXV, inclusive, are for finding the Hourly Motions of the Moon in Longitude and Latitude.

Table LXXXVI. Mean New Moons, and the Arguments for the Equations for New and Full Moon in January. The time of mean new moon in January of each year has been diminished by 15 hours, which has been added to the equations in Table LXXXIX. Thus, 4h. 20m. has been added to equation I; 10h. 10m. to equation II; 10m. to equation III; and 20m. to equation IV.

Tables LXXXVII and LXXXVIII, are used with the preceding in finding the Approximate Time of Mean New or Full Moon in any given month of the year.

Table LXXXIX, furnishes the Equations for finding the Approximate Time of New or Full Moon.

Table XC, contains the Mean Right Ascensions and Declinations of 50 principal Fixed Stars, for the beginning of the year 1840, with their Annual Variations.

Table XCI, is for finding the Aberration and Nutation of the Stars in the preceding catalogue.



Table XCII, contains the Mean Longitudes and Latitudes of some of the principal Fixed Stars, for the beginning of the year 1840, with their Annual Variations.

Tables XCIII, XCIV, XCV. Second, Third, and Fourth Differences. These tables are useful for finding, from the Nautical Almanac, the moon's longitude or latitude for any time between noon and midnight.

Table XCVI. Logistical Logarithms. This table is convenient in working proportions when the terms are minutes and seconds, or degrees and minutes, or hours and minutes,—especially when the first term is 1h. or 60m.

*To find the logistical logarithm of a number composed of minutes and seconds, or degrees and minutes of an arc; or of minutes and seconds, or hours and minutes of time.*

1. If the number consists of minutes and seconds, at the top or bottom of the table seek for the minutes, and in the same column opposite the seconds in the left-hand column will be found the logistical logarithm.

2. If the number is composed of hours and minutes, the hours must be used as if they were minutes, and the minutes as if they were seconds.

3. If the number is composed of degrees and minutes, the degrees must be used as if they were minutes, and the minutes as if they were seconds.

*To find the logistical logarithm of a number less than 3600.*

Seek in the second line of the table from the top the number next less than the given number, and the remainder, or the complement to the given number, in the first column on the left: then, in the column of the first number, and opposite the complement, will be found the logistical logarithm of the sum. Thus, to obtain the logarithm of 1531, we seek for the column of 1500, and opposite 31 we find 3713.

## PROBLEM I.

*To work, by logistical logarithms, a proportion the terms of which are degrees and minutes, or minutes and seconds of an arc; or hours and minutes, or minutes and seconds of time.*

With the degrees or minutes at the top, and minutes or seconds at the side, or if a term consists of hours and minutes, or minutes and seconds, with the hours or minutes at the top, and minutes or seconds at the side, take from Table XCVI the logistical logarithms of the three given terms; add together the logistical logarithms of the second and third terms and the arithmetical complement of that of the first term, rejecting 10 from the index.\* The result will be the logistical logarithm of the fourth term, with which take it from the table.

Note 1. The logistical logarithm of 60' is 0.

Note 2. If the second or third term contains tenths of seconds, (or tenths of minutes, when it consists of degrees and minutes), and is less than 6', or 6°, multiply it by 10, and employ the logarithm of the product in place of that of the term itself. The result obtained by the table divided by 10 will be the fourth term of the proportion, and will be exact to tenths.

Note 3. If none of the terms contain tenths of minutes or seconds, and it is desired to obtain a result exact to tenths, diminish the index of the logistical of the fourth term by 1, and cut off the right-hand figure of the number found from the table, for tenths.

Exam. 1. When the moon's hourly motion is 30' 12'', what is its motion in 16m. 24s.?

As 60m.	-	-	-	0
: 30' 12''	-	-	-	2981
:: 16m. 24s.	-	-	-	5633
: 8' 15''	-	-	-	8614

---

\* Instead of adding the arithmetical complement of the logarithm of the first term, the logarithm itself may be subtracted from the sum of the logarithms of the other two terms.

2. If the moon's declination change  $1^{\circ} 31'$  in 12 hours, what will be the change in 7h. 42m.?

As 12h.	-	-	Ar. Co.	9.3010
: $1^{\circ} 31'$	-	-	-	1.5973
:: 7h. 42m.	-	-	-	8917
				<hr/>
: $0^{\circ} 58'$	-	-	-	1.7900

3. When the moon's hourly motion in latitude is  $2' 26''.8$ , what is its motion in 36m. 22s.?

$2' 26''.8$				
60				
<hr/>				
$146''.8$				
10				
<hr/>				
1468	-	-	As 60m.	- 0
			: $1468''$	- 3896
			:: 36m. 22s.	- 2174
				<hr/>
			: $890''$	- 6070

Ans.  $1' 29''.0$ .

4. When the sun's hourly motion in longitude is  $2' 28''$ , what is its motion 49m. 11s.?

Ans.  $2' 1''$ .

5. If the sun's declination changes  $16' 33''$  in 24 hours, what will be the change in 14h. 18m.?

Ans.  $9' 52''$ .

6. If the moon's declination change  $54''.7$  in one hour, what will be the change in 52m. 18s.?

Ans.  $47''.7$ .

## PROBLEM II.

*To take from a table the quantity corresponding to a given value of the argument, or to given values of the arguments of the table.*

Case 1. *When quantities are given in the table for each sign and degree of the argument.*

With the signs of the given argument at the top or bottom, and the degrees at the side, (at the left side, if the signs are found at the top; at the right side, if they are found at the bottom), take out the corresponding quantity. Also take the difference between this quantity and the next following one in the table, and say,  $60'$ : this difference :: odd minutes and seconds of

given argument : a fourth term. This fourth term, added to the quantity taken out, when the quantities in the table are increasing, but subtracted when they are decreasing, will give the required quantity.

Note 1. When the quantities change but little from degree to degree, the required quantity may frequently be estimated without the trouble of making a proportion.

Note 2. In some of the tables the degrees or signs of the quantity sought, are to be had by referring to the head or foot of the column in which the minutes and seconds are found. (See Tables L, LI, LII, and LV.) The degrees there found are to be taken, if no horizontal mark intervenes ; otherwise, they are to be increased or diminished by  $1^\circ$ , or  $2^\circ$ , according as one or two marks intervene. They are to be increased, or diminished, according as their number is less or greater than the number of degrees at the other end of the column.

Note 3. If, as is the case with some of the tables, the quantities in the table have an algebraic sign prefixed to them, neglect the consideration of the sign in determining the correction to be applied to the quantity first taken out, and proceed according to the rule above given. The result will have the sign of the quantity first taken out. It is to be observed, however, that if the two consecutive quantities chance to have opposite signs, their numerical sum is to be taken instead of their difference ; also that the quantity sought will, in every such instance, be the numerical difference between the correction and the quantity first taken out, and, according as the correction is less or greater than this quantity, is to be affected with the same or the opposite sign.

Exam. 1. Given the argument  $7^s 6^\circ 24' 36''$ , to find the corresponding quantity in Table L.

$7^s 6^\circ$  gives  $0^\circ 43' 17''.4$ .

The difference between  $0^\circ 43' 17''.4$  and the next following quantity in the table is  $1' 7''.3$ .

$60' : 1' 7''.3 :: 24' 36'' : 27''.6.*$

---

\* The student can work the proportion either by common arithmetic, or by logistical logarithms, as he may prefer. In working this and all similar proportions by the arithmetical method, the seconds of the argument may be converted into the equivalent decimal part of a minute by means of Table XVII. It will be sufficient to take the fraction to the nearest tenth.

From	0° 43' 17".4
Take	27 .6
	0 42 49 .8

2. Given the argument  $2^s. 18^\circ 41' 20''$ , to find the corresponding quantity in Table XXV.

$$2^s. 18^\circ \text{ gives } 1^\circ 52' 32''.5.$$

The difference between  $1^\circ 52' 32''.5$  and the next following quantity in the table is  $21''.8$ .

$$60' : 21''.8 :: 41' 20'' : 15''.0.$$

To	1° 52' 32''.5
Add	15 .0
	1 52 47 .5

3. Given the argument  $9^s. 2^\circ 13' 33''$ , to find the corresponding quantity in Table XII.

$$9^s. 2^\circ \text{ gives } 29.8s.$$

The arithmetical sum of 29.8s. and the next following quantity in the table is 30.4s.

$$60' : 30.4s. :: 13^\circ 33' : 6.9s.$$

From	29.8s.
Take	6.9
	22.9s.

4. Given the argument  $5^s. 8^\circ 14' 52''$ , to find the corresponding quantity in Table LII. Ans.  $12' 36''.0$ .

5. Given the argument  $11^s. 11^\circ 23' 10''$ , to find the corresponding quantity in Table LVI. Ans.  $11' 48''.0$ .

6. Given the argument  $0^s. 26^\circ 20'$ , to find the corresponding quantity in Table XII. Ans. —  $41^s.0$ .

Case 2. *When the argument changes in the table by more or less than  $1^\circ$ ; or when it is given in lower denominations than signs.*

Take out of the table the quantity answering to the number in the column of arguments next less than the given argument. Take the difference between this quantity and the next following one, and also the difference of the consecutive values of the argument inserted in the table, and say, difference of arguments : difference of quantities :: excess of the given argument over the value next less in the table : a fourth term. This



fourth term applied to the quantity first taken out, according to the rule given in the preceding case, will give the quantity sought.

Note 3. In some of the tables the columns entitled Diff. are made up of the differences answering to a difference of 10' in the argument. In obtaining quantities from these tables, it will be found more convenient to take for the first and second terms of the proportion, respectively, 10', and the difference furnished by the table, and work the proportion by the arithmetical method. (See note at bottom of page 256).

Exam. 1. Given the argument  $0^s. 24^{\circ} 42' 15''$ , to find the corresponding quantity in Table LI.

$0^s. 24^{\circ} 30'$  gives  $9^{\circ} 47' 14''.3$ .

The difference between  $9^{\circ} 47' 14''.3$  and the next following quantity  $= 3 \times 63''.0 = 189''.0$ . The argument changes by 30'. And the excess of  $0^s. 24^{\circ} 42' 15''$  over  $0^s. 24^{\circ} 30'$  is  $12' 15''$ . Thus,

$$30' : 189''.0 :: 12' 15'' : 77''.2.$$

But the correction may be found more readily by the following proportion,

$$10' : 63''.0 :: 12'.25 : 77''.2.$$

To  $9^{\circ} 47' 14''.3$

Add  $77.2$

---

$9^{\circ} 48' 31.5$

2. Given the argument  $1^{\circ} 12'$ , to find the corresponding quantity in Table VIII.

$1^{\circ} 10'$  gives  $23' 13''$ ,

and

$5' : 27'' :: 2' : 11''$  the correction.

From  $23' 13''$

Take  $11$

---

$23 \quad 2$

3. Given the argument  $6^s. 6^{\circ} 7' 23''$ , to find the corresponding quantity in Table LV. Ans.  $90^{\circ} 20' 53''.5$ .

4. Given the argument  $49^{\circ} 27'$ , to find the corresponding quantity in Table LXIV. Ans.  $11' 19''.8$ .

Case 3. *When the argument is given in the table in hundredth, thousandth, or ten thousandth parts of a circle.*

The required quantity can be found in this case by the same

rule as in the preceding ; but it can be had more expeditiously by observing the following rules. If the argument varies by 10, multiply the difference of the quantities between which the required quantity lies by the excess of the given argument over the next less value in the table, and remove the decimal point one figure to the left ; the result will be the correction to be applied to the quantity taken out of the table. The same rule will apply in taking quantities from tables in which the differences, answering to a change of 10 in the argument, are given, although the argument should actually change by 50 or 100. If the argument changes by 100, multiply as above, and remove the decimal point two figures to the left. When the common difference of the arguments is 5, proceed as if it were 10, and double the result. In like manner, when the common difference is 50, proceed as if it were 100, and double the result.

Exam. 1. Given the argument 973, to find the corresponding quantity in Table XLV, column headed 13.

970 gives  $23''.5$ .

The difference is  $1''.2$ , and the excess 3.

$1''.2$	From	$23''.5$
3	Take	.4
<hr/>		<hr/>
Corr. .36		$23.1$

2. Given the argument 4834, to find the corresponding quantity in Table XLII, column headed 5.

4800 gives  $2' 3''.7$ .

The difference is  $6''.8$ , and the excess 34.

$6''.8$				From	$2' 3''.7$
34				Take	2.3
<hr/>					<hr/>
2.312	-	-	-		$2 1.4$

3. Given the argument 5444, to find the corresponding quantity in Table XLI.

Ans.  $15' 37''.7$ .

4. Given the argument 4225, to find the corresponding quantity in Table XLIII, column headed 8.

Ans.  $0' 47''.2$ .

Case 4. *When the table is one of double entry, or quantities are taken from it by means of two arguments.*

Take out of the table the quantity answering to the values of

the arguments of the table next less than the given values ; and find the respective corrections to be applied to it, due to the excess of the given value of each argument over the next less value in the table, by the general rule given in the preceding case. These corrections are to be added to the quantity taken out, or subtracted from it, according as the quantities increase or decrease with the arguments.

Note 1. If the tenths of seconds be omitted, the corrections above mentioned can be estimated, without the trouble of stating a proportion, or performing multiplications.

Note 2. The rule above given may, in some rare instances, give a result differing a few tenths of a second from the truth. The following rule will furnish more exact results. Find the quantities corresponding, respectively, to the value of the argument at the top next less than its given value, and the other given argument, and to the value next greater and the other given argument. Take the difference of the quantities found, and also the difference of the corresponding arguments at top, and say, difference of arguments : difference of quantities :: excess of given value of the argument at the top over its next less value in the table : a fourth term. This fourth term added to the quantity first found, if it is less than the other, but subtracted from it, if it is greater, will give the required quantity. The error of the first rule may be diminished without any extra calculation, by attending to the differences of the quantities answering to the value of the argument at the side next *greater* than its given value, and the values of the other argument, between which its given value lies.

Exam. 1. Given the argument 64 at the top and 77 at the side, to find the corresponding quantity in Table LXXXI.

50 and 70 give  $47''.7$ .

The difference between  $47''.7$  and the next quantity below it is  $1''.4$ . The excess of 77 over 70 is 7, and the argument at the side changes by 10.

	$1''.4$		
	$7$		
	<hr/>		
		From	$47''.7$
Corr. due excess 7,	$.98$ , or $1''.0$ .	Take	$1.0$
			<hr/>
Quantity corresponding to 50 and 77,			$46.7$

The difference between  $47''.7$  and the adjacent quantity in the next column on the right is  $3''.3$ . The excess of 64 over 50 is 14, and the argument at the top changes by 50.

	$3''.3$		
	$14$		
	<hr style="width: 50%; margin: 0;"/>		
	$.462$		
	$2$		
	<hr style="width: 50%; margin: 0;"/>		
Corr. due excess 14,	$.924$	From	$46''.7$
		Take	$0 \ .9$
			<hr style="width: 50%; margin: 0;"/>
			$45 \ .8$

2. Given the argument 223 at the top and 448 at the side, to find the corresponding quantity in Table XXX.

$220$  and  $440$  give  $16''.0$ .

The difference between  $16''.0$  and the quantity next below it is  $2''.2$ .

	$2''.2$		
	$8$		
	<hr style="width: 50%; margin: 0;"/>		
	$2 \ ) \ 1.76$		
	<hr style="width: 50%; margin: 0;"/>		
Corr. for excess 8,	$.88$ , or $0''.9$ .	From	$16''.0$
		Take	$0 \ .9$
			<hr style="width: 50%; margin: 0;"/>

Quantity corresponding to 220 and 448,  $15 \ .1$

The difference between  $16''.0$  and the adjacent quantity in the next column on the right is  $0''.7$ .

	$0''.7$		
	$3$		
	<hr style="width: 50%; margin: 0;"/>		
Corr. for excess 3,	$.21$	To	$15''.1$
		Add	$\ .2$
			<hr style="width: 50%; margin: 0;"/>
			$15 \ .3$

3. Given the argument 472 at the top and 786 at the side, to find the corresponding quantity in Table XXXI.

Ans.  $9''.7$ .

4. Given the argument 620 at the top and 367 at the side, to find the corresponding quantity in Table LXXXI.

Ans.  $55''.2$ .

5. Given the argument 348 at the top and 932 at the side, to find (by the rule given in Note 2) the corresponding quantity in Table XXXII.

Ans.  $15''.4$ .

## PROBLEM III.

*To convert Degrees, Minutes, and Seconds of the Equator into Time.*

Multiply the quantity by 4, and call the product of the seconds thirds; of the minutes, seconds; and of the degrees, minutes.

Exam 1. Convert  $83^{\circ} 11' 52''$  into time.

$$\begin{array}{r} 83^{\circ} \ 11' \ 52'' \\ \cdot \qquad \qquad \qquad 4 \\ \hline 5^{\text{h}} \ 32^{\text{m}} \ 47^{\text{s}} \ 28''' \end{array}$$

2. Convert  $34^{\circ} 57' 46''$  into time.

Ans. 2h. 19m. 51sec. 4'''.

## PROBLEM IV.

*To convert Time into Degrees, Minutes, and Seconds.*

Reduce the hours and minutes, to minutes; divide by 4, and call the quotient of the minutes, degrees; of the seconds, minutes; and multiply the remainder by 15, for the seconds.

Exam. 1. Convert 7h. 9m. 34sec. into degrees, &c.

$$\begin{array}{r} 7^{\text{h}} \ 9^{\text{m}} \ 34^{\text{s}} \\ 60 \\ \hline 4 \ ) \ 429 \ 34 \\ \hline 107^{\circ} \ 24' \ 30'' \end{array}$$

2. Convert 11h. 24m. 45s. into degrees, &c.

Ans.  $171^{\circ} 11' 15''$ .

## PROBLEM V.

*The Longitudes of two Places, and the Time at one of them being given, to find the corresponding time at the other.*

When the given time is in the morning, change it to astronomical time, by adding 12 hours, and diminishing the number of the



day by a unit. When the given time is in the evening, it is already in astronomical time.

Find the difference of longitude of the two places, by taking the numerical difference of their longitudes, when these are of the same name; that is, both east or both west; and the sum, when they are of different names; that is, one west and the other east. When one of the places is Greenwich, the longitude of the other is the difference of longitude.

Then, if the place at which the time is required, is to the *east* of the other place, *add* the difference of longitude, in time, to the given time; but, if it is to the *west*, *subtract* the difference of longitude, from the given time. The sum or remainder will be the required time.

Note. The longitudes of the places mentioned in the following examples, are given in Table I.

Exam. 1. When it is October 25th, 3h. 13m. 22sec. A. M., at Greenwich, what is the time, as reckoned at New York?

Time at Greenwich, October,	24 <sup>d</sup> .	15 <sup>h</sup> .	13 <sup>m</sup> .	22 <sup>s</sup> .
Diff. of Long.	-	-	-	4 56 4

---

Time at New York	-	-	24	10	17	18 P. M.
------------------	---	---	----	----	----	----------

2. When it is June 9h. 5h. 25m. 10sec. P. M. at Washington, what is the corresponding time at Greenwich?

Time at Washington, June,	9 <sup>d</sup> .	5 <sup>h</sup> .	25 <sup>m</sup> .	10 <sup>s</sup> .
Diff. of Long.	-	-	-	5 8 7

---

Time at Greenwich	-	-	9	10	33	17 P. M.
-------------------	---	---	---	----	----	----------

3. When it is January 15th, 2h. 44m. 23sec. P. M. at Paris, what is the time at Philadelphia?

Longitude of Paris,	-	-	-	0 <sup>h</sup> .	9 <sup>n</sup> .	21 <sup>s</sup> .	6 E
Do. of Philadelphia,	-	-	-	5	0	44	W.

---

	5	10	5.6
--	---	----	-----

Time at Paris, January,	-	15 <sup>d</sup> .	2 <sup>h</sup> .	44 <sup>m</sup> .	23 <sup>s</sup> .
Diff. of Long,	-	-	-	5	10 6

---

Time at Philadelphia,	-	-	14	21	34	17
-----------------------	---	---	----	----	----	----

Or January 15th, 9h. 34m. 17sec. A. M.

4. When it is March 31st, 8h. 4m. 21sec. P. M. at New Haven, what is the corresponding time at Berlin?

Ans. April 1st, 1h. 49m. 48sec. A. M.

5. When it is August 10th, 10h. 32m. 14sec. A. M. at Boston, what is the time at New Orleans?

Ans. Aug. 10th, 9h. 16m. 3sec. A. M.

6. When it is noon of the 23d of December at Greenwich, what is the time at New York?

Ans. Dec. 23d, 7h. 3m. 55sec. A. M.

### PROBLEM VI.

*The Apparent Time being given, to find the corresponding Mean Time; or the Mean Time being given, to find the Apparent.*

When the given time is not for the meridian of Greenwich, reduce it to that meridian by the last problem. Then find by the tables the sun's mean longitude corresponding to this time. Thus, from Table XVIII take out the longitude answering to the given year, and from Tables XIX, XX, and XXI take out the motions in longitude for the given month, days, hours and minutes, neglecting the seconds. The sum of the quantities taken from the tables, rejecting 12 signs, when it exceeds that quantity, will be the sun's mean longitude for the given time.

With the sun's mean longitude, thus found, take the Equation of Time from Table XII. Then, when Apparent Time is given to find the Mean, apply the equation with the sign it has in the table; but when Mean Time is given to find the Apparent, apply it with the contrary sign; the result will be the Mean or Apparent Time required.

This rule will be sufficiently exact for ordinary purposes, for several years before and after the year 1840. When the given date is a number of years distant from this epoch, take also with the sun's mean longitude the Secular Variation of the Equation of Time from Table XIII, and find by simple proportion the variation in the interval between the given year and 1840. The result, applied to the equation of time taken from Table XII, according to

its sign, if the given time is subsequent to the year 1840, but with the opposite sign, if it is prior to 1840, will give the equation of time at the given date, which apply to the given time as above directed.

Note 1. When the exact mean or apparent time to within a small fraction of a second is demanded, take the numbers in the columns entitled I, II, III, IV, V, N, in Tables XVII, XIX, XX, XXI, answering respectively to the year, months, days, hours, and minutes of the given time. With the respective sums of the numbers taken from each column, as arguments, enter Table XIV, and take out the corresponding quantities. These quantities added to the equation of time as given by Tables XII and XIII, and the constant 3.0s. subtracted, will give the true Equation of Time, if the given time is Mean Time. When Apparent Time is given, it will be farther necessary to correct the equation of time as given by the tables, by stating the proportion, 24 hours : change of equation for  $1^{\circ}$  of longitude : : equation of time : correction.

Note 2. The Equation of Time is given in the Nautical Almanac for each day of the year, at apparent, and also at mean noon, on the meridian of Greenwich, and can easily be found for any intermediate time by proportion. Directions for applying it to the given time are placed at the head of the column. The Equation is given on the first and second pages of each month.

Exam. 1. On the 16th of July, 1840, when it is 9h. 35m. 22s. P. M. mean time at New York, what is the apparent time at the same place?

Time at New York, July 1840, - 16<sup>d.</sup> 9<sup>h.</sup> 35<sup>m.</sup> 22<sup>s.</sup>

Diff. of Long. - - - - 4 56 4

Time at Greenwich, July 1840, 16 14 31 26

M. Long.

1840 - - - - - 9<sup>s.</sup> 10<sup>o</sup> 12' 49"

July - - - - - 5 29 23 16

16d. - - - - - 14 47 5

14h. - - - - - 34 30

31m. - - - - - 1 16

M. Long. - - - - 3 24 58 56

The equation of time in Table XII, corresponding to  $3^{\text{h}} 24^{\circ} 58'$   $56''$ , is  $+ 5^{\text{m}} 44^{\text{s}}$ .

Mean Time at New York, July 1840,  $16^{\text{h}} 9^{\text{m}} 35^{\text{s}} 22^{\text{s}}$ .

Equation of time, sign changed,  $- 5 44$

Apparent Time,  $16 9 29 38$  P. M.

2. On the 9th of May, 1842, when it is  $4^{\text{h}} 15^{\text{m}} 21^{\text{s}}$  A. M. apparent time at New York, what is the mean time at the same place, and also at Greenwich?

Time at New York, May 1842,  $8^{\text{d}} 16^{\text{h}} 15^{\text{m}} 21^{\text{s}}$ .

Diff. of Long.  $- - - 4 56 4$

Time at Greenwich,  $- - 8 21 11 25$

M. Long.

1842  $- - 9^{\text{h}} 10^{\circ} 43^{\text{m}} 18^{\text{s}}$

May  $- - 3 28 16 40$

8 d.  $- - 6 53 58$

21 h.  $- - 51 45$

11 m.  $- - 27$

M. Long.  $- 1 16 46 8$ . Equa. of time,  $- 3^{\text{m}} 45^{\text{s}}$ .

Apparent Time at Greenwich, May, 1842,  $8^{\text{d}} 21^{\text{h}} 11^{\text{m}} 25^{\text{s}}$ .

Equation of Time,  $- - - - - 3 45$

Mean Time at Greenwich,  $- - - 8 21 7 40$

Diff. of Long.  $- - - - - 4 56 4$

Mean Time at New York,  $- - - 8 16 11 36$

Or, May 9th,  $4^{\text{h}} 11^{\text{m}} 36^{\text{s}}$  A. M.

3. On the 3d of February, 1855, when it is  $2^{\text{h}} 43^{\text{m}} 36^{\text{s}}$  apparent time at Greenwich, what is the exact mean time at the same place?

Appar. Time at Greenwich, Feb., 1855,  $3^{\text{d}} 2^{\text{h}} 43^{\text{m}} 36^{\text{s}}$ .

	M. Long.				I.	II.	III.	IV.	V.	N.
1855 . .	9 <sup>s</sup>	10 <sup>o</sup>	34'	30"	433	279	806	889	866	863
Feb. . .	1	0	33	18	47	85	138	45	7	5
3 d. . .		1	58	17	68	5	9	3	0	0
2 h. . .			4	56	3					
43 m. . .			1	46						
	10	13	12	47	551	369	953	937	873	868

Appar. Time at Greenwich, Feb. 1855,	3 <sup>h</sup> .	2 <sup>h</sup> .	43 <sup>m</sup>	36 <sup>s</sup> .
Equation of time by Table XII, -	-	-	+ 14	8.6
100yrs. : 13s. (Sec. Var. Table XIII)				
:: 15yrs. : 1.9s. -	-	-	-	- 1.9
Approx. Mean Time at Greenwich, -	3	2	57	42.7
24h. : 6s. (change of equa. for 1° of				
long.) : 14m. : 0.1s. -	-	-	-	+ 0.1
II. III. -	-	-	-	0.8
II. IV. -	-	-	-	1.0
II. V. -	-	-	-	0.4
I. -	-	-	-	0.3
N. -	-	-	-	0.1
Constant. -	-	-	-	- 3.0

Mean Time at Greenwich	-	3	2	57	42.4
------------------------	---	---	---	----	------

4. On the 18th of November, 1841, when it is 2h. 12m. 26sec. A. M. mean time at Greenwich, what is the apparent time at Philadelphia?      Ans. Nov. 17th, 9h. 26m. 24s. P. M.

5. On the 2d of February, 1839, when it is 6h. 32m. 35sec. P. M. apparent time at New Haven, what is the mean time at the same place?      Ans. 6h. 46m. 38s. P. M.

6. On the 23d of September, 1850, when it is 9h. 10m. 12sec. mean time at Boston, what is the *exact* apparent time at the same place?      Ans. 9h. 8m. 1.0s.

## PROBLEM VII.

*To correct the Observed Altitude of a Heavenly Body for Refraction.*

With the given altitude take the corresponding refraction from Table VIII. Subtract the refraction from the given altitude, and the result will be the true altitude of the body at the given station.

This rule will give exact results if the barometer stands at 30 inches, and Fahrenheit's thermometer at 50°, and results sufficiently exact for ordinary purposes in any state of the atmosphere. When there is occasion for greater precision, take from Table IX.





### PROBLEM VIII.

*The Apparent Altitude of a Heavenly Body being given, to find its True Altitude.*

Correct the observed altitude for refraction by the foregoing problem. Then,

1. If the sun is the body whose altitude is taken, find its parallax in altitude by Table X, and add it to the observed altitude corrected for refraction. The result will be the true altitude, sought.

2. If it is the altitude of the moon that is taken, and the horizontal parallax at the time of the observation is known, find the parallax in altitude by the following formula :

$$\log. \sin (\text{par. in alt.}) = \log. \sin (\text{hor. par.}) + \log. \cos. (\text{app. alt.}) - 10;$$

and add it, as before, to the apparent altitude corrected for refraction.

3. If one of the planets is the body observed, the following formula will serve for the determination of the parallax in altitude when the horizontal parallax is known :

$$\log. (\text{par. in alt.}) = \log. (\text{hor. par.}) + \log. (\cos \text{ appar. alt.}) - 10.$$

Note 1. The equatorial horizontal parallax of the moon at any given time may be obtained from the tables appended to the work. (See Problem XIV). But it can be had much more readily from the Nautical Almanac. The equatorial horizontal parallax being known, the horizontal parallax at any given latitude may be obtained by subtracting the Reduction of Parallax, to be found in Table LXIV. The horizontal parallax of any planet, the altitude of which is measured, may also be derived from the Nautical Almanac.

Note 2. The fixed stars have no sensible parallax, and thus the observed altitude of a star, corrected for refraction, will be its true altitude at the centre of the earth as well as at the station of the observer.

Note 3. If the true altitude of a heavenly body is given, and it is required to find the apparent, the rules for finding the parallax in altitude and the refraction are the same as when the

apparent altitude is given; the true altitude being used in place of the apparent. But these corrections are to be applied with the opposite signs from those used in the determination of the true altitude from the apparent; that is, the parallax is to be subtracted, and the refraction added. It will also be more accurate to make use of equa. (12), p. 44, in the case of the moon.

Exam. 1. The observed altitude of the sun on the 1st of May, 1837, being  $26^{\circ} 40' 20''$ , what is its true altitude?

Obs. alt.	-	-	-	-	-	$26^{\circ} 40' 20''$
Refraction	-	-	-	-	-	$-1\ 56$

True alt. at the station,	-	-	-	$26\ 38\ 24$
Parallax in alt. (Table X)	-	-	-	$+8$

True altitude	-	-	-	-	$26\ 38\ 32$
---------------	---	---	---	---	--------------

2. Let the apparent altitude of the moon at New York on the 17th of March, 1837, Sh. P. M., be  $66^{\circ} 10' 44''$ ; the barometer 30.4 in. and the thermometer  $62^{\circ}$ ; required the true altitude.

Appar. alt.	-	-	$66^{\circ} 10' 44''$
-------------	---	---	-----------------------

Mean refrac.	-	$0\ 25.7$
--------------	---	-----------

Corr. for $+0.4$ in. bar.	$+0.3$
---------------------------	--------

Corr. for $+12^{\circ}$ ther.	$-0.6$
-------------------------------	--------

True refrac.	-	$0\ 25.4$
--------------	---	-----------

True alt. at N. York,	$66\ 10\ 18.6$
-----------------------	----------------

logarithms.  
cos. 9.60637

Equa. par. by N. Almanac,  $54' 13''$

Reduc. for lat.  $40^{\circ}$ ,  $4$

Hor. par. at New York,	$54\ 9$	-	-	sin. 8.19731
------------------------	---------	---	---	--------------

Par. in alt.	-	-	$21\ 52$	sin. 7.80368
--------------	---	---	----------	--------------

True altitude	-	$66\ 32\ 11$
---------------	---	--------------

3. On the 18th of February, 1837, the true meridian altitude of the planet Jupiter at Greenwich was  $56^{\circ} 54' 57''$ , what was its apparent altitude at the time of the meridian passage, the horizontal parallax being taken at  $1''.9$  as given by the Nautical Almanac?

True alt.	-	-	56° 54' 57"	-	cos.	9.7371
Hor. par.	1".9	-	-	-	log.	0.2787
Par. in alt.	-	-	-	— 1.0	-	log. 0.0158
Refraction	-	-	-	+ 21.0		
Appar. alt.	-	-	56 54 37			

4. What will be the true altitude of the sun on the 22d of September, 1840, at the time its apparent altitude is  $39^{\circ} 17' 50''$ ?

Ans.  $39^{\circ} 16' 46''$ .

5. Given  $29^{\circ} 33' 30''$  the apparent altitude of the moon at Philadelphia on the 15th of June, 1837, at 9h. 30m. P. M., and  $58' 33''$  the equatorial parallax of the moon at the same time, to find the true altitude.

Ans.  $30^{\circ} 22' 41''$ .

6. Given  $15^{\circ} 24' 23''$  the true altitude of Venus, and  $8''$  its horizontal parallax, to find the apparent altitude.

Ans.  $15^{\circ} 27' 41''$ .

## PROBLEM IX.

*To find the Sun's Longitude, Semi-diameter, and Hourly Motion, for a given time, from the Tables.*

*For the Longitude.*

When the given time is not for the meridian of Greenwich, reduce it to that meridian by Problem V; and when it is apparent time, convert it into mean time by the last problem.

With the mean time at Greenwich, take from Tables XVIII, XIX, XX, and XXI, the quantities corresponding to the year, month, day, hour, minute, and second (omitting those in the last two columns), and place them in separate columns headed as in Table XVIII, and take their sums.\* The sum in the column entitled *M. Long.* will be the tabular mean longitude of

---

\* In adding quantities that are expressed in signs, degrees, &c. reject 12 or 24 signs whenever the sum exceeds either of these quantities. In adding arguments expressed in 100 or 1000, &c. parts of the circle, when they consist of two figures, reject the hundreds from the sum; when of three figures, the thousands; and when of four figures, the ten thousands.

the sun ; the sum in the column entitled *Long. Perigee* will be the tabular longitude of the sun's perigee ; and the sums in the columns headed I, II, III, IV, V, N, will be the arguments for the small equations of the sun's longitude, and for the equation of the equinoxes, which forms one of them.

Subtract the longitude of the perigee from the sun's mean longitude, adding 12 signs when necessary to render the subtraction possible ; the remainder will be the sun's mean anomaly. With the mean anomaly take the equation of the sun's centre from Table XXV, and correct it by estimation, for the proportional part of the secular variation in the interval between the given year and 1840 ; also, with the arguments I, II, III, IV, V, take the corresponding equations from Tables XXVIII, XXX, XXXI and XXXII. The equation of the centre and the four other equations, added to the mean longitude, will give the sun's True Longitude, reckoned from the Mean Equinox.

With the argument N take the equation of the equinoxes or Lunar Nutation in longitude from Table XXVII. Also take the Solar Nutation in longitude, answering to the given date, from the same table. Apply these equations according to their signs to the true longitude from the mean equinox, already found, and add the constant 3", the result will be the True Longitude from the Apparent Equinox.

*For the Semi-diameter and Hourly Motion.*

With the sun's mean anomaly, take the Hourly Motion and Semi-diameter from Tables XXIII and XXIV.

Note 1. If the tenths of seconds be omitted in taking the equations from the tables of double entry, the error cannot exceed 2" ; in case the precaution is taken to add a unit, whenever the tenths exceed .5.

Note 2. The longitude of the sun, obtained by the foregoing rule, may differ about 3" from the same as derived from the most accurate solar tables now in use. When there is occasion for greater precision, take from Table XVIII, XIX, and XX, the quantities in the columns entitled VI and VII, along with those in the other columns. With the sums in these columns, and those in the columns I, II, as arguments, take the corresponding equations from Tables XXIX and XXXIII. Also with the sun's mean ano-



maly take the equation for the variable part of the aberration from Table XXXIV. Add these three equations along with the others to the mean longitude, and omit the addition of the constant 3". The result will be exact to within a fraction of a second.

Exam. 1. Required the sun's longitude, hourly motion, and semi-diameter, on the 25th October, 1837, at 11h. 27m. 38s. A. M. mean time at New York.

Mean time at N. York, Oct. 1837, 24<sup>d</sup>. 23<sup>h</sup>. 27<sup>m</sup>. 38<sup>s</sup>.

Diff. of Long. - - - 4 56 4

Mean time at Greenwich - 25 4 23 42

	M. Long.	Long. Perigee.	I.	II.	III.	IV.	V.	N.
	s o ' "	s o ' "						
1837 . . .	9 10 55 47.2	9 10 8 5	816	280	549	321	348	895
October . .	8 29 4 54.1	46	250	748	215	397	63	40
25 d. . . .	23 39 19.9	4	810	66	107	35	5	4
4 h. . . .	9 51.4		6	0	1			
23 m. . . .	56.7							
42 s. . . .	1.7	9 10 8 55	882	94	872	753	416	939
		7 3 50 51						
Eq. Sun's Cent.	7 3 50 51.0							
I. . . .	11 28 12 43.5	9 23 41 56	Mean Anomaly.					
II. III. . .	2.5							
II. IV. . .	9.0	Sun's Hourly Motion, . . . . 2' 29".7						
II. V. . . .	7.7							
Const. . .	19.3	Sun's Semi-diameter, . . . . 16' 17".2						
	3.0							
	7 2 4 16.0							
Lunar Nutation	— 6.1							
Solar Nutation	— 1.2							
	7 2 4 8.7							

2. Required the sun's longitude, hourly motion, and semi-diameter, on the 15th of July, 1837, at 8h. 20m. 40s. P. M. mean time at Greenwich.

	M. Long.	Long. Peri.	I.	II.	III.	IV.	V.	N.	VI.	VII.
	° ' " "	° ' " "								
1837 . . .	9 10 '55 47.2	9 10 8 5	816	280	549	321	348	895	787	600
July . . .	5 28 24 7.8	31	129	496	806	263	41	27	569	17
15 d. . . .	13 47 56.6	2	473	38	62	20	3	2	508	2
8 h. . . .	19 42.8		11	1	1				11	
20 m. . . .	49.3									
40 s. . . .	1.6	9 10 8 38	429	815	418	604	392	924	876	619
		3 23 28 25								
	3 23 28 25.3									
Eq. Sun's Cent.	11 29 33 10.1	6 13 19 47	Mean Anomaly.							
I. . . .	10.7		Sun's Hourly Motion, . . . . 2' 23".1							
II. III. . .	6.6		Sun's Semi-diameter, . . . . 15' 45".4							
II. IV. . .	5.0									
II. V. . . .	7.7									
I. VI. . . .	1.8									
II. VII. . .	0.2									
Aber. . . .	0.6									
	3 23 2 8.0									
Lunar Nutation	— 7.8									
Solar Nutation	+ 0.8									
Sun's true long.	3 23 2 1.0									

3. Required the sun's longitude, hourly motion, and semi-diameter, on the 10th of June, 1838, at 9h. 45m. 26s. A. M. mean time at Philadelphia, (omitting the three smallest equations of longitude).

Ans. Sun's longitude,  $2^{\text{s}} 19^{\circ} 11' 57''$ ; hourly motion,  $2' 23''.3$ ; semi-diameter,  $15' 46''.1$ .

4. Required the sun's longitude, hourly motion, and semi-diameter, on the 1st of February, 1837, at 12h. 30m. 15s. P. M. mean time at Greenwich.

Ans. Sun's longitude,  $10^{\text{s}} 13^{\circ} 1' 44''.6$ ; hourly motion,  $2' 32''.1$ ; semi-diameter,  $16' 14''.7$ .

### PROBLEM X.

*To find the Apparent Obliquity of the Ecliptic, for a given time, from the Tables.*

Take the mean obliquity for the given year from Table XXII. Then with the argument N, found as in the foregoing problem, and the given date, take from Table XXVII the lunar and solar nutations of obliquity; apply these according to their signs to

the mean obliquity; and the result will be the apparent obliquity.

Exam. 1. Required the apparent obliquity of the ecliptic on the 15th of March, 1839.

	N.				
1839, -	3				
March,	9				
15d. -	2				
	—	M. Obliquity,	23° 27' 36".9		
	14	- - - -	- - + 9 .1		
Solar Nutation for March 15th,		- -	+ 0 .5		
			—		
Apparent Obliquity, - - -			23 27 46 .5		

2. Required the apparent obliquity of the ecliptic on the 12th of July, 1845. Ans. 23° 27' 28".0.

## PROBLEM XI.

*Given the Sun's Longitude and the Obliquity of the Ecliptic, to find his Right Ascension and Declination.\**

Let  $\omega$  = obliquity of the ecliptic; L = sun's longitude; R = sun's right ascension; and D = sun's declination; then to find R and D, we have,

$$\log. \tan g R = \log. \tan g L + \log. \cos \omega - 10,$$

$$\log. \sin D = \log. \sin L + \log. \sin \omega - 10.$$

The right ascension must always be taken in the same quadrant as the longitude. The declination must be taken less than 90°; and it will be north or south according as its trigonometrical sine comes out positive or negative.

Note. The sun's right ascension and declination are given in the Nautical Almanac for each day in the year at noon on the meridian of Greenwich, and may be found at any intermediate time by a proportion.

Exam. 1. Given the sun's longitude 205° 23' 50", and the ob-

---

\* The obliquity of the ecliptic at any given time for which the sun's longitude is known, is found by the foregoing Problem.

liquity of the ecliptic  $23^{\circ} 27' 36''$ , to find his right ascension and declination.

$$L = 205^{\circ} 23' 50'' \quad - \quad - \quad - \quad \tan. \quad 9.67649$$

$$\omega = 23 \quad 27 \quad 36 \quad - \quad - \quad - \quad \cos. \quad 9.96253$$

$$R = 203 \quad 32 \quad 5 \quad - \quad - \quad - \quad \tan. \quad 9.63902$$

$$L = 205 \quad 23 \quad 50 \quad - \quad - \quad - \quad \sin. \quad 9.63235 \text{ ---}$$

$$\omega = 23 \quad 27 \quad 36 \quad - \quad - \quad - \quad \sin. \quad 9.60000$$

$$D = 9 \quad 49 \quad 52 \text{ S.} \quad - \quad - \quad - \quad \sin. \quad 9.23235 \text{ ---}$$

2. The obliquity of the ecliptic being  $23^{\circ} 27' 30''$ , required the sun's right ascension and declination when his longitude is  $44^{\circ} 18' 25''$ .

Ans. Right ascension  $41^{\circ} 50' 30''$ , and declination  $16^{\circ} 8' 40''$  N.

## PROBLEM XII.

*Given the Sun's Right Ascension, and the Obliquity of the Ecliptic, to find his Longitude and Declination.*

Using the same notation as in the last problem, we have, to find the longitude and declination,

$$\log. \tan g L = \log. \tan g R + \text{ar. co. log. } \cos \omega,$$

$$\log. \tan g D = \log. \sin R + \log. \tan g \omega - 10.$$

Exam. 1. What is the longitude and declination of the sun, when his right ascension is  $142^{\circ} 11' 34''$ , and the obliquity of the ecliptic  $23^{\circ} 27' 40''$ ?

$$R = 142^{\circ} 11' 34'' \quad - \quad - \quad - \quad \tan. \quad 9.88979 \text{ ---}$$

$$\omega = 23 \quad 27 \quad 40 \quad - \quad - \quad \text{Ar. Co. cos.} \quad 0.03747$$

$$L = 139 \quad 46 \quad 30 \quad - \quad - \quad - \quad \tan. \quad 9.92726 \text{ ---}$$

$$R = 142 \quad 11 \quad 34 \quad - \quad - \quad - \quad \sin. \quad 9.78746$$

$$\omega = 23 \quad 27 \quad 40 \quad - \quad - \quad - \quad \tan. \quad 9.63750$$

$$D = 14 \quad 53 \quad 55 \text{ N} \quad - \quad - \quad - \quad \tan. \quad 9.42496$$

2. Given the sun's right ascension  $310^{\circ} 25' 11''$ , and the obliquity of the ecliptic  $23^{\circ} 27' 35''$ , to find the longitude and declination.

Ans. Longitude  $307^{\circ} 59' 57''$ , and declination  $18^{\circ} 17' 0''$  S.

### PROBLEM XIII.

*The Sun's Longitude and the Obliquity of the Ecliptic being given, to find the Angle of Position.*

Let  $p$  = angle of position ;  $\omega$  = obliquity of the ecliptic ; and  $L$  = sun's longitude. Then,

$$\log. \tan p = \log. \cos L + \log. \tan \omega - 10.$$

The angle of position is always less than  $90^\circ$ . The northern part of the circle of latitude will be to the *west* or *east* of the northern part of the circle of declination, according as the sign of the tangent of the angle of position is *positive* or *negative*.

Exam. 1. Given the sun's longitude  $24^\circ 15' 20''$ , and the obliquity of the ecliptic  $23^\circ 27' 32''$ , required the angle of position.

$$L = 24^\circ 15' 20'' \quad - \quad - \quad - \quad - \quad \cos. \quad 9.95986$$

$$\omega = 23 \quad 27 \quad 32 \quad - \quad - \quad - \quad - \quad \tan. \quad 9.63745$$

$$p = 21 \quad 35 \quad 10 \quad - \quad - \quad - \quad - \quad \tan. \quad 9.59731$$

The northern part of the circle of latitude lies to the west of the circle of declination.

2. When the sun's longitude is  $120^\circ 18' 55''$ , and the obliquity of the ecliptic  $23^\circ 27' 30''$ , what is the angle of position ?

Ans.  $12^\circ 21' 17''$  ; and the northern part of the circle of latitude lies to the east of the circle of declination.

### PROBLEM XIV.

*To find from the Tables, the Moon's Longitude, Latitude, Equatorial Parallax, Semi-diameter, and Hourly Motion in Longitude and Latitude, for a given time.*

When the given time is not for the meridian of Greenwich, reduce it to that meridian, and when it is apparent time convert it into mean time.

Take from Table XXXV, and the following tables, the arguments numbered 1, 2, 3, &c., to 20, for the given year, and their variations for the given month, days, &c., and find the sums of the numbers for the different arguments respectively ; rejecting



the hundred thousands and also the units in the first, the ten thousands in the next eight, and the thousands in the others.

The resulting quantities will be the arguments for the first twenty equations of longitude.

With the same time, take from the same tables the remaining arguments with their variations, entitled Evection, Anomaly, Variation, Longitude, Supplement of the Node, II, V, VI, VII, VIII, IX, and X, and add the quantities in the column for the Supplement of the Node.

*For the Longitude.*

With the first twenty arguments of longitude, take from Tables XLI to XLVI, inclusive, the corresponding equations; and with the Supplement of the Node for another argument, take the corresponding equation from Table XLIX. Place these twenty-one equations in a single column, headed *Eqs. of Long.*; and write beneath them the constant 55". Find the sum of the whole, and place it in the column of Evection. Then the sum of the quantities in this column will be the corrected argument of Evection.

With the corrected argument of Evection, take the Evection from Table L, and add it to the sum in the column of *Eqs. of Long.* Place this in the column of Anomaly. Then the sum of the quantities in this column will be the corrected Anomaly.

With the corrected Anomaly, take the Equation of the Centre from Table LI, and add it to the last sum in the column of *Eqs. of Long.* Place the resulting sum in the column of Variation. Then the sum of the quantities in this column will be the corrected argument of Variation.

With the corrected argument of Variation, take the variation from Table LII, and add it to the last sum in the column of *Eqs. of Long.*; the result will be the sum of the principal equations of the Orbit Longitude, amounting in all to twenty-four, and the constants subtracted for the other equations. Place this sum in the column of Longitude. Then, the sum of the quantities in this column will be the Orbit Longitude of the Moon, reckoned from the mean equinox.

Add the orbit longitude to the supplement of the node, and the resulting sum will be the argument of Reduction.

With the argument of Reduction, take the Reduction from Table LIII, and add it to the Orbit Longitude. The sum will be the Longitude as reckoned from the mean equinox. With the Supplement of the Node, take the Nutation in Longitude from Table XXVII, and apply it, according to its sign, to the longitude from the mean equinox. The result will be the Moon's True Longitude from the Apparent Equinox.

*For the Latitude.*

The argument of the Reduction is also the 1st argument of Latitude. Place the sum of the first twenty-four equations of Longitude, taken to the nearest minute, in the column of Arg. II. Find the sum of the quantities in this column, and it will be the Arg. II of Latitude, corrected. The Moon's true Longitude is the 3d argument of Latitude. The 20th argument of Longitude is the 4th argument of Latitude. Take from Table LVIII the thousandth parts of the circle, answering to the degrees and minutes in the sum of the first twenty-four equations of longitude ; and place it in the columns V, VI, VII, VIII, and IX ; but not in the column X. Then the sums of the quantities in columns V, VI, VII, VIII, IX and X, rejecting the thousands, will be the 5th, 6th, 7th, 8th, 9th, and 10th arguments of Latitude.

With the Arg. I of Latitude, take the moon's distance from the North Pole of the Ecliptic, from Table LV ; and with the remaining nine arguments of latitude, take the corresponding equations from Tables LVI, LVII and LIX. The sum of these quantities increased by  $8''$ , will be the Moon's true distance from the North Pole of the Ecliptic. The difference between this distance and  $90^\circ$  will be the Moon's true latitude ; which will be *North* or *South*, according as the distance is less or greater than  $90^\circ$ .

*For the Equatorial Parallax.*

With the corrected arguments, Evection, Anomaly, and Variation, take out the corresponding quantities from Tables LXI, LXII, and LXIII. Their sum increased by  $7''$ , will be the Equatorial Parallax.

*For the Semi-diameter.*

With the Equatorial Parallax as an argument, take out the moon's semi-diameter from Table LXV.

*For the Hourly Motion in Longitude.*

With the arguments 2, 3, 4, 5, and 6 of Longitude, rejecting the two right-hand figures in each, take the corresponding equations of the hourly motion in longitude from Table LXVII. Find the sum of these equations, and the constant  $3''$ , and with this sum at the top, and the corrected argument of the Evection at the side, take the corresponding equation from Table LXIX; also with the corrected argument of the Evection, take the corresponding equation from Table LXVIII.

Add these equations to the sum just found, and with the resulting sum at the top, and the correct anomaly at the side, take the corresponding equation from Table LXX; also with the corrected anomaly, take the corresponding equation from Table LXXI.

Add these two equations to the sum last found, and with the resulting sum at the top, and the corrected argument of the Variation at the side, take the corresponding equation from Table LXXII. With the corrected argument of the Variation, take the corresponding equation from Table LXXIII.

Add these two equations to the sum last found, and with the resulting sum at the top, and the argument of the Reduction at the side, take the corresponding equation from Table LXXIV. Also, with the argument of the Reduction take the corresponding equation from Table LXXV. These two equations, added to the last sum, will give the sum of the principal equations of the hourly motion in longitude, and the constants subtracted for the others. To this add the constant  $27' 24''.0$ , and the result will be the Moon's Hourly Motion in Longitude.

*For the Hourly Motion in Latitude.*

With the Argument I of Latitude, take the corresponding equation from Table LXXIX. With this equation, and the sum of all the equations of the hourly motion in longitude, except the last two, take the corresponding equation from Table LXXXI. With the Argument II of Latitude, take the

corresponding equation from Table LXXXII. And with this equation at the top, and the sum of all the equations of the hourly motion in longitude, except the last two, take the equation from Table LXXXIII. Find the sum of these four equations and the constant  $1''$ . To the resulting sum apply the constant  $-2' 37''.2$ . The difference will be the Moon's true Hourly Motion in Latitude. The moon will be tending *North* or *South*, according as the sign is *positive* or *negative*.

Note. The errors of the results obtained by the foregoing rules, occasioned by the neglect of the smaller equations, cannot exceed for the longitude  $15''$ , for the latitude  $8''$ , for the parallax  $7''$ , for the hourly motion in longitude  $5''$ , and for the hourly motion in latitude  $3''$ ; and they will generally be very much less. When greater accuracy is required, take from Tables XXXV to XXXIX the arguments from 21 to 31, along with those from 1 to 20, and their variations. The sums of the numbers for the different arguments, respectively, will be the arguments of eleven small additional equations of longitude. Also take from the same tables the arguments entitled XI and XII, along with those in the preceding columns. Retain the right-hand figure of the sum in column 1 of arguments, and conceive a cypher to be annexed to each number in the columns of arguments of Table XLI. The numbers in the columns entitled *Diff. for 10*, will then be the differences for a variation of 100 in the argument.

*For the Longitude.* With the arguments 21 to 31, take the corresponding equations from Tables XLVII and XLVIII, and place them in the same column with the equations taken out with the arguments 1, 2, &c. to 20. Take also equation 32 from Table XLIX, as before. Find the sum, of the whole, (omitting the constant  $55''$ ) and then continue on as above. The longitude from the mean equinox being found, take the lunar nutation in longitude from Table LIV, and the solar nutation answering to the given date from Table XXVII. Apply them both, according to their sign, to the longitude from the mean equinox, and the result will be the more exact longitude from the apparent equinox, required.

*For the Latitude.* With the arguments XI and XII, take the corresponding equations from Table LIX. Add these with the other equations, and omit the constant  $8''$ . The difference between the sum and  $90^\circ$  will be the more exact latitude.

*For the Equatorial Parallax.* With the arguments 1, 2, 4, 5, 6, 8, 9, 12, 13, take the corresponding equations from Table LX. Find the sum of these and the other equations, omitting the constant  $7''$ , and it will be the more exact value of the Parallax.

*For the Hourly Motion in Longitude.* With the arguments 1, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, and 18, of longitude, along with the arguments 2, 3, 4, 5, and 6, heretofore used, take the corresponding equations from Table LXVII. Find the sum of the whole, omitting the constant  $3''$ , and proceed as in the rule already given.

To obtain the motion in longitude for the hour which precedes or follows the given time, with the arguments of Tables LXX, LXXII, and LXXIV, take the equation from Tables LXXVI and LXXVII. Also, with the arguments of Evection, Anomaly, Variation, and Reduction, take the equations from Tables LXXVIII. Find the sum of all these equations. Then, for the hour which follows the given time, add this sum to the hourly motion at the given time already found, and subtract  $2''.0$ ; for the hour which precedes, subtract it from the same quantity, and add  $2''.0$ .

It will expedite the calculation to take the equations of the second order from the tables, at the same time with those of the first order which have the same arguments.

*For the Hourly Motion in Latitude.* The moon's hourly motion in latitude may be had more exactly by taking with the arguments of Latitude V, VI, &c. to XII, the corresponding equations from Table LXXX, and finding the sum of these and the other equations of the hourly motion in latitude.

To obtain the moon's motion in latitude for the hour which precedes or follows the given time, with the Argument I of Latitude take the equation from Table LXXXIV, and with this equation and the sum of all the equations of the hourly motion in longitude except the two last, take the equation from Table LXXXV.



Find the sum of these two equations. Then, for the hour which follows the given time, add this sum to the Hourly Motion in Latitude already found, and subtract  $1''.3$ ; and for the hour which precedes, subtract it from the same quantity, and add  $1''.3$ .

It will also be more exact to enter Table LXXXI with the sum of all the other equations, diminished by  $6''$ , instead of the last equation, for the argument at the side. The numbers over the tops of the columns in Table LXXXI are the common differences of the consecutive numbers in the columns. The numbers in the last column are the common differences of the consecutive numbers in the same horizontal line.

Exam. 1. Required the moon's longitude, latitude, equatorial parallax, semi-diameter, and hourly motions in longitude and latitude, on the 14th of October, 1838, at 6h. 54m. 34s. P. M. mean time at New York.

Mean time at New York, October,	14 <sup>d</sup> .	6 <sup>h</sup> .	54 <sup>m</sup> .	34 <sup>s</sup> .
Diff. of Long. - - - -		4	56	4
Mean time at Greenwich, October,	14	11	50	38



Eqs. D's Long.			Eqs. D's Lat.			Arguments.		D's Eq. Par.		Hourly Motion in Longitude.		Hourly Motion in Latitude.	
Arg.	°	' "	°	' "	Arg.	Evection.	Anomaly.	Variation.	Constant.	Arguments.	Equa.	Arguments.	Equa.
1	0	23 25.7	87	57 33.1	I.	.	.	.	.	2 of long.	"	I.	"
2	2	2 2.1	1	20.0	II.	.	.	.	.	3 do.	5.0	Pre. Eq. & Sum Eq.	33.8
3	0	6.8		10.5	D's long.	.	.	.	.	4 do.	1.0	II.	46.9
4	2	35.3		14.9	20 long.	.	.	.	.	5 do.	0.0	Pre. Eq. & Sum Eq.	2.9
5	0	23.3		8.5	V.	.	.	.	.	6 do.	0.3	Constant	1.4
6	0	56.6		6.4	VI.	.	.	.	.	Constant	1.5		1.0
7	0	22.0		31.3	VII.	.	.	.	.	Sum	3.0		86.0
8	0	10.8		21.7	VIII.	.	.	.	.	Evect. & Sum Eqs.	10.8		- 237.2
9	1	48.5		19.8	IX.	.	.	.	.	Evect.	0.2		- 151.2
10	0	10.7		6.6	X.	.	.	.	.	Sum	32.8		2' 31".2
11	0	18.0	88	1 0.8	Const.	.	.	.	.	An. & Sum Eqs.	11.6		
12	2	24.5	90	0 0.0		.	.	.	.	Anomaly	21.7		
13	8	4				.	.	.	.	Sum	66.1		
14	13	6				.	.	.	.	Var. & Sum Eqs.	9.4		
15	1	1.0				.	.	.	.	Variation	44.9		
16	8	2				.	.	.	.	Sum	120.4		
17	11	7				.	.	.	.	Red. & Sum Eqs.	2.6		
18	3	3				.	.	.	.	Reduction	4.1		
19	0	0.8				.	.	.	.	Sum	127.1		
20	10	1				.	.	.	.	Constant	2' 7".1		
21	10	1				.	.	.	.	Moon's Hourly Mot. in Long.	27' 24".0		
22	10	3				.	.	.	.				
23	10	3				.	.	.	.				
24	10	3				.	.	.	.				
25	10	3				.	.	.	.				
26	10	3				.	.	.	.				
27	10	3				.	.	.	.				
28	10	3				.	.	.	.				
29	10	3				.	.	.	.				
30	10	3				.	.	.	.				
31	10	3				.	.	.	.				
32	10	3				.	.	.	.				
33	10	3				.	.	.	.				
34	10	3				.	.	.	.				
35	10	3				.	.	.	.				
36	10	3				.	.	.	.				
37	10	3				.	.	.	.				
38	10	3				.	.	.	.				
39	10	3				.	.	.	.				
40	10	3				.	.	.	.				
41	10	3				.	.	.	.				
42	10	3				.	.	.	.				
43	10	3				.	.	.	.				
44	10	3				.	.	.	.				
45	10	3				.	.	.	.				
46	10	3				.	.	.	.				
47	10	3				.	.	.	.				
48	10	3				.	.	.	.				
49	10	3				.	.	.	.				
50	10	3				.	.	.	.				
51	10	3				.	.	.	.				
52	10	3				.	.	.	.				
53	10	3				.	.	.	.				
54	10	3				.	.	.	.				
55	10	3				.	.	.	.				
56	10	3				.	.	.	.				
57	10	3				.	.	.	.				
58	10	3				.	.	.	.				
59	10	3				.	.	.	.				
60	10	3				.	.	.	.				
61	10	3				.	.	.	.				
62	10	3				.	.	.	.				
63	10	3				.	.	.	.				
64	10	3				.	.	.	.				
65	10	3				.	.	.	.				
66	10	3				.	.	.	.				
67	10	3				.	.	.	.				
68	10	3				.	.	.	.				
69	10	3				.	.	.	.				
70	10	3				.	.	.	.				
71	10	3				.	.	.	.				
72	10	3				.	.	.	.				
73	10	3				.	.	.	.				
74	10	3				.	.	.	.				
75	10	3				.	.	.	.				
76	10	3				.	.	.	.				
77	10	3				.	.	.	.				
78	10	3				.	.	.	.				
79	10	3				.	.	.	.				
80	10	3				.	.	.	.				
81	10	3				.	.	.	.				
82	10	3				.	.	.	.				
83	10	3				.	.	.	.				
84	10	3				.	.	.	.				
85	10	3				.	.	.	.				
86	10	3				.	.	.	.				
87	10	3				.	.	.	.				
88	10	3				.	.	.	.				
89	10	3				.	.	.	.				
90	10	3				.	.	.	.				
91	10	3				.	.	.	.				
92	10	3				.	.	.	.				
93	10	3				.	.	.	.				
94	10	3				.	.	.	.				
95	10	3				.	.	.	.				
96	10	3				.	.	.	.				
97	10	3				.	.	.	.				
98	10	3				.	.	.	.				
99	10	3				.	.	.	.				
100	10	3				.	.	.	.				







Exam. 2. Required the moon's longitude, latitude, equatorial parallax, semi-diameter, and hourly motions in longitude and latitude, on the 9th of April, 1838, at 8h. 58m. 19sec. P. M. mean time at Washington.

Ans. Long.  $6^{\text{s.}} 19^{\circ} 45' 31''.2$ ; lat.  $36' 21''.9$  S.; equat. par.  $54' 36''.3$ ; semi-diameter  $14' 52''.7$ ; hor. mot. in long.  $30' 15''.2$ ; and hor. mot. in lat.  $2' 47''.0$ , tending south.\*

### PROBLEM XV.

*The Moon's Equatorial Parallax, and the Latitude of a Place, being given, to find the Reduced Parallax and Latitude.*

With the latitude of the place, take the reductions from Table LXIV, and subtract them from the Parallax and Latitude.

Exam. 1. Given the equatorial parallax  $55' 15''$ , and the latitude of New York  $40^{\circ} 42' 49''$  N., to find the reduced parallax and latitude.

Equatorial parallax	-	-	-	-	$55' 15''$
Reduction	-	-	-	-	$5$

Reduced parallax	-	-	-	-	$55 10$
------------------	---	---	---	---	---------

Latitude of New York	-	-	-	$40^{\circ} 42' 40''$ N.
----------------------	---	---	---	--------------------------

Reduction	-	-	-	-	$11 20$
-----------	---	---	---	---	---------

Reduced Lat. of New York	-	-	$40 31 20$
--------------------------	---	---	------------

2. Given the equatorial parallax  $60' 36''$  and the latitude of Baltimore  $39^{\circ} 17' 13''$  N., to find the reduced parallax and latitude.

Ans. Reduced par.  $60' 32''$ , and reduced lat.  $39^{\circ} 5' 59''$ .

3. Given the equatorial parallax  $57' 22''$ , and the latitude of New Orleans  $29^{\circ} 57' 45''$  N., to find the reduced parallax and latitude.

Ans. Reduced par.  $57' 19''$ , and reduced lat.  $29^{\circ} 47' 50''$ .

---

\* The smaller equations were omitted in working this example.

## PROBLEM XVI.

*To find the Longitude and Altitude of the Nonagesimal Degree of the Ecliptic, for a given time and place.*

For the given time reduced to mean time at Greenwich, find the sun's mean longitude and the argument N from Tables XVIII, XIX, XX, and XXI. To the sun's mean longitude, apply according to its sign the nutation in right ascension, taken from Table XXVII with argument N; and the result will be the sun's mean longitude, reckoned from the true equinox.

Reduce the mean time of day at the given place, expressed astronomically, to degrees, &c., and add it to the sun's mean longitude from the true equinox. The sum, rejecting  $360^\circ$ , when it exceeds that quantity, will be *the right ascension of the mid-heaven, or the sidereal time in degrees.*

Next, find the reduced latitude of the place by Problem XV; and when it is *north*, *subtract* it from  $90^\circ$ ; but when it is *south*, *add* it to  $90^\circ$ ; the sum or difference will be *the reduced distance of the place from the north pole.*

Also take the obliquity of the ecliptic for the given year from Table XXII.\*

These three quantities having been found, the longitude and altitude of the nonagesimal degree may be computed from the following formulæ :

$$\log. \cos. \frac{1}{2} (H - \omega) - \log. \cos. \frac{1}{2} (H + \omega) = A \dots (1);$$

$$\log. \tan \frac{1}{2} (H - \omega) + 10 - \log. \tan \frac{1}{2} (H + \omega) = B \dots (2);$$

$$\log. \tan E = A + \log. \tan \frac{1}{2} (S - 90^\circ) \dots (3);$$

$$\log. \tan F = \log. \tan E + B \dots (4);$$

$$N = E + F + 90^\circ \dots (5);$$

$$\log. \tan \frac{1}{2} h = \log. \cos. E + \log. \tan \frac{1}{2} (H + \omega) + \text{ar. co. log.} \cos. F - 20 \dots (6);$$

---

\* If great precision is required, the apparent obliquity is to be used in place of the mean. (See Prob. X.)

in which,

$H$  = the reduced distance of the place from the north pole ;

$\omega$  = the Obliquity of the Ecliptic ;

$S$  = the Sidereal Time converted into degrees ;

$N$  = the required Longitude of the Nonagesimal ;

$h$  = the required Altitude of the Nonagesimal ;

$E$  and  $F$  are auxiliary angles.

We first find the logarithmic sums  $A$  and  $B$ . With these we determine the angles  $E$  and  $F$  by formulæ (3) and (4), and with these again  $N$  and  $h$  by formulæ (5) and (6).

The angles  $E$ ,  $F$  are to be taken less than  $180^\circ$ ; and less or greater than  $90^\circ$ , according as the sign of their tangent proves to be positive or negative.

Note 1. In case the given place lies within the arctic circle, we must take, in place of formula (5), the following :

$$N = E - F + 90^\circ.$$

Note 2. As the obliquity of the ecliptic varies but slowly from year to year, the values which have once been found for the logarithms  $A$ ,  $B$ , and  $C$ , will answer for several years from the date of their determination, unless very great accuracy is required.

Note 3. The angle  $h$  derived from formula (6), is the distance of the zenith of the given place from the north pole of the ecliptic. This is not always equal to the altitude of the nonagesimal. Throughout the southern hemisphere, and frequently in the northern near the equator, it is the supplement of the altitude. In employing this angle in the following Problem, it is, however, for the sake of simplicity, called the altitude of the nonagesimal in all cases.

Exam. 1. Required the longitude and altitude of the nonagesimal degree of the ecliptic at New York, on the 18th of September, 1838, at 3h. 52m. 56sec. P. M. mean time.

The sun's mean longitude taken from the tables, for the given time, is  $5^h 27^\circ 19' 17''$ , and the argument  $N$  is 987. The nutation taken from Table XXVII with argument  $N$  is  $-1''$ . Hence, the sun's mean longitude from the true equinox is  $5^h 27^\circ$

19' 16". The given time of day, expressed astronomically, is 3h. 52m. 56sec. ; which in degrees is 58° 14' 0".

The reduced latitude of New York, found by Problem XV, is 40° 31' 20", and this taken from 90° leaves the polar distance 49° 28' 40". The obliquity of the ecliptic, derived from Table XXII, is 23° 27' 37".

Given time in degrees	-	-	-	58°	14'	0"
Sun's mean longitude	-	-	-	177	19	16
Sidereal time in degrees (S)	-	-	-	235	33	16
				90		
				2 )	145	33 16
H	-	-	-	49°	28'	40"
ω	-	-	-	23	27	37
Diff.	-	-	-	26	1	3
Sum	-	-	-	72	56	17
$\frac{1}{2}$ diff.	-	-	-	13	0	31
$\frac{1}{2}$ sum	-	-	-	36	28	8
				cos.	9.98870	- tan. + 10,19.36366
				cos.	9.90535	- tan. C. 9.86871
				A.	0.08335	B. 9.49495
					0.50866	
$\frac{1}{2}$ (S -- 90°)	72	46	38			
E	-	-	-	75	38	55
				tan.	0.59201	- cos. 9.39422
				B.	9.49495	C. 9.86871
F	-	-	-	50	41	55
				90	0	0
				$\frac{1}{2}$ alt. non.	16°	7' 54"
						tan. 9.46125
long. non.	216	20	50			
				alt. non.	32	15 48

2. Required the longitude and altitude of the nonagesimal degree of the ecliptic at New York, on the 10th of May, 1838, at 11h. 33m. 56sec. P. M. mean time.

Ans. Long. 200° 12' 23", and alt. 37° 0' 34".

## PROBLEM XVII.

*To find the Apparent Longitude and Latitude, as affected by Parallax, and the Augmented Semi-diameter of the Moon; the Moon's True Longitude, Latitude, Horizontal Semi-diameter, and Equatorial Parallax, and the Longitude and Altitude of the Nonagesimal Degree of the Ecliptic, being given.*

We have for the resolution of this Problem the following formulæ :

$$\log. x = \log. P + \log. \cos. h + \text{ar. co. log. cos. } \lambda - 10 \dots (1);$$

$$c = \log. x + \log. \text{tang } h - 10 \dots (2);$$

$$\log. u = c + \log. \sin K - 10 \dots (3);$$

$$\log. u' = c + \log. \sin (K + u) - 10 \dots (4);$$

$$\log. p = c + \log. \sin (K + u') - 10 \dots (5);$$

$$\text{Appar. long.} = \text{true long.} + p \dots (6);$$

$$\log. \text{tang } \lambda' = \log. p + \text{ar. co. log. cos. } \lambda + \text{ar. co. log. } u + \log. \sin (\lambda - x) - 10 \dots (7);$$

$$\log. v = \log. P + \log. \cos. h + \log. \cos. \lambda' - 10 \dots (8);$$

$$\log. z = \log. v + \log. \text{tang } h + \log. \text{tang } \lambda' + \log. \cos. (K + \frac{1}{2} p) - 30 : \dots (9);$$

$$\pi = v - z \dots (10);$$

$$\text{Appar. lat.} = \text{true lat.} - \pi \dots (11);$$

$$\log. R' = \log. p + \text{ar. co. log. cos. } \lambda + \text{ar. co. log. } u + \log. \cos. \lambda' + \log. R - 10 \dots (12);$$

in which,

$P$  = the Reduced Parallax of the Moon ;

$h$  = the Altitude of the Nonagesimal ;

$\lambda$  = the True Latitude of the Moon (minus when south) ;

$K$  = the Longitude of the Moon, minus the longitude of the Nonagesimal ;

$p$  = the required Parallax in Longitude ;

$\lambda'$  = the *approximate* Apparent Latitude of the Moon ;

$\pi$  = the required Parallax in Latitude ;



$R'$  = the True Semi-diameter of the Moon ;

$R'$  = the Augmented Semi-diameter of the Moon ;

$x, u, u', v, z$ , are auxiliary arcs.

Formulæ (1), (2), (3), (4), and (5), being resolved in succession, we derive the apparent longitude from formula (6) ; then the apparent latitude from equations (7), (8), (9), (10), (11) ; and lastly, the augmented semi-diameter from equation 12.

The latitude of the moon must be affected with the *negative* sign when *south* ; and the apparent latitude will be *south* when it comes out *negative*. In performing the operations, it is to be remembered that the *cosine* of a negative arc has the *same* sign as the cosine of a positive arc of an equal number of degrees ; but that the *sine* or *tangent* of a negative arc has the *opposite* sign from the sine or tangent of an equal positive arc. Attention must also be paid to the signs in the addition and subtraction of arcs. Thus, two arcs affected with essential signs, which are to be added to each other, are to be added *arithmetically*, when they have like signs, but subtracted if they have unlike signs ; and when one arc is to be taken from another, its sign is to be changed, and the two united according to their signs. An arithmetical sum, when taken, will have the same sign as each of the arcs ; and an arithmetical difference the same sign as the greater arc.

The use of negative arcs may be avoided, though the calculation would be somewhat longer, by using the true polar distance  $d$ , and the approximate apparent polar distance  $d'$ , in place of  $\lambda$  and  $\lambda'$ , substituting  $\sin d$  for  $\cos. \lambda$ ,  $\cos. (d + x)$  for  $\sin (\lambda - x)$ ,  $\sin d'$  for  $\cos. \lambda'$ ,  $\log. \text{co-tang } d'$  for  $\log. \text{tang } \lambda'$  ; and observing that  $p$  is to be subtracted from the true longitude in case the longitude of the nonagesimal exceeds the longitude of the moon ; that  $z$ , when it comes out negative, is to be added to  $v$ , which is always positive to the north of the tropic, otherwise subtracted ; and that the parallax in latitude is to be applied according to its sign to the true polar distance.

In seeking for the logarithms of the trigonometrical lines, it will be sufficient to take those answering to the nearest tens of seconds.

Note 1. When great accuracy is not desired,  $u'$  may be taken

for  $p$ , from which it can never differ more than a fraction of a second.

2. In solar eclipses, the moon's latitude is very small, and formula (7) may be changed into the following,

$\log. \lambda' = \log. p + \text{ar. co. log. cos. } \lambda + \text{ar. co. log. } u + \log. (\lambda - x) - 10$   
and  $\cos. \lambda'$  omitted in formula (12) without material error.

Formulæ (8), (9), (10), and (11), may also now be dispensed with, unless very great precision is desired, and the value of  $\lambda'$  given by the above formula taken for the apparent latitude.

It is to be observed also, that in eclipses of the sun  $P$  is taken equal to the reduced parallax of the moon minus the sun's horizontal parallax. By this the parallax of the sun in longitude and latitude is referred to the moon, and the relative apparent places of the sun and moon are correctly obtained, without the necessity of a separate computation of the sun's parallax in longitude and latitude.

Exam. 1. About the time of the middle of the occultation of the star Antares, on the 10th of May, 1838, the moon's longitude, by the *Connaissance des Temps*, was  $247^{\circ} 37' 6''.7$ ; latitude  $4^{\circ} 14' 14''.7$  S.; semi-diameter  $15' 24''.2$ ; and equatorial parallax  $56' 31''.7$ ; and the longitude of the nonagesimal at New York was  $200^{\circ} 12' 23''$ ; the altitude  $37^{\circ} 0' 34''$ ; required the apparent longitude and latitude, and the augmented semi-diameter of the moon at New York, at the time in question.

Equat. par.	56' 31''.7	Moon's long.	247° 37' 7"
Reduction	4 .6	Long. nonag.	200 12 23

$P = 56 \ 27 \ .1$

$K = 47 \ 24 \ 44$

$h = 37 \ 0 \ 34$

$\lambda = -4 \ 14 \ 14.7$

P	.	.	.	.	3387''.1	.	.	log.	3.52983
---	---	---	---	---	----------	---	---	------	---------

h	.	.	.	37°	0'	34''	.	.	cos.	9.90230
---	---	---	---	-----	----	------	---	---	------	---------

$a. \ 3.43213$

$\lambda$	.	.		- 4	14	15	.	Ar. co.	cos.	0.00119
-----------	---	---	--	-----	----	----	---	---------	------	---------

x	.	.	.		45	12	.	2712''	.	log.	3.43332
---	---	---	---	--	----	----	---	--------	---	------	---------

h	.	.	.	37	0	34	.	.	.	tan.	9.87725
---	---	---	---	----	---	----	---	---	---	------	---------

$c. \ 3.31057$

							<i>c.</i> 3.31057
K	-	-	-	47° 24' 44"	-	-	sin. 9.86701
<i>u</i>	-	-	-	25 5	- 1505"	-	log. 3.17758
							<i>c.</i> 3.31057
K + <i>u</i>	-	-	-	47 49 49	-	-	sin. 9.86991
<i>u'</i>	-	-	-	25 15	- 1515".2	-	log. 3.18048
							<i>c.</i> 3.31057
K + <i>u'</i>	-	-	-	47 49 59	-	-	sin. 9.86993
<i>p</i>	-	-	-	25 15.3	1515".3	-	log. 3.18050
True long.	-	-	-	247 37 6.7			
Appar. long.	-	-	-	248 2 22.0			
<i>p</i>	-	-	-	-	-	-	log. 3.18050
$\lambda - x$	-	-	-	4 59 27	-	-	sin. 8.93957—
$\lambda$	-	-	-	-	-	-	Ar. co. cos. 0.00119
<i>u</i>	-	-	-	-	-	-	Ar. co. log. 6.82242
$\lambda'$	-	-	-	5 1 10	-	-	tan. 8.94368—
$\lambda'$	-	-	-	5 1 10	-	-	cos. 9.99833
							<i>a.</i> 3.43213
<i>v</i>	-	-	-	44 54.4	2694".4	-	log. 3.43046
<i>h</i>	-	-	-	-	-	-	tan. 9.87725
$\lambda'$	-	-	-	-	-	-	tan. 8.94368—
K + $\frac{1}{2} p$	-	-	-	47 37 22	-	-	cos. 9.82867
<i>z</i>	-	-	-	—2 0.2	120".2	-	log. 2.08006—
<i>v</i> — <i>z</i>	-	-	-	46 54.6			
<i>v</i> — <i>z</i> (sign changed)	-	-	-	—46 54.6			
True lat.	-	-	-	—4 14 14.7			
Appar. lat.	-	-	-	5 1 9.3 S.			
<i>p</i>	-	-	-	-	-	-	log. 3.18050
$\lambda$	-	-	-	-	-	-	Ar. co. cos. 0.00119
<i>u</i>	-	-	-	-	-	-	Ar. co. log. 6.82242
$\lambda'$	-	-	-	-	-	-	cos. 9.99833
R	-	-	-	15 24.2	924".2	-	log. 2.96577
Augm. semi-diam.				15 29.4	929".4	-	log. 2.96821

Exam. 2. About the middle of the eclipse of the sun on the 18th of September, 1838, the moon's longitude was  $175^{\circ} 29' 19''.0$ , latitude  $47' 47''.5$ , equatorial parallax  $53' 53''.5$ , and semi-diameter  $14' 41''.1$ ; and the longitude of the nonagesimal at New York was  $216^{\circ} 20' 50''$ , the altitude  $32^{\circ} 15' 48''$ : required the apparent longitude and latitude, and the augmented semi-diameter of the moon.

Equat. paral.	$53' 53''.5$	Moon's long.	$175^{\circ} 29' 19''$
Reduction,	$4.4$	Long. nonag.	$216 \quad 20 \quad 50$
	<hr/>		<hr/>
	$53 \quad 49.1$	K =	$-40 \quad 51 \quad 31$
Sun's paral.	$8.6$	h =	$32 \quad 15 \quad 48$
	<hr/>		
	P = $53 \quad 40.5$	$\lambda =$	$0 \quad 47 \quad 47.5$
P	- - - - $3220''.5$	- - -	log. $3.50792$
h	- - $32^{\circ} 15' 48''$	- - -	cos. $9.92716$
$\lambda$	- - - $47 \quad 47.5$	- - -	Ar. co. cos. $0.00004$
			<hr/>
x	- - $45 \quad 23.5$	- $2723''.5$	- log. $3.43512$
h	- - $32 \quad 15 \quad 48$	- - -	- tan. $9.80023$
			<hr/>
			c. $3.23535$
K	- - $-40 \quad 51 \quad 31$	- - -	sin. $9.81570$ —
			<hr/>
u	- - $-18 \quad 45$	- $1125''$	- log. $3.05105$ —
			<hr/>
			c. $3.23535$
K + u	- - $-41 \quad 10 \quad 16$	- - -	sin. $9.81844$ —
			<hr/>
u'	- - $-18 \quad 52.9$	- $1132''.9$	- log. $3.05379$ —
			<hr/>
			c. $3.23535$
K + u'	- - $-41 \quad 10 \quad 24$	- - -	sin. $9.81844$ —
			<hr/>
p	- - $-18 \quad 52.9$	- $1132''.9$	- log. $3.05379$ —
True long.	$175 \quad 29 \quad 19.0$		
	<hr/>		
Appar. long.	$175 \quad 10 \quad 26.1$		

$p$	-	-	-	-	-	-	-	log.	3.05379
$\lambda$	-	-	-	-	-	-	Ar. co.	cos.	0.00004
$u$	-	-	-	-	-	-	Ar. co.	log.	6.94895
$\lambda - x$	-		$2' 24''.0$	-	$144''.0$	-	-	log.	2.15836

Appar. latitude	$2' 24''.9$	N.	$144''.9$	-	-	log.	2.16114
-----------------	-------------	----	-----------	---	---	------	---------

$p$	-	-	-	-	-	-	-	log.	3.05379
$\lambda$	-	-	-	-	-	-	Ar. co.	cos.	0.00004
$u$	-	-	-	-	-	-	Ar. co.	log.	6.94895
$R$	-	-	$14' 41''.1$	-	$881''.1$	-	-	log.	2.94502

Augm. semi-diam.	$14 46.7$	-	$886''.7$	-	-	log.	2.94780
------------------	-----------	---	-----------	---	---	------	---------

### PROBLEM XVIII.

*To find the Mean Right Ascension and Declination, or Longitude and Latitude of a Star, for a given time, from the Tables.*

Take the difference between the given year and 1840. Then seek in Table XV for the fraction of the year answering to the given month and days, and add it to this difference, if the given time is after the beginning of the year 1840; but if it is before, subtract it. Multiply the sum or difference by the annual variation given in the catalogue (Table XC), and the product will be the variation in the interval between the given time and the epoch of the catalogue. Apply this product to the quantity given in the catalogue, according to its sign, if the given time is after the beginning of the year 1840, but with the opposite sign if it is before, and the result will be the quantity sought.

Exam. 1. Required the mean right ascension and declination of the star Sirius on the 15th of August, 1842.

Interval between given time and beginn. of 1840 ( $t$ ),	2.619 yrs.
Annual variation of right ascension,	- - - 2.646s.

Variation of right ascension for interval $t$ ,	- - 6.93s.
---	------------

A similar operation gives for the variation of declination in the same interval,  $11''.65$ .



Mean right ascen. beginning of 1840, Table XC,	6 <sup>h</sup> . 38 <sup>m</sup> . 5.76 <sup>s</sup> .
Variation for interval $t$ ,	+ 6.93
	<hr/>

Mean right ascension required,	6 38 12.69
	<hr/>

Mean declination beginning of 1840,	16° 30' 4".79 S.
-------------------------------------	------------------

Variation for interval $t$ ,	+ 11.65
	<hr/>

Mean declination required,	16 30 16.44 S.
----------------------------	----------------

2. Required the mean longitude and latitude of Aldebaran on the 20th of October, 1838.

Interval between given time and begin. of 1840, ( $t$ )	1.200 yrs.
---	------------

Annual variation of longitude,	50".208
	<hr/>

Variation of longitude for interval $t$ ,	60".2
---	-------

A similar operation gives for the variation of latitude in the same interval 0".4.

Mean longitude beginning of 1840,	2 <sup>s</sup> . 7° 33' 5".4
-----------------------------------	------------------------------

Variation for interval $t$ ,	— 1 0.2
	<hr/>

Mean longitude required,	2 7 32 5.2
	<hr/>

Mean latitude beginning of 1840,	5° 28' 38".0 S.
----------------------------------	-----------------

Variation for interval $t$ ,	+ 0.4
	<hr/>

Mean latitude required,	5 28 38.4 S.
-------------------------	--------------

3. Required the mean right ascension and declination of Capella on the 9th of February, 1839 ?

Ans. Mean right ascension 5<sup>h</sup>. 4<sup>m</sup>. 48.74<sup>s</sup>., and mean declination 45° 49' 38".53 N.

4. Required the mean longitude and latitude of Aldebaran on the 16th of April, 1845 ?

Ans. Mean longitude 2<sup>s</sup>. 7° 37' 30".9, and mean latitude 5° 28' 36".2.

# PROBLEM XIX.

*To find the Aberration of a Star in Right Ascension and Declination, for a given Day.*

This problem may be resolved for any of the stars in the catalogue of Table XC by means of the following formulæ :

$$\log. (\text{aber. in right ascen.}) = \log. M + \log. \sin. (\odot + \varphi) - 10.$$

$$\log. (\text{aber. in declin.}) = \log. N + \log. \sin. (\odot + \theta) - 10,$$

in which M, N, are constants,  $\odot$  the longitude of the sun on the given day, and  $\varphi, \theta$ , auxiliary angles. Log M, log N, and the angles  $\varphi, \theta$ , are given for each of the stars in the catalogue, in Table XCI.  $\odot$  may be derived from an ephemeris of the sun, or it may be computed from the solar tables by Problem IX.

Exam. 1. What was the amount of aberration, in right ascension and declination of  $\alpha$  Orionis on the 20th of December, 1837, the sun's longitude on that day being  $8^{\text{s}}.28^{\circ}28'$ ?

Right Ascension.

Table XCI, $\varphi$	-	$6^{\text{s}}.3^{\circ}13'$	log. M	-	0.1361
$\odot$	-	$8\ 28\ 28$			

$$\odot + \varphi \quad \underline{3\ 1\ 41} \quad - \quad - \quad \sin. \quad \underline{9.9998}$$

$$\text{Aberration} = 1''.37 \quad - \quad - \quad - \quad - \quad \log. \quad 0.1359$$

Declination.

Table XCI, $\theta$	-	$8^{\text{s}}.28^{\circ}23'$	log. N	-	0.7521
$\odot$	-	$8\ 28\ 28$			

$$\odot + \theta \quad - \quad \underline{5\ 26\ 51} \quad - \quad - \quad \sin. \quad \underline{8.7399}$$

$$\text{Aberration} = 0''.31 \quad - \quad - \quad - \quad - \quad \log. \quad \underline{1.4920}$$

2. Required the aberrations in right ascension and declination of  $\alpha$  Andromedæ on the 1st of May, 1838, the sun's longitude being  $1^{\text{s}}.10^{\circ}38'$ .

Ans. Aberr. in right ascension  $-1''.07$ , and aberr. in declination  $-11''.70$ .

## PROBLEM XX.

*To find the Nutation of a Star in Right Ascension and Declination, for a given Day.*

This Problem may be solved by means of the formulæ,

$$\log. (\text{nuta. in right asc.}) = \log. M' + \log. \sin (\Omega + \varphi') - 10;$$

$$\log. (\text{nuta. in declinat.}) = \log. N' + \log. \sin (\Omega + \theta') - 10;$$

in which  $M'$ ,  $N'$  are constants,  $\Omega$  the mean longitude of the moon's ascending node, and  $\varphi'$ ,  $\theta'$  auxiliary angles.  $\log. M'$ ,  $\log. N'$ , and the angles  $\varphi'$ ,  $\theta'$  are given for each of the stars in the catalogue, in Table XCI. The mean longitude of the moon's ascending node is given for every tenth day of the year in the Nautical Almanac, page 266, and may be easily found for any intermediate day from the daily motion inserted at the foot of the column of longitudes. It may also be had by finding the supplement of the moon's node, for the given time, from the tables, and subtracting it from  $12^{\text{s.}} 0^{\circ} 7'$ .

Exam. 1. What was the amount of the nutation, in right ascension and declination, of  $\alpha$  Orionis on the 20th of December, 1837, the mean longitude of the moon's node on that day being  $18^{\circ} 54'$ ?

Right Ascension.

Table XCI, $\varphi'$	-	6 <sup>s.</sup> 0° 15'	log. $M'$	-	-	0.0481	)
$\Omega$	-	0 18 54					

$\Omega + \varphi'$	6	19	9	-	-	-	sin. 9.5159—
---------------------	---	----	---	---	---	---	--------------

Nutation =	—	0".37	-	-	log. 1.5640—
------------	---	-------	---	---	--------------

Declination.

Table XCI, $\theta'$	-	3 <sup>s.</sup> 2° 37'	log. $N'$	-	-	0.9657
$\Omega$	-	0 18 54				

$\Omega + \theta'$	3	21	31	-	-	-	sin. 9.9686
--------------------	---	----	----	---	---	---	-------------

Nutation =	8".60	-	-	log. 0.9343
------------	-------	---	---	-------------

2. Required the nutations in right ascension and declination of  $\alpha$  Andromedæ on the 1st of May, 1838.

Ans. Nutation in right ascension —  $0''.54$ , and nutation in declination —  $1''.43$ .

Note. When the apparent place of a star is desired with great accuracy, the *solar* nutations must also be estimated and allowed for. These may be determined by repeating the process for finding the lunar nutations, only using twice the sun's longitude in place of the longitude of the moon's node, and multiplying the results by the decimal .075.

The calculation of the solar nutations in Example 1st, is as follows :

Right Ascension.					
Table XCI,	$\phi'$	-	6 <sup>s</sup> .	0° 15',	log. M' - - 0.0481
	2 $\odot$	-	4	56 56	
<hr/>					
	2 $\odot$ + $\phi'$		10	57 11 - - -	sin. 9.7455—
<hr/>					
				— 0".06	- log. 1.7936—
				.075	

Solar Nutat. = — 0".00

Declination.					
Table XCI,	$\theta'$	-	3 <sup>s</sup> .	2° 37',	log. N' - - 0.9657
	2 $\odot$	-	4	56 56	
<hr/>					
			7	59 33 - - -	sin. 9.9999—
<hr/>					
				— 9".24	- - 0.9656—
				.075	

Solar Nutat. = — 0 .69

In Example 2d, we find for the solar nutation in right ascension, — 0".08, and for the solar nutation in declination, — 0".57.

## PROBLEM XXI.

*To find the Apparent Right Ascension and Declination of a Star on a given Day.*

Find the mean right ascension and declination for the given day by Problem XVIII ; then compute the aberrations in right ascension and declination by Problem XIX, and the lunar and solar nutations in right ascension and declination by Problem

XX. Apply the aberrations and nutations according to their signs, to the mean right ascension and declination on the given day, observing that the declination when south is to be marked negative, and the results will be the apparent right ascension and declination sought.

Exam. 1. What was the apparent right ascension and declination of  $\alpha$  Orionis on the 20th of December, 1837?

	h. m. s.			° ' "		
Table XC, M. right ascen.	5	46	30.71,	M. dec.	7	22 17.14N.
Variations	-		— 6.59	-	-	— 2.42
	<hr/>				<hr/>	
	5	46	24.12		7	22 14.72
Aberr. -	-		+ 1.37	-	-	+ 0.31
Lun. nutat. -	-		— 0.37	-	-	+ 8.60
Sol. nutat. -	-		0.00	-	-	— 0.69
	<hr/>				<hr/>	

App. right asc. 5 46 25.12, App. dec. 7 22 22.94N.

2. Required the apparent right ascension and declination of  $\alpha$  Andromedæ on the 1st of May, 1838.

Ans. Appar. right ascen. 0h. 0m. 0.90s., and appar. dec.  $28^{\circ} 11' 39'' 90$ .

## PROBLEM XXII.

*To find the Aberration of a Star in Longitude and Latitude, for a given Day.*

The formulæ for the computation are,

$$\log. (\text{aber. in long}) = 1.30880 + \log. \cos. (6s. + \odot - L) + \text{ar. co. log. } \cos. \lambda - 10;$$

$$\log. (\text{aber. in lat.}) = 1.30880 + \log. \sin. (6s. + \odot - L) + \log. \sin. \lambda - 20;$$

in which  $\odot$  = longitude of the sun on the given day;  $L$  = mean longitude of the star; and  $\lambda$  = mean latitude of the star.

Exam. 1. Required the aberrations in longitude and latitude of  $\alpha$  Scorpii, on the 26th of February, 1838. the sun's longitude on that day being  $11^{\text{s.}} 7^{\circ} 29'$ .



By Prob. XVIII,  $L = 8^s \ 7^\circ 30'$ , and  $\lambda = 4^\circ 32' \text{ S.}$

$$6s. + \odot - 17 \ 7 \ 29 \quad \text{Const. log. } 1.3088$$

$$6s. + \odot - L \ 8 \ 29 \ 59 \quad - \quad - \quad \cos. \ 6.4637—$$

$$\lambda \quad - \quad - \quad 4 \ 32 \quad \text{Ar. co. cos. } 0.0014$$

$$\text{Aberr. in long.} = - \ 0''.00 \ \log. \ 3.7739—$$

$$\text{Const. log. } 1.3088$$

$$6s. + \odot - L \ 8^s \ 29^\circ 59' \quad - \quad \sin. \ 9.2474—$$

$$\lambda \quad - \quad - \quad 4 \ 32 \quad - \quad \sin. \ 9.7099$$

$$\text{Aberr. in lat.} = - \ 1''.61, \ \log. \ 0.2661—$$

2. Required the aberrations in longitude and latitude of *Arc-turus* on the 5th of October, 1838, the sun's longitude being  $6^s \ 11^\circ 47'$ .

Ans. Aberr. in long. —  $23''.34$ , and aberr. in lat.  $1''.85$ .

Note. The *nutation* in longitude of a fixed star may be found after the same manner as the nutation in longitude of the sun. (See Problem IX, page 271).

### PROBLEM XXIII.

*To find the Apparent Longitude and Latitude of a Star, for a given Day.*

Find the mean longitude and latitude on the given day by Problem XVIII. Find also the aberrations in longitude and latitude by Problem XXII, and the nutation in longitude, as in Problem IX. Apply the aberration and nutation in longitude, according to their signs, to the mean longitude, and the result will be the apparent longitude; and apply the aberration in latitude according to its sign, to the mean latitude, and the result will be the apparent latitude.

Exam. 1. Required the apparent longitude and latitude of  $\alpha$  *Scorpii* on the 26th of February, 1838.

Table XC, M. long.	8 <sup>s</sup> . 7° 31' 45".2,	M. lat. 4° 32' 51".6 S.
Var.	— 1 32 .57	0 .78
	<hr/>	<hr/>
	8 7 30 12 .63	4 32 50 .82
Aberr.	0 .00	— 1 .61
Nutat.	— 4 .40	
	<hr/>	<hr/>

App. long. 8 7 30 8 .23, App. lat. 4 32 49 .21S.

2. Required the apparent longitude and latitude of Arcturus on the 5th of October, 1838.

Ans. Appar. long. 6<sup>s</sup>. 21° 58' 40".4, and appar. lat. 30° 51' 19".1.

## PROBLEM XXIV.

*To compute the Longitude and Latitude of a Heavenly Body from its Right Ascension and Declination, the Obliquity of the Ecliptic being given.*

This Problem may be solved by means of the following formulæ:

$$\log. \tan x = \log. \tan D + \text{ar. co. log. sin } R;$$

$$\log. \tan L = \log. \cos. (x - \omega) + \log. \tan R + \text{ar. co. log. cos. } x - 10;$$

$$\log. \tan \lambda = \log. \tan (x - \omega) + \log. \sin L - 10;$$

in which

R = the Right Ascension;

D = the Declination (minus when south);

L = the Longitude;

$\lambda$  = the Latitude;

$\omega$  = the Obliquity of the ecliptic;

$x$  is an auxiliary arc. It must be taken according to the sign of its tangent, but always less than 180°. The longitude will always be in the same quadrant as the right ascension. The latitude must be taken less than 90°, and will be *north* or *south*, according as the sign is *positive* or *negative*.

Note. When the mean longitude and latitude are to be derived from the mean right ascension and declination, the mean obli-

quity of the ecliptic is taken. When the apparent longitude and latitude are to be derived from the apparent right ascension and declination, found as in Problem XXI, the apparent obliquity is taken. The mean obliquity of the ecliptic at any assumed time is easily deduced from Table XXII. The apparent obliquity is found by Problem X.

Exam. 1. On the 20th of June, 1838, the right ascension of Capella was  $76^{\circ} 11' 29''$ , the declination  $45^{\circ} 49' 35''$  N., and the obliquity of the ecliptic  $23^{\circ} 27' 37''$ : required the longitude and latitude.

D = $45^{\circ} 49' 35''$	-	-	-	tan.	0.0125295
R = $76 11 29$	-	-	-	Ar. co. sin.	0.0127367
<hr/>					
x = $46 39 54$	-	-	-	tan.	0.0252662
$\omega = 23 27 37$					<hr/>
x — $\omega = 23 12 17$	-	-	-	cos.	9.9633641
R = $76 11 29$	-	-	-	tan.	0.6094483
x = $46 39 54$	-	-	-	Ar. co. cos.	0.1635095
<hr/>					
Long. = $79 36 3$	-	-	-	tan.	0.7363219
<hr/>					
L = $79 36 3$	-	-	-	sin.	9.9928071
x — $\omega = 23 12 17$	-	-	-	tan.	9.6321516
<hr/>					
Lat. = $22 51 47$	-	-	-	tan.	9.6249587

2. Given the right ascension of *Spica*  $199^{\circ} 11' 35''$ , and declination  $10^{\circ} 19' 24''$  S., and the obliquity of the ecliptic  $23^{\circ} 27' 36''$ , on the 1st of January, 1840, to find the longitude and latitude.

Ans. Long.  $201^{\circ} 36' 32''$ , and lat.  $2^{\circ} 2' 30''$  S.

## PROBLEM XXV.

*To compute the Right Ascension and Declination of a Heavenly Body from its longitude and latitude, the obliquity of the ecliptic being given.*

The formulæ for the solution of this problem are,

$$\log. \text{tang } y = \log. \text{tang } \lambda + \text{ar. co. log. sin } L;$$

$$\log. \text{tang } R = \log. \cos. (y + \omega) + \log. \text{tang } L + \text{ar. co. log. cos. } y - 10;$$

$$\log. \text{tang } D = \log. \text{tang } (y + \omega) + \log. \sin. R - 10;$$

in which

$L$  = the Longitude;

$\lambda$  = the Latitude (*minus* when South);

$R$  = the Right Ascension;

$D$  = the Declination;

$\omega$  = the Obliquity of the ecliptic;

$y$  is an auxiliary arc. It must be taken according to the sign of its tangent, but always less than  $180^\circ$ . The right ascension will always be in the same quadrant with the longitude. The declination must be taken less than  $90^\circ$ , and will be *north* or *south*, according as the sign is *positive* or *negative*.

Note. The mean or apparent obliquity of the ecliptic is taken, according as the given and required elements are mean or apparent.

Exam 1. On the 1st of January, 1830, the longitude of *Sirius* was  $3^h 11^m 44^s 18''$ , the latitude  $39^\circ 34' 1''$  S., and the obliquity of the ecliptic  $23^\circ 27' 41''$ : required the right ascension and declination.

$\lambda$	=	$39^\circ 34' 1''$	-	-	tan. 9.9171381 —
$L$	=	101 44 18	-	-	Ar. co. sin. 0.0091788
$y$	=	139 50 14	-	-	tan. 9.9263169 —
$\omega$	=	23 27 41	-	-	—
$y + \omega$	=	163 17 55	-	-	cos. 9.9812819 —
$L$	=	101 44 18	-	-	tan. 0.6823798 —
$y$	=	139 50 14	-	-	Ar. co. cos. 0.1167843 —
Right ascen.	=	99 24 48	-	-	tan. 0.7804460 —
$R$	=	99 24 48	-	-	sin. 9.9941121
$y + \omega$	=	163 17 55	-	-	tan. 9.4771803 —
Dec.	=	16 29 20 S.	-	-	tan. 9.4712924 —

2. Given the longitude of Aldebaran  $67^\circ 33' 5''$ , and latitude  $5^\circ 28' 38''$  S., and the obliquity of the ecliptic  $23^\circ 27' 36''$ , on the 1st of January, 1840, to find the right ascension and declination.

Ans. Right ascension  $66^\circ 41' 4''$ , and declination  $16^\circ 10' 57''$  N.

## PROBLEM XXVI.

*The Longitude and Declination of a Body being given, and also the Obliquity of the Ecliptic, to find the Angle of Position.*

The formula is

$$\log. \sin p = \log. \sin \omega + \log. \cos. L + \text{ar. co. log. cos. } D - 10;$$

$p$  = Angle of Position (required);

$L$  = Longitude;

$D$  = Declination;

$\pi$  = Obliquity of the ecliptic.

The angle of position  $p$  must be taken less than  $90^\circ$ . It is to be observed also that when the longitude is less than  $90^\circ$ , or more than  $270^\circ$ , the northern part of the circle of latitude lies to the *west* of the circle of declination, but that when the longitude is between  $90^\circ$  and  $270^\circ$ , it lies to the *east*.

Note. The angle of position may also be computed from the right ascension and latitude, by means of a formula similar to that just given, namely,

$$\log. \sin p = \log. \sin \omega + \log. \cos. R + \text{ar. co. log. cos. } \lambda - 10.$$

Exam. 1. Given the longitude of *Regulus*  $147^\circ 27' 54''$ , and declination  $12^\circ 47' 45''$  N., and the obliquity of the ecliptic  $23^\circ 27' 41''$ , to find the angle of position.

$\omega = 23^\circ 27' 41''$	-	-	sin. 9.6000260
$L = 147 \quad 27 \quad 54$	-	-	cos. 9.9258601
$D = 12 \quad 47 \quad 45$	-	Ar. co. cos.	0.0109217

Angle of pos. = $20 \quad 7 \quad 58$	-	-	sin. 9.5368078
---------------------------------------	---	---	----------------

The circle of latitude lies to the east of the circle of declination.

2. Given the longitude of Fomalhaut  $351^\circ 51' 38''$ , and declination  $30^\circ 31' 14''$  S., and the obliquity of the ecliptic  $23^\circ 27' 41''$ , to find the angle of position. Ans.  $27^\circ 13' 36''$ .

The circle of latitude lies to the west of the circle of declination.



## PROBLEM XXVII.

*To find from the Tables the Time of New or Full Moon, for a given Year and Month.*

*For the New Moon.*

Take from Table LXXXVI, the time of mean new moon in January, and the Arguments I, II, III and IV, for the given year. Take from Table LXXXVII, as many lunations with the corresponding variations of Arguments I, II, III, and IV, as the given month is months past January, and add these quantities to the former, rejecting the ten thousands from the sums in the columns of the first two arguments, and the hundreds from the sums in the columns of the other two. Seek the number of days from the first of January to the first of the given month, in the *second* or *third* column of Table LXXXVIII, according as the given year is a *common* or *bissextile* year, and subtract it from the sum in the column of mean new moon : the remainder will be tabular time of mean new moon for the given month. It will sometimes happen that the number of days taken from Table LXXXVIII, will exceed the number of days of the sum in the column of mean new moon : in this case one lunation more, with the corresponding arguments, must be added.

With the sums in the columns I, II, III, and IV, as arguments, take the corresponding equations from Table LXXXIX, and add them to the time of mean new moon : the sum will be *Approximate* time of new moon for the given month, expressed in mean time at Greenwich.

Next, for the approximate time of new moon calculate the true longitudes and hourly motions in longitude of the sun and moon ; subtract the less longitude from the greater, and the hourly motion of the sun from the hourly motion of the moon ; and say, as the difference between the hourly motions : the difference between the longitudes : : 60 minutes : the correction of the approximate time. The correction *added* to the approximate time, when the sun's longitude is *greater* than the moon's, but *subtracted*, when it is *less*, will give the true time of new moon required, in mean

time at Greenwich. This time may be reduced to the meridian of any given place by Problem V.

For Full Moon.

Take from Table LXXXVI, the time of mean new moon, and the corresponding Arguments I, II, III, and IV, for January of the given year, and from Table LXXXVII, a half lunation with the corresponding changes of the arguments. Then, when the time of mean new moon for January is on or after the 16th, subtract the latter quantities from the former, increasing, when necessary to render the subtraction possible, either or both of the first two arguments by 10,000, and of the last two by 100 ; but add them when the time is before the 16th. The result will be the tabular time of mean full moon and the corresponding arguments, for January. Proceed to find the approximate time of full moon after the same manner as directed for the new moon.

For the approximate of full moon calculate the true longitudes and hourly motions in longitude of the sun and moon. Subtract the sun's longitude from the moon's, adding 360° to the latter if necessary. Take the difference between the remainder and VI signs, and call the result R. Also subtract the hourly motion of the sun from the hourly motion of the moon. Then say, as the difference between the hourly motions : R :: 60m. : the correction of the approximate time. The correction *added* to the approximate time of full moon, when the excess of the moon's longitude over the sun's is *less* than VI signs, but *subtracted* when it is *greater*, will give the true time of full moon.

Exam. 1. Required the time of new moon in September, 1838, expressed in mean time at New-York.

	M. New Moon.	I.	II.	III.	IV.
	d. h. m.				
1838,	24 16 53	0681	9175	99	85
8 lun.	236 5 52	6468	5737	22	93
	260 22 45	7149	4912	21	78
Days,	243				
Sept'r,	17 22 45				
I.	0 16				
II.	9 35				
III.	3				
IV.	10				
Sept'r,	18 8 49	Approximate time.			

Moon's true long. found for approx. time, is  $5^{\text{s.}} 25^{\circ} 29' 19''$   
 Sun's do. do. do.  $5 25 27 27$

Difference, - - - - -  $1 52$

Moon's hourly motion in long. is - - -  $29 28$

Sun's do. do. - - -  $2 27$

Difference, - - - - -  $27 1$

As  $27' 1'' : 1' 52'' :: 60^{\text{m}} : 4^{\text{h}} 9^{\text{s.}}$ , the correction.

Approx. time of new moon, September, -  $18^{\text{d.}} 8^{\text{h.}} 49^{\text{m.}} 0^{\text{s.}}$

Correction, - - - - -  $—4 9$

True time, in mean time at Greenwich, -  $18 8 44 51$

Diff. of meridians, - - - - -  $4 56 4$

True time, in mean time at New York, -  $18 3 48 47$

Exam. 2. Required the time of full moon in April, 1838, expressed in mean time at New York.

	M. Full Moon.	I.	II.	III.	IV.
1838,	d. h. m.				
$\frac{1}{2}$ lun.	24 16 53	0681	9175	99	85
	14 18 22	404	5359	58	50
3 lun.	9 22 31	0277	3816	41	35
	88 14 12	2425	2151	46	97
Days,	98 12 43	2702	5967	87	32
	90				
April,	8 12 43				
I.	8 29				
II.	16 7				
III.	15				
IV.	30				
April,	9 14 4	Approximate time.			

Moon's true long. found for approx. time, is  $6^{\text{s.}} 19^{\circ} 44' 17''$

Sun's do. do. do.  $0 19 45 22$

$5 29 58 55$

$6 0 0 0$

R  $1 5$

Moon's hourly motion in long. is	-	-	30'	15"
Sun's do. do.	-	-	2	27
Difference	-	-	-	27 48

As 27' 48" : 1' 5" :: 60<sup>m.</sup> : 2<sup>m.</sup> 20<sup>s.</sup>, the correction.

Approximate time of full moon, April,	9 <sup>d.</sup>	14 <sup>h.</sup>	4 <sup>m.</sup>	0 <sup>s.</sup>
Correction, - - - - -			+ 2	20
True time, in mean time at Greenwich,	9	14	6	20
Diff. of meridians, - - - - -		4	56	4
True time, in mean time at New York,	9	9	10	16

3. Required the time of new moon in September, 1837, expressed in mean time at Philadelphia; taking the longitudes for the approximate time from the Nautical Almanac.

Ans. 29d. 3h. 0m. 5s.

4. Required the time of full moon, in October, 1837, expressed in mean time at Boston.

Ans. 13d. 6h. 30m. 25s.

PROBLEM XXVIII.

*To determine the number of Eclipses of the Sun and Moon that may be expected to occur in any given Year, and the Times nearly at which they will take place.*

*For the Eclipses of the Sun.*

Take, for the given year, from Table LXXXVI the time of mean new moon in January, the arguments and the number N. If the number N differs less than 37 from either 0, 500, or 1000, an eclipse must occur at that new moon. If the difference is between 37 and 53, there may be an eclipse, but it is doubtful, and the doubt can only be removed by a calculation of the true places of the moon and sun. If the difference exceeds 53, an eclipse is impossible.

If an eclipse may or must occur at the new moon in January, calculate the approximate time of new moon by Problem XXVII, and it will be the time nearly of the middle of the eclipse, ex-

pressed in mean time at Greenwich. This may be reduced to the meridian of any other place by Problem V.

To find the first new moon after January, at which an eclipse of the sun may be expected, seek in column N of Table LXXXVII the first number after that answering to the half lunation, that, added to the number N for the given year, will make the sum come within 53 of 0, 500, or 1000. Take the corresponding lunations, changes of the arguments, and the number N, and add them, respectively, to the mean new moon in January, the arguments, and the number N, for the given year. Take from the *second* or *third* column of Table LXXXVIII, according as the given year is a *common* or *bissextile* year, the number of days next less than the days of the sum in the column of mean new moon, and subtract it from this sum; the remainder will be the tabular time of mean new moon in the month corresponding to the days taken from Table LXXXVIII. At this new moon there may be an eclipse of the sun; and if the sum in the column N is within 37 of the numbers mentioned above, there must be one. Find the approximate time of new moon, and it will be the time nearly of the middle of the eclipse.

If any of the other numbers in the last column of Table LXXXVII are found, when added to the number N of the given year, to give a sum that falls within the limit 53, proceed in a similar manner to find the approximate times of the eclipses.

Note. When the sum of the numbers N, or the number N itself, in case the eclipse happens in January, is a little above 0, or a little less than 500, the moon will be to the north of the sun, and there is a *probability* that the eclipse will be visible at any given place in north latitude at which the approximate time of the eclipse found as just explained and reduced to the meridian of the place comes during the day-time. When the number N found for the eclipse is more than 500, the moon will be to the south of the sun, and the eclipse will seldom be visible in the northern hemisphere except near the equator.

*For the Eclipses of the Moon.*

Find the time of full moon and the corresponding arguments and number N, for January of the given year, as explained in Problem XXVII. Then proceed to find the times at which



eclipses of the moon may or must occur, after the same manner as for eclipses of the sun, only making use of the limits 35 and 25, instead of 53 and 37.\*

Note. An eclipse of the moon will be visible at a given place, if the time of the eclipse thus found nearly and reduced to the meridian of the place comes in the night.

Exam. 1. Required the eclipses that may be expected in the year 1840, and the times nearly at which they will take place.

*For the Eclipses of the Sun.*

	M. New Moon.	I.	II.	III.	IV.	N.
	d. h. m.					
1840,	3 10 30	0085	6386	65	63	844
2 lun.	59 1 28	1617	1434	31	98	170
	62 11 58	1702	7820	96	61	014
	60					
March	2 11 58	As the sum of the numbers N comes within 37 of 0, there must be an eclipse.				
I.	8 3					
II.	19 38					
III.	12					
IV.	13					
March	3 16 4	Mean time at Greenwich.				

	M. New Moon.	I.	II.	III.	IV.	N.
	d. h. m.					
1840,	3 10 30	0085	6386	65	63	844
8 lun.	236 5 52	6468	5737	22	93	682
	239 16 22	6553	2123	87	56	526
	213					
Aug.	26 16 22	As the sum of the numbers N comes within 37 of 500, there must be an eclipse.				
I.	0 54					
II.	0 49					
III.	15					
IV.	16					
Aug.	26 18 36	Mean time at Greenwich.				

\* The numbers 53, 37, and 35, 25, are the lunar and solar ecliptic limits, as determined by Delambre. The limits given in the text, converted into thousandth parts of the circle, are 55, 37, and 37, 21.

*For the Eclipses of the Moon.*

	M. Full Moon.	I.	II.	III.	IV.	N.
	d. h. m.					
1840,	3 10 30	0085	6386	65	63	844
$\frac{1}{2}$ lun.	14 18 22	404	5359	58	50	43
	18 4 52	489	1745	23	13	887
1 lun.	29 12 44	808	717	15	99	85
	47 17 36	1297	2462	38	12	972
	31					
Febr.	16 17 36	As the sum of the numbers N, although it comes within 35 of 1000, does not come within 25, the eclipse may be considered doubtful.				
I.	7 27					
II.	0 23					
III.	5					
IV.	27					
Febr.	17 1 58	Mean time at Greenwich.				

	M. Full Moon.	I.	II.	III.	IV.	N.
	d. h. m.					
1840,	18 4 52	489	1745	23	13	887
7 lun.	206 17 8	5659	5020	7	94	596
	224 22 0	6148	6765	30	07	483
	213					
Aug.	11 22 0	As the sum of the numbers N comes within 25 of 500, there must be an eclipse.				
I.	1 37					
II.	19 16					
III.	3					
IV.	25					
Aug.	12 19 21	Mean time at Greenwich.				

2. Required the eclipses that may be expected in the year 1839, and the times nearly at which they will take place, expressed in mean civil time at New York.

Ans. One of the sun on the 15th of March at 9h. 20m. A. M.; and one of the sun on the 7th of September at 5h. 24m. P. M.

3. Required the eclipses that may be expected in the year 1841, and the times nearly at which they will take place, expressed in mean civil time at New York.

Ans. Four of the sun, namely, one on the 22nd of January at 12h. 18m. P. M.; one on the 21st of February at 6h. 17m. A. M.; one on the 18th of July at 9h. 24m. A. M.; and one on the 16th

of August at 4h. 28m. P. M. : and two of the moon, namely, one on the 5th of February at 9h. 10m. P. M. ; and one on the 2d of August at 5h. 5m. A. M.

The eclipses of the sun in January and August may be considered as doubtful.

## PROBLEM XXIX.

*To calculate an Eclipse of the Moon.*

The calculation of the circumstances of a lunar eclipse is effected with the following fundamental data, derived from the tables of the sun and moon :

Approximate Time of Full Moon (at Greenwich)	T
Sun's Longitude at that time, - - -	L
Do. Hourly Motion, - - - -	s
Do. Semi-diameter, - - - -	$\delta$
Do. Parallax, - - - -	p
Moon's Longitude, - - - -	l
Do. Latitude, - - - -	$\lambda$
Do. Equatorial Parallax, - - -	P
Do. Semi-diameter, - - - -	d
Do. Hourly Motion in longitude, - -	m
Do. Hourly Motion in latitude, - -	n

We obtain the time T by Problem XXVII ; the quantities appertaining to the sun, namely, L, s, and  $\delta$ , by Problem IX ;\* and those which have relation to the moon, namely, l,  $\lambda$ , P, d, m, and n, by Problem XIV.

From these quantities we derive the following :

True Time of Full Moon, (at given place) -	T'
Moon's Latitude at that time, - - -	$\lambda'$
Semi-diameter of earth's shadow, - - -	S
Inclination of Moon's relative orbit, - -	I

T being known, T' is found as explained in Problem XXVII. To obtain  $\lambda'$ , we state the following proportion,

$$1 \text{ hour} : \text{correction for the time of full moon} :: n : x ;$$

---

\* p may be taken = 9".

from this we deduce the value of  $x$ ; and thence find  $\lambda'$  by the equation,

$$\lambda' = \lambda \pm x.$$

When the true time of full moon, expressed in mean time at Greenwich, is *later* than the approximate time, the *upper* sign is to be used, if the latitude is *increasing*, the *lower* if it is *decreasing*; but when the true time is *earlier* than the approximate time, the *lower* sign is to be used if the latitude is *increasing*; the *upper*, if it is *decreasing*.

The value of  $S$  is derived from the equation,

$$S = (P + p - \delta) + \frac{1}{60} (P + p - \delta);$$

and the angle  $I$  from the formula,

$$\log. \tan g I = \log. n + \text{ar. co. log. } (m - s).$$

The foregoing quantities having all been determined, the various circumstances of the eclipse may be calculated by the following formulæ :

*For the Time of the Middle of the Eclipse.*

$$3.55630 + \log. \cos. I + \text{ar. co. log. } (m - s) - 10 = R;$$

$$\log. t = R + \log. \lambda' + \log. \sin I - 10;$$

$$M = T' \pm t:$$

$t$  = interval between time of middle of eclipse and time of full moon;  $M$  = time of middle of the eclipse.

The *upper* sign is to be taken in the last equation when the latitude is *decreasing*; the *lower*, when it is *increasing*.

*For the Times of Beginning and End.*

$$\log. c = \log. \lambda' + \log. \cos. I - 10;$$

$$\log. v = \frac{\log. (S + d + c) + \log. (S + d - c)}{2} + R;$$

$$B = M - v, \text{ and } E = M + v:$$

$v$  = half duration of the eclipse;  $B$  = time of beginning; and  $E$  = time of end.

Note. If  $c$  is equal to or greater than  $S + d$ , there cannot be an eclipse.

*For the Times of Beginning and End of the Total Eclipse.*

$$\log. v' = \frac{\log. (S - d + c) + \log. (S - d - c)}{2} + R;$$

$$B' = M - v', \text{ and } E' = M + v';$$

$v'$  = half duration of the total eclipse;  $B'$  = time of beginning of total eclipse; and  $E'$  = time of end of total eclipse.

Note. When  $c$  is greater than  $S - d$ , the eclipse cannot be total.

*For the Quantity of the Eclipse.*

$$\log. Q = 0.77815 + \log. (S + d - c) + \text{ar. co. log. } d - 10;$$

$Q$  = the quantity of the eclipse in digits.

Note 1. An eclipse of the moon begins on the eastern limb, and ends on the western. In partial eclipses the southern part of the moon is eclipsed when the latitude is north, and the northern part when the latitude is south.

Note 2. When the eclipse commences before sunset, and ends after sunset, the moon will rise more or less eclipsed. To obtain the quantity of the eclipse at the time of the moon's rising, find the moon's hourly motion on the relative orbit by the equation,

$$\log. h = \log. (m - s) + \text{ar. co. log. cos. } I;$$

in which  $h$  = hourly motion or relative orbit. Also find the interval between the time of sunset and the time of the middle of the eclipse, which call  $i$ . Then,

$$1 \text{ hour} : i :: h : x.$$

Deduce the value of  $x$  from this proportion, and substitute it in the equation,

$$c' = \sqrt{c^2 + x^2};$$

in which  $c$  designates the same quantity as in previous formulæ. Find the value of  $c'$ , and use it in place of  $c$  in the above formula for the quantity of the eclipse, and it will give the quantity of the eclipse at the time of the moon's rising. When the eclipse begins before and ends after sunrise, the quantity of the eclipse at the time of the moon's setting may be found in the same manner, only using sunrise instead of sunset.

Example. Required to calculate, for the meridian of New-York, the eclipse of the moon in October, 1837.



*Elements.*

Approximate time of full moon,	-	T = 11 <sup>h</sup> . 10 <sup>m</sup> . (Oct. 13)
Sun's longitude at that time,	- -	L = 6 <sup>s</sup> . 20° 24' 28"
Do. hourly motion,	- - -	s = 2 29
Do. semi-diameter,	- - -	δ = 16 4
Do. parallax,	- - -	p = 9
Moon's longitude,	- - -	l = 0 20 21 51
Do. latitude,	- - -	λ = 11 28
Do. equatorial parallax,	- - -	P = 59 32
Do. semi-diameter,	- - -	d = 16 13
Do. hourly motion in long.	- -	m = 35 54
Do. hourly motion in lat. (tending north),	n =	3 19

Approx. time of full moon, October,	-	13 <sup>d</sup> . 11 <sup>h</sup> . 10 <sup>m</sup> . 00 <sup>s</sup> .
Correction found by Prob. XXVII,	-	+ 4 42

True time, in mean time at Greenwich,	-	13 11 14 42
Diff. of meridians,	- - -	4 56 4

True time, in mean time at New York, T' =	13 6 18 38
---	------------

$$60^m : 4^m. 42^s : : 3' 19'' : x = 16''.$$

Moon's lat. at approx. time,	- - -	λ = 11' 28" S.
Correction,	- - -	x = — 16
Moon's lat. at true time,	- - -	λ' = 11 12
Moon's equatorial parallax,	- - -	P = 59' 32"
Sun's do.	- - -	p = 9
Sum,	- - -	59 41
Sun's semi-diameter,	- - -	δ = 16 4
Diff.	- - -	P + p — δ = 43 37
Add	- - -	$\frac{1}{60}$ (P + p — δ) = 44
Semi-diameter of earth's shadow,	- - -	S = 44 21

Moon's hor. mot. less sun's ( $m - s$ ) = 2005" - Ar. co. log. 6.69789

Moon's hor. motion in latitude,  $n = 199$  - log. 2.29885

Inclination of rel. orbit, I = 5° 40' - tan. 8.99674

*Time of Middle.*

									3.55630
I	-	-	-	-	5° 40'	-	-	cos.	9.99787
m — s	-	-	-	-	-	2005''	Ar. co.	log.	6.69789
									<hr/>
								R.	0.25206
λ	-	-	-	-	-	672''	-	log.	2.82737
I	-	-	-	-	5° 40'	-	-	sin.	8.99450
									<hr/>
t	-	-	-	-	0 <sup>h</sup> . 1 <sup>m</sup> . 58 <sup>s</sup> .	= 118 <sup>s</sup> .	-	log.	2.07393
T'	-	-	-	-	6 18 38	P. M.			
									<hr/>
Middle,	-	-	-	-	6 20 36	P. M.			

*Times of Beginning and End.*

λ'	-	-	-	-	-	-	-	log.	2.82737
I	-	-	-	-	-	-	-	cos.	9.99787
									<hr/>
c	-	-	-	-	11' 9'' = 669''	-	-	log.	2.82524
S + d + c	-	-	-	-	-	4303''	-	log.	3.63377
S + d — c	-	-	-	-	-	2965	-	log.	3.47202
									<hr/>
							2 )	7.10579	
									<hr/>
								3.55289	
								R.	0.25206
									<hr/>
v	-	-	-	-	1 <sup>h</sup> . 46 <sup>m</sup> . 22 <sup>s</sup> .	= 6382 <sup>s</sup> .	-	log.	3.80495
Middle,	-	-	-	-	6 20 36				
									<hr/>
Beginning,	-	-	-	-	4 34 14	P. M.			
End,	-	-	-	-	8 6 58	P. M.			
									<hr/>
S — d + c	-	-	-	-	-	2357''	-	log.	3.37236
S — d — c	-	-	-	-	-	1019	-	log.	3.00817
									<hr/>
							2 )	6.38053	
									<hr/>
								3.19026	
								R.	0.25206
									<hr/>
					0 <sup>h</sup> . 46 <sup>m</sup> . 9 <sup>s</sup> .	= 2769 <sup>s</sup> .	-	log.	3.44232



$\theta$  is an auxiliary arc. The value of  $\theta$  being derived from the first equation, the second will then make known the value of  $\Delta$ .

$\alpha$  and  $\lambda'$  are in every instance to be effected with the positive sign.\*

*For the Approximate Times of Beginning, Greatest  
Obscuration, and End*

Let the time for which the above calculations are made, be denoted by  $T$ . If the apparent distance of the centres of the sun and moon, found for the time  $T$ , is less than the sum of their apparent semi-diameters, there is an eclipse at this time. But if it is greater, either the eclipse has not yet commenced, or it has already terminated. It has not commenced if the apparent longitude of the moon is less than the longitude of the sun; and has terminated, if the apparent longitude of the moon is greater than the longitude of the sun.

1. If there should be an eclipse at the time  $T$ , from the sun's longitude and hourly motion in longitude, and the moon's longitude and latitude, and hourly motions in longitude and latitude, found for this time, calculate the longitudes and the moon's latitude for two instants respectively an hour before, and an hour after the time  $T$ . The semi-diameter of the sun, and the equatorial parallax and semi-diameter of the moon, may, in our present inquiry, be regarded as remaining the same during the eclipse. Find the apparent longitude and latitude, and the augmented semi-diameter of the moon (using in all cases the relative parallax), and thence compute by the formulæ already given, the apparent distance of the centres of the sun and moon at the two instants in question.

Observe for each result, whether it is less or greater than the sum of the apparent semi-diameters of the two bodies. If the

\* The apparent distance or the centres  $\Delta$  may be found without the aid of logarithms by means of the following equation:

$$\Delta = \sqrt{a^2 + \lambda'^2}.$$

If the logarithmic formulæ are used, it will be sufficient here to take out the angle  $\theta$  to the nearest minute. When we have occasion to obtain the distance of the centres exact to within a small fraction of a second,  $\theta$  must be taken to the nearest tens of seconds, if it exceeds  $20^\circ$  or  $30^\circ$ .

moon is apparently on the same side of the sun at the times  $T$  and  $T + 1h.$ , take the difference of the distances of the two bodies in apparent longitude at these times, but if it is on opposite sides, take their sum, and it will be the variation of this distance in the hour following  $T$ . Find in like manner the variation of the distance during the hour preceding  $T$ . Then, if the apparent distance of the centres at the times  $(T - 1h.)$ ,  $(T + 1h.)$  is less than the sum of the apparent semi-diameters, deduce from these results the variations of the distance in apparent longitude during the preceding and following hours, allowing for the second difference, and observing whether the two bodies are approaching each other or receding from each other. Thence, find the distance in apparent longitude at the times  $(T - 2h.)$ ,  $(T + 2h.)$ . Find by the same method the apparent latitude of the moon at the instants  $(T - 2h.)$ ,  $(T + 2h.)$ , observing that the variation of the apparent latitude in any given interval is the difference between the latitudes at the beginning and end of it, if they are both of the same name; their sum, if they are of opposite names.

From these results derive the apparent distance of the centres of the sun and moon at the two instants in question.

If there should still be an eclipse at the time  $(T + 2h.)$  or  $(T - 2h.)$ , find by the same method the distance of the centres at the time  $(T + 3h.)$  or  $(T - 3h.)$ . These calculations being effected, the times of the beginning, greatest obscuration, and end of the eclipse will fall between some of the instants  $T$ ,  $(T - 1h.)$ ,  $(T + 1h.)$  &c., for which the apparent distance of the centres is computed.

2. If the eclipse occurs after the time  $T$ , the different phases will happen between the instants  $T$ ,  $(T + 1h.)$ ,  $(T + 2h.)$ , &c. Find the apparent distance of the centres of the sun and moon for the times  $(T + 1h.)$ ,  $(T + 2h.)$ , by the same method as that by which it is found for the times  $(T + 1h.)$ ,  $(T - 1h.)$ , in the case just considered. Then, if the eclipse has not terminated, deduce the distance of the moon from the sun in apparent longitude, and the moon's apparent latitude, for the time  $(T + 3h.)$ , from these distances and latitudes at the times  $T$ ,  $(T + 1h.)$ ,  $(T + 2h.)$ ; as in the preceding case the distance and latitude for the time  $(T + 2h.)$  were deduced from the same at the times  $(T - 1h.)$ ,  $T$ ,  $(T + 1h.)$ . With the results obtained compute



the apparent distance of the centres of the two bodies at the time ( $T + 3h$ .)

3. In case the eclipse occurs before the time  $T$ , the apparent distance of the centres must be found by similar methods for the times ( $T - 1h$ .), ( $T - 2h$ .), &c.

The calculation is to be continued until the distance, from being less, becomes greater than the sum of the semi-diameters.

Now, let  $h$  = variation of apparent distance of centres in the interval of one hour comprised between the first two of the instants for which the distance is computed;  $d$  = difference between the sum of the semi-diameters of the sun and moon and the apparent distance of their centres at the first instant; and  $t$  = interval between first instant and the time of the beginning of the eclipse. Then,

$$h : d :: 60^m : t \text{ (nearly).}$$

Find the value of  $t$  given by this proportion, and add it to the time at the first instant, and the result will be a first approximation to the time of the beginning of the eclipse, which call  $b$ . Find, by interpolation,\* the distance of the moon from the sun in apparent longitude ( $a$ ), and the moon's apparent latitude ( $\lambda'$ ), for this time, and thence compute the apparent distance of the centres. Take  $h$  = variation of apparent distance in the interval between the time  $b$  and the nearest of the two instants above mentioned, between which the beginning falls, and  $d$  = difference between the apparent distance of the centres at the time  $b$  and the sum of the semi-diameters, and compute again the value of  $t$ . Add

---

\* The second differences may easily be taken into the account in finding the quantities  $a$  and  $\lambda'$  for the time  $b$ . Thus, let  $k$  = variation of  $a$  for the interval of an hour comprised between the instants above mentioned,  $k'$  = same for the succeeding hour, and  $i$  = interval between  $b$  and the nearer of the two instants, (in minutes). Then, if we put  $f = \frac{k}{6}$ ,  $c = \frac{k - k'}{36}$ , and  $v$  = var. of  $a$  in interval  $i$ ,

$$v = \frac{i \left\{ f \pm \left( c + \frac{c}{2} \right) \right\}}{10}.$$

The upper sign is to be used when the time  $b$  is nearer the first than the second instant, the lower when it is nearer the second than the first. The error by this method will not exceed the number  $c$  (supposing the changes of  $k, k'$  from 10m. to 10m. to increase or decrease by equal degrees).

this to the time  $b$ , or subtract it from it, according as  $b$  is before or after the beginning, and the result will be a second approximation to the time of the beginning, which call  $B$ . If necessary, a result still more approximate may be had, by taking  $h$  = variation of apparent distance of centres in the interval  $B - b$ ,  $d$  = difference between apparent distance at the time  $B$  and sum of semi-diameters, finding anew the value of  $t$  given by the preceding proportion, and adding it to or subtracting it from, as the case may be, the time  $B$ .

The end of the eclipse will fall between the last two of the several instants for which the apparent distance of the centres of the moon and sun have been computed. The approximate time of the end is found by the same method as that of the beginning.\*

The middle of the interval between the approximate times of the beginning and end of the eclipse, will be a first approximation to the time of greatest obscuration.

Note. When the object is merely to prepare for an observation, results sufficiently near the truth may be obtained by a graphical construction. The elements of the construction are the difference of the apparent longitudes of the moon and sun, and the apparent latitude of the moon, found as above, for two or more instants during the continuance of the eclipse. Draw a right line  $EF$  (Fig. 78), to represent the ecliptic, assume on it some point  $C$  for the position of the sun at the instant of apparent conjunction, and lay off  $CA$ ,  $CA'$ , equal to the two differences of apparent longitude; and to the right or left, according as the moon is to the west or east of the sun at the instants for which the calculations have been made. Erect the perpendiculars  $Ap$ ,  $A'p'$ , and mark off  $Aa$ ,  $A'a'$  equal to the two apparent latitudes. Through  $a$ ,  $a'$ , draw a right line, and it will be the apparent relative orbit of the moon, or will differ but little from it. From  $C$  let fall the perpendicular  $Cm$  upon the relative orbit,  $m$  will be the apparent place of the moon at the instant of greatest obscuration. Take a distance in the dividers equal to

---

\* In effecting the reductions of the quantities  $a$  and  $\lambda'$  to the first approximate time of end,  $k'$  must stand for the variation of  $a$  during the hour preceding that comprised between the last two instants, and the last instant must be substituted for the first. (See Note, p. 323.)

the sum of the apparent semi-diameters of the moon and sun, and placing one foot of it at C, mark off with the other the points  $f, f'$ , for the beginning and end of the eclipse, and by means of a square mark on E F the points  $b, e$ , which answer to the beginning and end. If the eclipse be total or annular, mark the points of immersion and emersion,  $g, g'$ , with an opening in the dividers equal to the difference of the semi-diameters, and find the corresponding points  $b', e'$  on the line E F.

If the calculations are made from hour to hour, the distance A A' is the apparent relative hourly motion of the sun and moon in longitude. This distance laid off repeatedly to the right and left, will determine the points 1, 2, &c., answering to 1h., 2h., &c. before and after the times for which the calculations are made. If the spaces in which the points  $b, e$ , answering to the beginning and end of the eclipse, occur, be divided into quarters, and then sub-divided into three equal parts or five minute spaces, the approximate times of the beginning and end of the eclipse will become known.

From the point  $m$ , as a centre, describe the lunar disc; and from the point C, as a centre, describe the sun's disc, and we shall have the figure of the greatest eclipse. The quantity of the eclipse will result from the proportion,

$$SN : MN :: 12 : \text{number of digits eclipsed.}$$

Draw from the centre C to the place of commencement  $f$ , the line C  $f$ ; and through the same point C raise a perpendicular to the ecliptic. With the declination and longitude of the sun at the time of the beginning, calculate its angle of position by Problem XIII, and lay it off in the figure, placing the circle of declination C P to the left if the tangent of the angle of position be positive, to the right if it be negative.

Compute also for the time of beginning, the angle of the vertical of the sun with the circle of declination, that is, the angle P S Z in Figure 18, for which we have in the triangle P S Z the side P S = co-declination, the side P Z = co-latitude, and the included angle Z P S. (The requisite formulæ are given in the Appendix). Form this angle in the figure at the point C, placing C Z to the right or left of C P, according as the time is in the forenoon or afternoon, C Z will be the vertical, and Z the *vertex*, or highest point of the sun. The arc Z  $t$  on the limb of

the sun, will be the angular distance from the vertex of the point on the limb at which the eclipse commences.

*For the True Times of Beginning, Greatest Obscuration, and End.*

The approximate times of beginning, greatest obscuration, and end of the eclipse, being calculated by the rules which have been given, find from the tables, the moon's longitude, latitude, equatorial parallax, semi-diameter, and hourly motions in longitude and latitude, for the approximate time of greatest obscuration. With the moon's longitude and latitude, and hourly motions in longitude and latitude, found for this time, calculate the longitude and latitude for the approximate times of beginning and end. The parallax and semi-diameter may, without material error, be considered the same during the eclipse. With the moon's true longitude, latitude and semi-diameter at the approximate times of beginning, greatest obscuration, and end, calculate its apparent longitude and latitude, and augmented semi-diameter, for these several times, (making use of the relative parallax). With the sun's longitude and hourly motion previously found for the approximate time of new moon, find his longitude at the approximate times of beginning, greatest obscuration, and end. The sun's semi-diameter found for the approximate time of new moon will serve also for any time during the eclipse. With the data thus obtained, calculate by the formulæ given on page 320, the apparent distance of the centres of the sun and moon at the approximate times of the three phases.

Note. When very great accuracy is required, the moon's longitude, latitude, equatorial parallax, semi-diameter, and hourly motions in longitude and latitude, must be calculated directly from the tables, for the approximate times of the beginning and end, as well as for that of the greatest obscuration.

*For the Beginning.*

Subtract the apparent longitude of the moon at the approximate time of beginning from the true longitude of the sun at the same time, and denote the difference by  $a$ . Do the same for the approximate time of greatest obscuration. Subtract the latter result from the former, paying attention to

the signs, and call the remainder  $k$ . Next, take the difference between the apparent latitudes of the moon at the approximate times of beginning and greatest obscuration, if they are of the same name; their sum, if they are of opposite names; and denote the difference or sum, as the case may be, by  $n$ . This done, compute the correction to be applied to the approximate time of beginning, by means of the following formulæ:

$$\log. b = \log. a + \log. k + \text{ar. co. log. } n - 10;$$

$$c = \lambda' - b, \quad S = d + \delta - 5'';$$

$$\log. t = \log. (S + \Delta) + \log. (S - \Delta) + \text{ar. co. log. } n + \text{ar. co. log. } c + \log. L + 1.47712 - 20;$$

in which,

$t$  = Correction of approx. time of beginn. (required);

$a$  = Diff. of appar. long. of Moon and Sun at approx. time;

$L$  = Half duration of eclipse in minutes (known approximately);

$k$  = Appar. relative motion of Sun and Moon in long. in the interval  $L$ ;

$n$  = Moon's appar. motion in lat. in same interval;

$\lambda'$  = Moon's appar. lat.;

$d$  = Augmented semi-diameter of Moon;

$\delta$  = Semi-diam. of Sun;

$\Delta$  = Appar. distance of centres of Sun and Moon.

$b$  and  $c$  are auxiliary quantities.

First find the value of  $b$  by the first equation, and substitute it in the second. Then derive the values of  $c$  and  $S$  from the second and third equations, and substitute them in the fourth, and it will make known the value of  $t$ , which is to be applied to the approximate time of the beginning of the eclipse according to its sign.

The quantities  $a$ ,  $k$ ,  $n$ , &c. are all to be expressed in seconds. The apparent latitude  $\lambda'$  must be affected with the *negative* sign, when it is *south*. The motion in latitude,  $n$ , must also have the negative sign in case the moon is apparently receding from the north pole.  $a$  and  $k$  are always positive.



The result may be verified, and corrected, by computing the apparent distance of the centres at the time found, and comparing it with the sum of the semi-diameters minus  $5''$ .

Note. When great precision is desired, the quantities  $k$  and  $n$  must be found for some shorter interval than the half duration of the eclipse. Let some instant be fixed upon, some five or ten minutes before or after the approximate time of the beginning of the eclipse, according as the contact takes place before or after. For this time deduce the longitude and latitude of the moon, from the longitude and latitude at the approximate time of beginning, by means of their hourly variations; and thence calculate the apparent longitude and latitude, and the augmented semi-diameter. Find the longitude of the sun for the time in question, from its longitude and hourly motion already known for the approximate time of beginning. Then proceed according to the rule given above, only using the quantities thus found for the time assumed, in place of the corresponding quantities answering to the approximate time of greatest obscuration.  $L$  will always represent the interval for which  $k$  and  $n$  are determined.

*For the End.*

Subtract the longitude of the sun at the approximate time of the end from the apparent longitude of the moon at the same time. Do the same for the approximate time of greatest obscuration. Then proceed according to the rule for the beginning, only substituting every where the approximate time of the end for the approximate time of the beginning, and taking in place of the formula  $c = \lambda' - b$ , the following:

$$c = \lambda' + b.$$

*For the Greatest Obscuration.*

Take the sum of the distances of the moon from the sun in apparent longitude at the approximate times of the beginning and end of the eclipse, and call it  $k$ . Take the difference of the apparent latitudes of the moon at the same times, if the two are of the same name; but if they are of different names, take their sum. Denote the difference or sum by  $n$ . Let  $a' =$  the distance of the moon from the sun in apparent longitude at the true time of greatest obscu-

ration ;  $\lambda'$  = the apparent latitude of the moon at the approximate time of greatest obscuration,

$$k : n :: \lambda' : a'.$$

Find the value of  $a'$  by this proportion, affecting  $\lambda'$ ,  $n$ ,  $k$ , always with the positive sign.

Ascertain whether the greatest obscuration has place before or after the apparent conjunction, by observing whether the apparent latitude of the moon is increasing or decreasing about this time ; the rule being, that when it is *increasing*, the greatest obscuration will occur *before* apparent conjunction ; when it is *decreasing*, *after*. If the approximate and true times of greatest obscuration are both before or both after apparent conjunction, from the value found for  $a'$  subtract the distance of the moon from the sun in apparent longitude at the approximate time ; but if one of the times is before and the other after apparent conjunction, take the sum of the same quantities. Denote the difference or sum by  $m$ . Also let  $D$  = duration of eclipse, and  $t$  = correction to be applied to the approximate time of greatest obscuration. Then to find  $t$ , we have the proportion

$$k : m :: D : t.$$

If the apparent latitude of the moon is decreasing,  $t$  is to be applied according to the sign of  $m$  ; but if the apparent latitude is increasing, it is to be applied according to the opposite sign.

A still more exact result may be had by repeating the foregoing calculations, making use now of the apparent latitude at the time just found. When the greatest accuracy is required, the values of  $k$  and  $n$  may be found more exactly after the same manner as for the beginning or end.

### *For the Quantity of the Eclipse.*

Find by interpolation the apparent latitude of the moon at the true time of greatest obscuration. With this, and the distance in longitude  $a$  obtained by the proportion above given, compute by the formulæ on page 320, the apparent distance of the centres of the sun and moon at the time of greatest obscuration. Subtract this distance from the sum of the apparent semi-diameter of the two bodies, and denote the remainder by  $R$ . Then,

Sun's semi-diam. :  $R$  :: 6 digits : number of digits eclipsed.

When the apparent distance of the centres of the sun and moon at the time of greatest obscuration is less than the difference between the sun's semi-diameter and the augmented semi-diameter of the moon, the eclipse is either *annular* or *total*; *annular*, when the sun's semi-diameter is the *greater* of the two; *total*, when it is the *less*.

*For the Beginning and End of the Annular or Total Eclipse.*

The times of the beginning and end of the annular or total eclipse may be found as follows: the greatest obscuration will take place very nearly at the middle of the eclipse in question, and will not differ, at most, more than five or eight minutes (according as the eclipse is total or annular) from the beginning and end: to obtain the half duration of the eclipse, and thence the times of the beginning and end, we have the formulæ

$$\log. \tan \delta = \log. \lambda' + \text{ar. co. log. } a, \log. k' = \log. k + \text{ar. co. log. } \sin \theta$$

$$S = \delta - d - 1'', \text{ or } S = d - \delta + 1'';$$

$$\log. c = \frac{\log. (S + \Delta) + \log. (S - \Delta)}{2};$$

$$\log. t = \text{ar. co. log. } k' + \log. c + \log. D + 1.77815 - 10.$$

$$\text{Time of Begin.} = M - t, \text{ Time of End} = M + t,$$

in which.

$M$  = Time of greatest obscuration.

$\lambda'$  = Moon's apparent latitude at that time.

$a$  = Distance of moon from sun in appar. long.

$k$  = Variation of this distance during the whole eclipse, or relative mot. in appar. long. during this interval.

$k'$  = Moon's appar. mot. on relative orbit for same interval.

$\theta$  = Inclination of relative orbit.

$\delta$  = Semi-diameter of sun.

$d$  = Augm. semi-diam. of moon.

$\Delta$  = Appar. distance of centres.

$D$  = Duration of eclipse (partial *and* annular or total).

$t$  = Half duration of annular or total eclipse.

The *first* value of  $S$  is used when the eclipse is *annular*, the *second* when it is *total*. The quantities may all be regarded as positive. The results may be verified and corrected by finding

directly the apparent distance of the centres for the times obtained, and comparing it with the value of  $S$ .

*For the Point of the Sun's Limb at which the Eclipse commences.*

Find the angle of position of the sun, and the angle between its vertical circle and circle of declination, at the beginning of the eclipse, as explained at page 325. Let the former be denoted by  $p$ , and the latter by  $v$ . Give to each the negative sign, if laid off towards the right; the positive sign, if laid off towards the left. Let  $a$  = distance of the moon from the sun in apparent longitude at the beginning of the eclipse;  $\lambda'$  = the moon's apparent latitude at the same time; and  $\theta$  = angular distance of the point of contact from the ecliptic. Compute the angle  $\theta$  by the formula,

$$\log. \tan \theta = \log. \lambda' + \text{ar. co. log. } a;$$

taking it always less than  $90^\circ$ , and positive or negative according to the sign of its tangent.  $\lambda'$  is negative when south;  $a$  is always positive.

Let  $A$  = distance on the limb of the point of contact from the vertex. The above operations being performed, the value of  $A$  results from the equation,

$$A = p + v + 90^\circ - \theta;$$

$p$ ,  $v$ , and  $\theta$  being taken with their signs.

If the result is affected with the positive sign, the point first touched will lie to the right of the vertex. If with the negative sign, it will lie to the left of the vertex.

Note. The circumstances of an occultation of a fixed star by the moon may be calculated in nearly the same manner as those of a solar eclipse. The star in the occultation holds the place of the sun in the eclipse. The immersion and emersion of the star correspond to the beginning and end of the eclipse. The elements which ascertain the relative apparent place and motion of the moon and star, take the place of those which ascertain the relative apparent place and motion of the moon and sun. Thus the star's longitude, corrected for aberration and nutation (see Problem XXIII), must be used instead of the sun's longitudes; the apparent distances of the moon from the star in latitude, instead of the moon's apparent latitudes; and the moon's augmented

semi-diameter, instead of the sum of the semi-diameters of the sun and moon. The difference of the longitudes, and the relative motion in longitude, must also now be reduced to a parallel to the ecliptic passing through the star, (see Art. 458, page 187). If  $\lambda$  = apparent latitude of star,  $a$  = diff. of appar. longitudes of moon and star, and  $k$  = relative motion in longitude, we must substitute in the formulæ for the eclipse, for  $\lambda'$ ,  $\lambda' - \lambda$ ; for  $a$ ,  $a \cos \lambda$ ; and for  $k$ ,  $k \cos \lambda$ .  $n$  will stand for the relative motion in latitude, or for the variation of  $\lambda' - \lambda$ .

Example. Required to calculate an eclipse of the sun, for the latitude and meridian of New York, that will occur on the 18<sup>th</sup> of September, 1838.

*For the Approximate Times of the Phases.*

Approximate time of New Moon.

Sept. 18<sup>d</sup>. 8<sup>h</sup>. 49<sup>m</sup>.

Sun's longitude, - - - - -	175° 27' 31".4
Do. hourly motion, - - - - -	2 26 .7
Do. semi-diameter, - - - - -	15 57 .0
Moon's longitude, - - - - -	175 29 19
Do. latitude, - - - - -	47 47
Do. equatorial parallax, - - - - -	53 53
Do. semi-diameter, - - - - -	14 41
Do. hor. mot. in long. - - - - -	29 29
Do. hor. mot. in lat. - - - - -	2 41
Do. appar. long. (Prob. XVII), - - - - -	175 10 26
Do. appar. lat. ( $\lambda'$ ), - - - - -	2 25 N.
Do. augm. semi-diameter, - - - - -	14 47
Diff. of appar. long. ( $a$ ), - - - - -	17 5
Appar. dist. of cen. ( $\Delta$ ), - - - - -	17 15
Sum of semi-diameters, - - - - -	30 44

7<sup>h</sup>. 49<sup>m</sup>.

Sun's longitude, - - - - -	175° 25' 4"
Moon's appar. long. - - - - -	174 47 3
Do. appar. lat. ( $\lambda'$ ), - - - - -	8 12 N.
Do. augm. semi-diameter, - - - - -	14 49
Diff. of appar. long. ( $a$ ), - - - - -	38 1



Appar. dist. of cen. ( $\Delta$ ),	-	-	-	-	38' 53"
Sum of semi-diameters,	-	-	-	-	30 46

9h. 49m.

Sun's longitude, -	-	-	-	-	175° 29' 58"
Moon's appar. long.	-	-	-	-	175 36 15
Do. appar. lat. ( $\lambda'$ ),	-	-	-	-	2 18 S.
Do. augm. semi-diameter,	-	-	-	-	14 44
Diff. of appar. long. ( $\alpha$ ),	-	-	-	-	6 17
Appar. dist. of cen. ( $\Delta$ ),	-	-	-	-	6 42
Sum of semi-diameters, -	-	-	-	-	30 41

	$a$	diff. or $k$	$\lambda'$	diff. or $n$	$\Delta$	diff.	sum semid.
7h 49m.	2281"		492" N		2333"		1846"
8 49	1025	1256"	145 N	347"	1035	1298"	1844
9 49	377	1402	138 S	283	402		1841
10 49	1925	1548	357 S	219	1958	1556	1839

*For the Approximate Time of Beginning.*

$$h = 1298'', d = 2333'' - 1846'' = 487'';$$

$$1298'' : 487'' :: 60^m : t = 22^m.5$$

7h. 49m.

22

1st Approxi. 8h. 11m.

7h. 49m.	-	$a = 2281''$	-	$\lambda' = 492''$ N
Correction for 22m.	447		-	133 (See Note, p. 323)

$$8h. 11m. - a = 1834 - \lambda' = 359$$

$$a = 1834'' \text{ ar. co. log. } 6.73660 - \text{ log. } 3.26340$$

$$\lambda' = 359 \text{ log. } 2.55509$$

$$\theta = 11^\circ 4' 30'' \text{ tan. } 9.29169 - \text{ ar. co. cos. } 0.00817$$

$$\text{Appar. dist. of cen. } \Delta = 1869'' - \text{ log. } 3.27157$$

$$\text{Sum of semi-diam. } - - 1846$$

$$487'' : 23'' :: 22^m. : t = 1^m. 2^s.$$

8<sup>h</sup>. 11<sup>m</sup>.

+ 1

2d Approx. 8<sup>h</sup>. 12<sup>m</sup>.*For the Approximate Time of the End.*

$$h = 1556'', d = 1958'' - 1839'' = 119''.$$

$$1556'' : 119'' :: 60^m : t = 4^m.6.$$

10<sup>h</sup>. 49<sup>m</sup>.

— 5

1st Approx. 10<sup>h</sup>. 44<sup>m</sup>.

$$10^h. 49^m. - a = 1925'' - - \lambda' = 357'' S.$$

$$\text{Correction for } 5^m. \quad 132 - - \quad 17$$

$$10^h. 44^m. - a = 1793 - - \lambda' = 340 S.$$

$$a = 1793'' \quad \text{Ar. co. log. } 6.74642 - \log. 3.25358$$

$$\lambda' = 340 - - \log. 2.53148$$

$$\delta = - - - \tan. 9.27790 - \text{Ar. co. cos. } 0.00767$$

$$\text{Appar. dist. of cen. } \Delta = 1825'' - 3.26125$$

1839

$$133'' : 14'' :: 5^m : t = 0^m.5.$$

10<sup>h</sup>. 44<sup>m</sup>.

0 .5

2d Approx. 10<sup>h</sup>. 44<sup>m</sup>.5*For the Approximate Time of Greatest Obscuration.*

$$\text{Approx. time of begin.} - 8^h. 12^m.$$

$$\text{Approx. time of end,} - 10 \quad 44$$

$$2 \quad ) \quad 18 \quad 56$$

$$\text{1st Approx.} - 9 \quad 28$$

*For the True Times of the Phases.*

	Approx. time of Beginning. 8 <sup>h</sup> . 12 <sup>m</sup> .	Approx. time of Greatest Obscur. 9 <sup>h</sup> . 28 <sup>m</sup> .	Approx. time of End. 10 <sup>h</sup> . 44 <sup>m</sup> .
Sun's longitude, 175° 26' 1.0,	175° 29' 6.8,	175° 32' 12.6	
Do. semi-diam., 15 57.0,	15 57.0,	15 57.0	
Moon's app. long. 174 55 36.7,	175 27 7.7,	176 2 17.2	
Do. app. lat. 5 45.3 N,	0 43.5 S,	5 32.4 S.	
Do. augm. semid. 14 48.0,	14 45.1,	14 41.7	

	<i>a</i>	<i>k</i>	$\lambda'$	<i>n</i>	$\Delta$	<i>S</i>
8 <sup>h</sup> . 12 <sup>m</sup> .	1824".3	1705".2	345".3 N	388".8	1856".7	1840".0
9 28	119 .1	1923 .7	43 .5 S	288 .9	1835 .0	1833 .7
10 44	1804 .6		332 .4 S			

*For the True Time of Beginning.*

<i>a</i> - 1824".3	-	-	-	-	-	log. 3.26109
<i>k</i> - 1705 .2	-	-	-	-	-	log. 3.23178
<i>n</i> - 388 .8	-	-	-	-	Ar. co. log. 7.41028—	
<i>b</i> = — 8001 .1	-	-	-	-	-	log. 3.90315—
$\lambda'$ - 345 .3						
$\lambda' - b = c = 8346 .4$	-	-	-	-	Ar. co. log. 6.07850	
<i>S</i> + $\Delta$ - 3696 .7	-	-	-	-	-	log. 3.56781
<i>S</i> — $\Delta$ - —16 .7	-	-	-	-	-	log. 1.22272—
<i>n</i> - -	-	-	-	-	Ar. co. log. 7.41028—	
<i>T</i> - 76m.	-	-	-	-	-	log. 1.88081
					Const. log. 1.47712	
Corr. of approx. time, + 43 <sup>s</sup> .4	-	-	-	-	-	log. 1.63724 +
Approx. time, 8 <sup>h</sup> . 12 <sup>m</sup> . 0 .0						
True time of begin. 8 12 43 .4,						in Greenwich time.
Diff. of merid. - 4 56 4						
True time of begin. 3 16 39 .4,						in New York time.

*For the True Time of End.*

$a$	-	-	1804".6	-	-	-	-	log. 3.25638
$k$	-	-	1923 .7	-	-	-	-	log. 3.28414
$n$	-	-	288 .9	-	-	-	Ar. co. log. 7.53925—	
$b =$	—		12016 .3	-	-	-	-	log. 4.07977—
$\lambda'$	-	—	332 .4					
<hr/>								
$\lambda' + b = c =$	—		12348 .7	-	-	-	Ar. co. log. 5.90838—	
$S + \Delta$	-	-	3668 .7	-	-	-	-	log. 3.56451
$S - \Delta$	-	-	—1 .3	-	-	-	-	log. 0.11394—
$n$	-	-	-	-	-	-	Ar. co. log. 7.53925—	
$T$	-	-	76m.	-	-	-	-	1.88081
							Const. log. 1.47712	
<hr/>								
Corr. of approx. time,			— 3 <sup>s</sup> .0	-			log. 0.48401—	
Approx. time,	-		10 <sup>h</sup> . 44 <sup>m</sup> . 0 .0					
<hr/>								
True time of end,		10	43	57 .0,	in Greenwich time.			
Diff. of merid.	-		4	56	4			
<hr/>								
True time of end,		5	47	53,	in New York time.			

*For the True Time of Greatest Obscuration.*

True time of beginning,	-	-	-	18 <sup>h</sup> . 12 <sup>m</sup> 43 <sup>s</sup> .4
Do. of end,	-	-	-	10 43 57 .0

---

2) 18 56 40 .4

---

2d Approx. 9 28 20 .2

9<sup>h</sup>. 49<sup>m</sup>. - -  $\lambda' = 138''$  S.

9 28 - -  $\lambda' = 43 .5$  S.

---

Diff. 21

---

Diff. 94 .5

21<sup>m</sup>. : 20<sup>s</sup>. :: 94<sup>''</sup>.5 : 1<sup>''</sup>.5

" 43 .5

---

9<sup>h</sup>. 28<sup>m</sup>. 20<sup>s</sup>.

---

$\lambda' = 45 .0$





$$\begin{array}{rcll} a & - & 12''.0 & - \text{ Ar. co. log. } 8.92082, & - & - & \text{ log. } 1.07918 \\ \lambda' & - & 65.1 & - & - & - & 1.81358 \end{array}$$

$$\theta \quad - \quad - \quad - \quad - \quad \tan. 0.73440, \quad - \quad \text{ Ar. co. cos. } 0.74165$$

$$\text{Shortest distance of centres,} \quad - \quad 66''.2 \quad - \quad - \quad \text{ log. } 1.82083$$

$$\text{Sum of semi-diameters,} \quad - \quad 1837.0$$

$$\hline 1770.8$$

$$15' 57'' : 1770''.8 :: 6 : 11.1 \text{ digits eclipsed.}$$

*For the Situation of the Point at which the Obscuration commences.*

$$8^h. 12^m. \quad - \quad - \quad a = 1824'', \quad - \quad - \quad \lambda' = 345''.3 \text{ N.}$$

$$76^m. : 43^s. :: 1705'' : 16, 76^m. : 43^s. :: 389'' : 3.7$$

$$\text{At the beginn.} \quad - \quad a = 1808, \quad - \quad - \quad \lambda' = 341.6$$

$$a \quad - \quad 1808 \quad - \quad \text{ Ar. co. log. } 6.74275$$

$$\lambda' \quad - \quad 341.6 \quad - \quad \text{ log. } 2.53352$$

$$\theta = 10^\circ 41' 53'' \quad - \quad \tan. 9.27627$$

$$\text{Obliq. of eclip. (Prob. X), } 23^\circ 27' 47'' \text{ sin. } 9.60005 \quad - \quad \tan. 9.63753$$

$$\text{Sun's longitude,} \quad - \quad 175 \quad 26 \quad 3 \text{ sin. } 8.90093 \quad - \quad \cos. 9.99862$$

$$\hline \text{sin. } 8.50098, \quad \tan. 9.63615$$

$$\text{Sun's declination, } 1^\circ 49' 0''; \text{ Angle of pos. } 23^\circ 23' 59''.$$

$$\text{Mean time of begin. } 3^h. 16^m. 39^s, \text{ Lat. } 40^\circ 42' 40'', \text{ Dec. } 1^\circ 49' 0''$$

$$\text{Equa. of time,} \quad - \quad 5 \quad 58 \quad 90 \quad 90$$

$$\text{Appar. time,} \quad - \quad 3 \quad 22 \quad 37, \text{ PZ} = 49 \quad 17 \quad 20, \text{ PS} = 88 \quad 11$$

$$\hline 4) \quad 202 \quad 37$$

$$\text{Hour angle P} = 50^\circ 39' 15'' \quad - \quad \cos. 9.80210$$

$$\text{Co. lat. PZ} = 49 \quad 17 \quad 20 \quad - \quad \tan. 0.06526$$

$$m = 36^\circ 23' 0'' \quad - \quad - \quad \tan. 9.86736$$

$$\text{Co. dec. PS} = 88 \quad 11 \quad 0$$

$$\hline m' = 51 \quad 48 \quad 0 \quad - \quad \text{ Ar. co. sin. } 0.10466$$

$$m' = 51 \ 48 \ 0 \quad - \quad \text{Ar. co. sin. } 0.10466$$

$$m = 36 \ 23 \ 0 \quad - \quad - \quad \text{sin. } 9.77320$$

$$P = 50 \ 39 \ 15 \quad - \quad - \quad \text{tan. } 0.08627$$

---


$$S = 42 \ 38 \ 10 \quad - \quad - \quad \text{tan. } 9.96413$$

$$\text{Angle of position,} \quad - \quad - \quad - \quad 23^\circ 23' 50''$$

$$\text{Angle from eclip. } (\theta), \quad - \quad - \quad - \quad 10 \ 41 \ 50$$

$$\text{Angle of dec. circle from vertex (S),} \quad 42 \ 38 \ 10$$

$$90$$

---

Angular dist. of point first touched from vertex, 98 32, to the right.

*For the Beginning and End of the Annular Eclipse.*

Approx. time, 9<sup>h</sup>. 32<sup>m</sup>. 29<sup>s</sup>. 2 = true time of greatest obscur.

At this time,  $a = 12''.2$ ,  $\lambda' = 63''.7$ .

$$a = 12''.2 \quad - \quad \text{Ar. co. log. } 8.91364 \quad - \quad - \quad \text{log. } 1.08636$$

$$\lambda' = 63''.7 \quad - \quad - \quad \text{log. } 1.80414$$

---


$$\theta = 79^\circ 9' 30'' \quad - \quad \text{tan. } 0.71778 \quad - \quad \text{Ar. co. cos. } 0.72564$$

---


$$\Delta = 64''.9 \quad - \quad - \quad \text{log. } 1.81200$$

$$S + \Delta = 135''.8 - \text{log. } 2.13290, \quad \theta = 79^\circ 9' 30'' - \text{Ar. co. sin. } 0.00783$$

$$S - \Delta = 6''.2 - \text{log. } 0.79239, \quad k = 3628.9 \quad - \quad \text{log. } 3.55977$$

---


$$2) 2.92529, \quad h \quad - \quad - \quad \text{Ar. co. log. } 6.43240$$

---


$$1.46264 \quad - \quad - \quad - \quad - \quad 1.46264$$

$$L = 152^m. \quad - \quad - \quad - \quad \text{log. } 2.18184$$

$$\text{Const. log. } 1.77815$$

---


$$t = 0^h. \ 1^m. \ 11^s. \ 6 \quad - \quad \text{log. } 1.85503$$

$$\text{Time of greatest obscur.} \quad - \quad 4 \ 36 \ 25 \ .2$$

---


$$\text{Formation of ring,} \quad - \quad - \quad 4 \ 35 \ 13 \ .6, \text{ New-York time.}$$

---


$$\text{Rupture of} \quad \text{do.} \quad - \quad - \quad 4 \ 37 \ 36 \ .8 \quad \text{“} \quad \text{“}$$



## APPENDIX.

---

### TRIGONOMETRICAL FORMULÆ.

#### I. *Relative to a Single Arc or Angle $a$ .*

1.  $\sin^2 a + \cos^2 a = 1$
2.  $\sin a = \tan a \cos a$
3.  $\sin a = \frac{\tan a}{\sqrt{1 + \tan^2 a}}$
4.  $\cos a = \frac{1}{\sqrt{1 + \tan^2 a}}$
5.  $\tan a = \frac{\sin a}{\cos a}$
6.  $\cot a = \frac{1}{\tan a} = \frac{\cos a}{\sin a}$
7.  $\sin a = 2 \sin \frac{1}{2} a \cos \frac{1}{2} a$
8.  $\cos a = 1 - 2 \sin^2 \frac{1}{2} a$
9.  $\cos a = 2 \cos^2 \frac{1}{2} a - 1$
10.  $\tan \frac{1}{2} a = \frac{\sin a}{1 + \cos a}$
11.  $\cot \frac{1}{2} a = \frac{\sin a}{1 - \cos a}$
12.  $\tan^2 \frac{1}{2} a = \frac{1 - \cos a}{1 + \cos a}$
13.  $\sin 2a = 2 \sin a \cos a$
14.  $\cos 2a = 2 \cos^2 a - 1 = 1 - 2 \sin^2 a$

II. *Relative to Two Arcs  $a$  and  $b$ , of which  $a$  is supposed to be the greater.*

$$15. \quad \sin (a + b) = \sin a \cos b + \sin b \cos a$$

$$16. \quad \sin (a - b) = \sin a \cos b - \sin b \cos a$$

$$17. \quad \cos (a + b) = \cos a \cos b - \sin a \sin b$$

$$18. \quad \cos (a - b) = \cos a \cos b + \sin a \sin b$$

$$19. \quad \tan (a + b) = \frac{\tan a + \tan b}{1 - \tan a \tan b}$$

$$20. \quad \tan (a - b) = \frac{\tan a - \tan b}{1 + \tan a \tan b}$$

$$21. \quad \sin a + \sin b = 2 \sin \frac{1}{2} (a + b) \cos \frac{1}{2} (a - b)$$

$$22. \quad \sin a - \sin b = 2 \sin \frac{1}{2} (a - b) \cos \frac{1}{2} (a + b)$$

$$23. \quad \cos a + \cos b = 2 \cos \frac{1}{2} (a + b) \cos \frac{1}{2} (a - b)$$

$$24. \quad \cos b - \cos a = 2 \sin \frac{1}{2} (a + b) \sin \frac{1}{2} (a - b)$$

$$25. \quad \tan a + \tan b = \frac{\sin (a + b)}{\cos a \cos b}$$

$$26. \quad \tan a - \tan b = \frac{\sin (a - b)}{\cos a \cos b}$$

$$27. \quad \cot a + \cot b = \frac{\sin (a + b)}{\sin a \sin b}$$

$$28. \quad \cot b - \cot a = \frac{\sin (a - b)}{\sin a \sin b}$$

$$29. \quad \frac{\sin a + \sin b}{\sin a - \sin b} = \frac{\tan \frac{1}{2} (a + b)}{\tan \frac{1}{2} (a - b)}$$

$$30. \quad \frac{\cos b + \cos a}{\cos b - \cos a} = \frac{\cot \frac{1}{2} (a + b)}{\tan \frac{1}{2} (a - b)}$$

$$31. \quad \frac{\tan a + \tan b}{\tan a - \tan b} = \frac{\cot b + \cot a}{\cot b - \cot a} = \frac{\sin (a + b)}{\sin (a - b)}$$

$$32. \quad \frac{\cot b - \tan a}{\cot b + \tan a} = \frac{\cot a - \tan b}{\cot a + \tan b} = \frac{\cos (a + b)}{\cos (a - b)}$$

$$33. \quad \sin^2 a - \sin^2 b = \sin (a + b) \sin (a - b)$$

$$34. \quad \cos^2 a - \sin^2 b = \cos (a + b) \cos (a - b)$$

$$35. \quad 1 \pm \sin a = 2 \sin^2 (45^\circ \pm \frac{1}{2} a)$$

$$36. \quad \frac{1 \pm \sin a}{1 \mp \sin a} = \tan^2 (45^\circ \pm \frac{1}{2} a)$$



$$37. \quad \frac{1 \pm \sin a}{\cos a} = \tan (45^\circ \pm \tfrac{1}{2} a)$$

$$38. \quad \frac{1 - \sin a}{1 - \cos a} = \frac{\sin^2 (45^\circ - \tfrac{1}{2} a)}{\sin^2 \tfrac{1}{2} a}$$

$$39. \quad \frac{1 + \sin b}{1 + \cos a} = \frac{\sin^2 (45^\circ + \tfrac{1}{2} b)}{\cos^2 \tfrac{1}{2} a}$$

$$40. \quad \frac{1 + \tan b}{1 - \tan b} = \tan (45^\circ + b)$$

$$41. \quad \frac{1 - \tan b}{1 + \tan b} = \tan (45^\circ - b)$$

$$42. \quad \sin a \cos b = \tfrac{1}{2} \sin (a + b) + \tfrac{1}{2} \sin (a - b)$$

$$43. \quad \cos a \sin b = \tfrac{1}{2} \sin (a + b) - \tfrac{1}{2} \sin (a - b)$$

$$44. \quad \sin a \sin b = \tfrac{1}{2} \cos (a - b) - \tfrac{1}{2} \cos (a + b)$$

$$45. \quad \cos a \cos b = \tfrac{1}{2} \cos (a + b) + \tfrac{1}{2} \cos (a - b)$$

### III. *Trigonometrical Series.*

$$46. \quad \left\{ \begin{array}{l} \sin a = a - \frac{a^3}{2 \cdot 3} + \frac{a^5}{2 \cdot 3 \cdot 4 \cdot 5} - \&c. \\ \cos a = 1 - \frac{a^2}{2} + \frac{a^4}{2 \cdot 3 \cdot 4} - \frac{a^6}{2 \cdot 3 \cdot 4 \cdot 5 \cdot 6} + \&c. \\ \tan a = a + \frac{a^3}{3} + \frac{2a^5}{3 \cdot 5} + \frac{17a^7}{3^2 \cdot 5 \cdot 7} + \&c. \\ \cot a = \frac{1}{a} - \frac{a}{3} - \frac{a^3}{3^2 \cdot 5} - \frac{2a^5}{3^3 \cdot 5 \cdot 7} - \&c. \end{array} \right.$$

Let  $a$  = length of an arc of a circle of which the radius is 1, and  $(a'')$  = number of seconds in this arc, then to replace an arc expressed by its length, by the number of seconds contained in it, we have the formula,

$$47. \quad a = (a'') \sin 1''; \log. \sin 1'' = \overline{6.685574867}.$$

### IV. *Differences of Trigonometrical Lines.*

$$48. \quad \Delta \sin x = + 2 \sin \tfrac{1}{2} \Delta x. \cos (x + \tfrac{1}{2} \Delta x)$$

$$49. \quad \Delta \cos x = - 2 \sin \tfrac{1}{2} \Delta x. \sin (x + \tfrac{1}{2} \Delta x)$$

$$50. \quad \Delta \tan x = + \frac{\sin \Delta x}{\cos x. \cos (x + \Delta x)}$$

$$51. \quad \Delta \cot x = - \frac{\sin \Delta x}{\sin x \cdot \sin (x + \Delta x)}$$

V. Resolution of Right Angled Spherical Triangles.

<i>Given.</i>	<i>Required.</i>	<i>Solution.</i>
Hypothen. and an angle	side op. giv. ang. 52	$\sin x = \sin h \cdot \sin a$
	side adj. giv. ang. 53	$\tan x = \tan h \cdot \cos a$
	the other angle 54	$\cot x = \cos h \cdot \tan a$
Hypothen. and a side	the other side 55	$\cos x = \frac{\cos h}{\cos s}$
	ang. adj. giv. side 56	$\cos x = \tan s \cdot \cot h$
	ang. op. giv. side 57	$\sin x = \frac{\sin s}{\sin h}$
A side and the angle opposite	the hypothen. 58	$\sin x = \frac{\sin s}{\sin a}$
	the other side 59	$\sin x = \tan s \cdot \cot a$
	the other angle 60	$\sin x = \frac{\cos a}{\cos s}$
A side and the angle adjacent	the hypothen. 61	$\cot x = \cos a \cdot \cot s$
	the other side 62	$\tan x = \tan a \cdot \sin s$
	the other angle 63	$\cos x = \sin a \cdot \cos s$
The two sides	the hypothen. 64	$\cos x = \text{rectang. cos. of the giv. sides}$
	an angle 65	$\cot x = \sin \text{adj. side} \times \cot. \text{op. side}$
The two angles	the hypothen. 66	$\cos x = \text{rectang. cot. of the given angles}$
	a side 67	$\cos x = \frac{\cos. \text{opp. ang.}}{\sin. \text{adj. ang.}}$

the ambiguous cases

In these formulæ,  $x$  denotes the quantity sought.

$a$  = the *given* angle

$s$  = the *given* side

$h$  = the hypotenuse.

The formulæ for the resolution of right angled spherical triangles are all embraced in two rules discovered by Lord Napier, and called *Napier's Rules for the Circular Parts*. The circular parts, so called, are the two legs of the triangle, the complement of the hypotenuse and the complements of the acute angles. The right angle is omitted. In resolving a right angled spherical triangle, there are always three of the circular parts under consideration, namely, the two given parts and the required part. When the three parts in question are contiguous to each other, the middle one is called the *middle part*, and the others the *adjacent parts*. When two of them are contiguous, and the third is separated from these by a part on each side, the part thus separated is called the middle part, and the other two the *opposite parts*. The rules for the use of the circular parts are (the radius being taken = 1),

1. Sine of the middle part = the rectangle of the tangents of the adjacent parts.

2. Sine of the middle part = the rectangle of the cosines of the opposite parts.

Equations 52 to 67, are sufficient to resolve all the cases of right angled spherical triangles; but they lack precision if the unknown quantity is very small and determined by means of its cosine or cotangent; or, if the unknown quantity is near  $90^\circ$ , and given by a sine or a tangent: in these cases the following formulæ may be used,

$$68. \quad \tan^2 \frac{1}{2} a = - \frac{\cos (B + C)}{\cos (B - C)},$$

$$69. \quad \tan^2 \frac{1}{2} B = \frac{\sin (a - c)}{\sin (a + c)},$$

$$70. \quad \tan^2 \frac{1}{2} c = \tan \frac{1}{2} (a + b) \tan \frac{1}{2} (a - b),$$

$$71. \quad \tan (45^\circ - \frac{1}{2} b) = \sqrt{\tan (45^\circ - x)}, \tan x = \sin a \sin B.$$

$$72. \quad \tan^2 \frac{1}{2} b = \tan \left( \frac{B - C}{2} + 45^\circ \right) \tan \left( \frac{B + C}{2} - 45^\circ \right)$$

$a$  is the hypotenuse,  $B, C$ , the acute angles, and  $b, c$ , the sides opposite the acute angles.

# VI. Resolution of Oblique Angled Spherical Triangles.

If A, B, C, denote the three angles of a spherical triangle, and  $a$ ,  $b$ ,  $c$ , the sides which are opposite to them respectively.

$$73. \quad \frac{\sin A}{\sin a} = \frac{\sin B}{b} = \frac{\sin C}{\sin c}$$

or, the sines of the angles are proportional to the sines of the opposite sides.

$$74. \quad \cos c = \cos a \cos b + \sin a \sin b \cos C$$

$$75. \quad \cos c = \cos (a - b) - 2 \sin a \sin b \sin^2 \frac{1}{2} C$$

$$76. \quad \cos C = \sin A \sin B \cos c - \cos A \cos B$$

$$77. \quad \sin a \cos c = \sin c \cos a \cos B + \sin b \cos C$$

$$78. \quad \sin a \cot c = \cos a \cos B + \sin B \cot C$$

$$79. \quad \sin a \cos B = \sin c \cos b - \sin b \cos c \cos A$$

## I. Given the three sides, $a, b, c$ .

To find one of the angles.

$$80. \quad \sin^2 \frac{1}{2} A = \frac{\sin (k - b) \sin (k - c)}{\sin b \sin c}$$

$$81. \quad \cos^2 \frac{1}{2} A = \frac{\sin k \sin (k - a)}{\sin b \sin c}$$

$$82. \quad 2k = a + b + c$$

## II. Given the three angles A, B, C.

To find one of the sides.

$$83. \quad \sin^2 \frac{1}{2} a = \frac{\cos K \cos (K - A)}{\sin B \sin C}$$

$$84. \quad \cos^2 \frac{1}{2} a = \frac{\cos (K - B) \cos (K - C)}{\sin B \sin C}$$

$$85. \quad 2K = A + B + C.$$

## III. Given two sides $a$ and $b$ , and the included angle C.

1°. To find the two other angles A and B.

$$\left. \begin{aligned} 86. \quad \tan \frac{1}{2} (A + B) &= \cot \frac{1}{2} C. \frac{\cos \frac{1}{2} (a - b)}{\cos \frac{1}{2} (a + b)} \\ 87. \quad \tan \frac{1}{2} (A - B) &= \cot \frac{1}{2} C. \frac{\sin \frac{1}{2} (a - b)}{\sin \frac{1}{2} (a + b)} \end{aligned} \right\} \text{Napier's Analogies}$$

2°. To find the third side  $c$ .

$$88. \begin{cases} \tan \frac{1}{2} c = \tan \frac{1}{2} (a - b) \cdot \frac{\sin \frac{1}{2} (A + B)}{\sin \frac{1}{2} (A - B)} \\ \tan \frac{1}{2} c = \tan \frac{1}{2} (a + b) \cdot \frac{\cos \frac{1}{2} (A + B)}{\cos \frac{1}{2} (A - B)} \end{cases}$$

or equa. 74.

iv. *Given two angles A and B, and the adjacent side c.*

1°. To find the other two sides,  $a$  and  $b$ .

$$\begin{aligned} 80. \quad \tan \frac{1}{2} (a + b) &= \tan \frac{1}{2} c \cdot \frac{\cos \frac{1}{2} (A - B)}{\cos \frac{1}{2} (A + B)} \\ 90. \quad \tan \frac{1}{2} (a - b) &= \tan \frac{1}{2} c \cdot \frac{\sin \frac{1}{2} (A - B)}{\sin \frac{1}{2} (A + B)} \end{aligned} \quad \left. \vphantom{\begin{aligned} 80. \\ 90. \end{aligned}} \right\} \begin{array}{l} \text{Napier's} \\ \text{Analogies.} \end{array}$$

2°. To find the third angle  $C$ .

$$91. \begin{cases} \cot \frac{1}{2} C = \tan \frac{1}{2} (A - B) \cdot \frac{\sin \frac{1}{2} (a + b)}{\sin \frac{1}{2} (a - b)} \\ \cot \frac{1}{2} C = \tan \frac{1}{2} (A + B) \cdot \frac{\cos \frac{1}{2} (a + b)}{\cos \frac{1}{2} (a - b)} \end{cases}$$

or equa. 76.

v. *Given two sides  $a, b$ , and an opposite angle A.*

To find the other opposite angle  $B$ ; take equation 73, or the proportion; sines of the angles are as sines of the opposite sides. (For the methods of determining the remaining angle and side, see page 348, Case 3.)

vi. *Given two angles A, B, and an opposite side a.*

To find the other opposite side  $b$ ; sines of the angles are proportional to the sines of the opposite sides. (For the methods of determining the remaining side and angle, see page 349, Case 4.)

### *Other Methods of Resolving Oblique Angled Spherical Triangles.*

Except when three sides or three angles are given, the data always include an angle  $A$ , and the adjacent side  $b$ , besides a third part. The required parts in the different cases may be found by the following formulæ, and formula 73.

$$92. \quad \tan m = \tan b \cos A \qquad 93. \quad \cot n = \tan A \cos b$$



94.  $c = m + m'$

95.  $C = n + n'$

96.  $\frac{\cos a}{\cos b} = \frac{\cos m'}{\cos m}$

97.  $\frac{\cos A}{\cos B} = \frac{\sin n}{\sin n'}$

98.  $\frac{\tan A}{\tan B} = \frac{\sin m'}{\sin m}$

99.  $\frac{\tan a}{\tan b} = \frac{\cos n}{\cos n'}$

100.  $\sin k = \sin A \sin b.$

From the angle C, (Fig. 79), a perpendicular C D is let fall upon the opposite side  $c$ , which divides the triangle into two right angled triangles, that are resolved separately. In the one, A C D, A and  $b$  are known, and it is easy to find the other parts, which, joined to the third given part, serve to resolve the second right angled triangle B C D, and determine the unknown quantity required.  $m, m'$  denote the two segments of the base;  $n, n'$  the two parts of the angle C; and  $k$  the perpendicular arc C D.

It must be observed, that if the perpendicular C D fell without the triangle,  $m$  and  $m'$ ,  $n$  and  $n'$  would have contrary signs: this happens when the angles A and B at the base are of different kinds, (the one  $\angle$ , the other  $>90^\circ$ ). When it is not known whether this circumstance has place or not, the problem is susceptible of two solutions.

The perpendicular arc is to be let fall from that one of the vertices B or C, for which it does not divide into two parts the 3d element given with A and  $b$ .

The detail of the different cases is as follows: the data are A,  $b$ , and another arc or angle.

Case 1. *Given two sides and the included angle,  $b, c, A$ .*

Equation 92 makes known  $m$ , 94  $m'$ , which may be negative, (what the calculation shows), 96  $a$ , 98 B, and equation 73, (page 346), C, which is known in kind.

Case 2. *Given two angles and the adjacent side, A,  $c, b$ .*

Equation 93 makes known  $n$ , 95  $n'$ , which may be negative, (what the calculation shows), 97 B, 99  $a$ ; finally, equation 73 (page 346), gives C, which is known in kind.

Case 3. *Given two sides and an opposite angle,  $b, a, A$ .*

Equation 92 gives  $m$ , 96  $m'$ , 94  $c$ , 98 and 73 B and C; or else, 93 gives  $n$ , 99  $n'$ , 95 C, 97 and 73 B and  $c$ .

This problem admits in general of two solutions. In effect, the

arc  $m'$  or  $n'$  being given by its cos., may have either the sign + or —; there are then two values for  $c$ , and also for  $C$ .  $m'$  and  $n'$  enter into equations 97 and 98 by their sines, whence result two values of  $B$ ; same for  $C$  and  $c$ .

Case 4. *Given two angles and an opposite side, A, B, b.*

Equation 92 gives  $m$ , 98  $m'$ , 94  $c$ , 96  $a$ , and equation 73 makes known  $C$ ;

or else 93 gives  $n$ , 97  $n'$ , 95  $C$ , 99 and 73  $a$  and  $c$ .

There are also two solutions in this case; for,  $m'$  or  $n'$  is given by a sin., and therefore two supplementary arcs satisfy the question. Thus  $c$  in 94, or  $a$  in 99, receives two values; same for  $a$  in 96 and  $c$  in 95, &c.

When the triangle is *isocetes*,  $B = C$ ,  $b = c$ , the perpendicular arc must be let fall from the vertex  $A$ , and the equations become very simple. We find

$$101. \quad \sin \frac{1}{2} a = \sin \frac{1}{2} A \sin b$$

$$102. \quad \tan \frac{1}{2} a = \tan b \cos B$$

$$103. \quad \cos b = \cot B \cot \frac{1}{2} A$$

$$104. \quad \cot \frac{1}{2} A = \cos \frac{1}{2} a \sin B$$

The knowledge of two of the four elements  $A$ ,  $B$ ,  $a$ ,  $b$ , which form the *isocetes* triangle, is sufficient for the determination of the two others.

## INVESTIGATION OF ASTRONOMICAL FORMULÆ.

*Formulæ for the Parallax in Right Ascension and Declination, and in Longitude and Latitude.* (Referred to from Article 106, page 47).

Let  $s$  (Fig. 80) be the *true place* of a star seen from the centre of the earth,  $s'$  the *apparent place*, seen from a point on the surface of which  $z$  is the zenith, the latitude being  $l$ . The displacement  $s s' = p$  is the parallax in altitude, which takes effect in the vertical circle  $z s'$ ;  $p$  is the pole; the hour angle  $z p s = q$  is changed into  $z p s'$ , and  $s p s' = a$  is the *variation of the hour angle, or the parallax in right ascension*; the polar distance

$p s = d$  is changed into  $p s'$ ; the difference  $\delta$  of these arcs is the *parallax in declination* or of *polar distance*. We have (For. 73, p. 346),

$$\begin{aligned} \sin s' : \sin p s (d) &:: \sin s p s' (\alpha) : \sin s s' (p), \\ \sin s p s' (q + \alpha) : \sin z s' (Z) &:: \sin s' : \sin p z (90^\circ - l). \end{aligned}$$

Multiplying, term by term, we obtain,

$$\sin s' \sin (q + \alpha) : \sin d \sin Z :: \sin \alpha \sin s' : \sin p \cos l;$$

$$\text{whence,} \quad \sin \alpha = \frac{\sin p \cos l}{\sin d \sin Z} \sin (q + \alpha)$$

Or, substituting for  $p$  its value given by equ. (10,) p. 43, and replacing  $H$  by  $P$ ,

$$\sin \alpha = \frac{\sin P \cos l}{\sin d} \sin (q + \alpha). \dots (A).$$

This equation makes known  $\alpha$  when the apparent hour angle  $z p s' = q + \alpha$ , seen from the earth's surface, is given; but if we know the true hour angle  $z p s = q$ , seen from the centre of the earth, developing  $\sin (q + \alpha)$ , (For. 15, p. 342), and putting  $\frac{\sin P \cos l}{\sin d} = m$

$$\sin \alpha = m (\sin q \cos \alpha + \sin \alpha \cos q),$$

or, dividing by  $\sin \alpha$ ,

$$1 = m (\sin q \cot \alpha + \cos q);$$

whence, by transformation,

$$\tan \alpha = \frac{m \sin q}{1 - m \cos q} = m \sin q + m^2 \sin q \cos q.$$

Restoring the value of  $m$ ,

$$\tan \alpha = \frac{\sin P \cos l}{\sin d} \sin q + \left( \frac{\sin P \cos l}{\sin d} \right)^2 \sin q \cos q.$$

Putting the arc  $\alpha$  in place of its tangent, and  $P$  in place of  $\sin P$ , and expressing these arcs in seconds, (For. 47, p. 343), there results,

$$\alpha = \frac{P \cos l}{\sin d} \sin q + \left( \frac{P \cos l}{\sin d} \right)^2 \sin q \cos q \sin 1'' \dots (B).$$

The *parallax in declination*  $\delta$  is the difference of the arcs  $p s = d$ ,  $p s' = d + \delta$ . Let  $z s = z$ , and  $z s' = Z$ . The triangles  $z p s$  and  $z p s'$  give (For. 74 and 73),

$$1^{\circ}. \cos p \approx s = \frac{\cos d - \sin l \cos z}{\cos l \sin z} = \frac{\cos (d + \delta) - \sin l \cos Z}{\cos l \sin Z},$$

$$2^{\circ}. \sin p \approx s = \frac{\sin d \sin q}{\sin z} = \frac{\sin (d + \delta) \sin (q + \alpha)}{\sin Z}.$$

From the first equation we derive,

$$\begin{aligned} \cos (d + \delta) &= \frac{\cos d \sin Z - \sin l \cos z \sin Z}{\sin z} + \sin l \cos Z \\ &= \frac{\cos d \sin Z - \sin l (\cos z \sin Z - \sin z \cos Z)}{\sin z} \\ &= \frac{\cos d \sin Z - \sin l \sin (Z - z)}{\sin z}, \end{aligned}$$

or, (equ. 10, p. 43),

$$= \frac{\sin Z}{\sin z} (\cos d - \sin P \sin l);$$

from the second,

$$\frac{\sin Z}{\sin z} = \frac{\sin (d + \delta)}{\sin d} \cdot \frac{\sin (q + \alpha)}{\sin q};$$

substituting,

$$\cos (d + \delta) = \frac{\sin (d + \delta)}{\sin d} \cdot \frac{\sin (q + \alpha)}{\sin q} (\cos d - \sin P \sin l)$$

$$\frac{\cos (d + \delta)}{\sin (d + \delta)} = \frac{\sin (q + \alpha)}{\sin q} \left( \frac{\cos d}{\sin d} - \frac{\sin P \sin l}{\sin d} \right)$$

$$\cot (d + \delta) = \frac{\sin (q + \alpha)}{\sin q} \left( \cot d - \frac{\sin P \sin l}{\sin d} \right). \quad (C)$$

$$\text{Put } \tan x = -\frac{\sin P \sin l}{\sin d};$$

$$\begin{aligned} \text{then, } \cot (d + \delta) &= \frac{\sin (q + \alpha)}{\sin q} (\cot d - \tan x) \\ &= \frac{\sin (q + \alpha)}{\sin q} \left( \frac{\cos d}{\sin d} - \frac{\sin x}{\cos x} \right) \\ &= \frac{\sin (q + \alpha)}{\sin q} \cdot \frac{\cos d \cos x - \sin d \sin x}{\sin d \cos x} \\ &= \frac{\sin (q + \alpha) \cos (d + x)}{\sin q \sin d \cos x} \dots (D). \end{aligned}$$

The apparent polar distance  $(d + \delta)$  being computed by either of the formulæ (C) and (D), we have  $\delta = (d + \delta) - d$ .

Formulæ may be obtained that will give the parallax in latitude without first finding the apparent latitude (except approximately.)

From equa. (C) we obtain,

$$\frac{\sin P \sin l}{\sin d} = \cot d - \frac{\sin q \cot (d + \delta)}{\sin (q + \alpha)},$$

and we also have,

$$\cot d - \cot (d + \delta) = \frac{\cos d}{\sin d} - \frac{\cos (d + \delta)}{\sin (d + \delta)} = \frac{\sin \delta}{\sin d \sin (d + \delta)};$$

the sum of these equations gives

$$\frac{\sin P \sin l}{\sin d} = \cot (d + \delta) \left( 1 - \frac{\sin q}{\sin (q + \alpha)} \right) + \frac{\sin \delta}{\sin d \sin (d + \delta)}.$$

$$\begin{aligned} \text{Now,} \quad 1 - \frac{\sin q}{\sin (q + \alpha)} &= \frac{\sin (q + \alpha) - \sin q}{\sin (q + \alpha)} \\ &= \frac{2 \sin \frac{1}{2} \alpha \cos (q + \frac{1}{2} \alpha)}{\sin (q + \alpha)} = \frac{\sin \alpha \cos (q + \frac{1}{2} \alpha)}{\sin (q + \alpha) \cos \frac{1}{2} \alpha} \quad (\text{For. 22, 13.}) \\ &= \frac{\cos (q + \frac{1}{2} \alpha) \sin P \cos l}{\sin d \cos \frac{1}{2} \alpha}, \text{ by equa. (A).} \end{aligned}$$

Substituting,

$$\begin{aligned} \frac{\sin P \sin l}{\sin d} &= \cot (d + \delta) \frac{\cos (q + \frac{1}{2} \alpha) \sin P \cos l}{\sin d \cos \frac{1}{2} \alpha} + \\ &\quad \frac{\sin \delta}{\sin d \sin (d + \delta)}, \end{aligned}$$

$$\begin{aligned} \text{or,} \quad \sin \delta &= \sin P \sin l \sin (d + \delta) - \\ &\quad \frac{\cos (d + \delta) \cos (q + \frac{1}{2} \alpha) \sin P \cos l}{\cos \frac{1}{2} \alpha} \dots (\text{E}). \end{aligned}$$

$$= \sin P \sin l [\sin (d + \delta) - \tan. y \cos (d + \delta)],$$

$$\text{making} \quad \tan y = \frac{\cot l \cos (q + \frac{1}{2} \alpha)}{\cos \frac{1}{2} \alpha};$$

$$\text{whence,} \quad \sin \delta = \frac{\sin P \sin l}{\cos y} \sin (d + \delta - y) \dots (\text{F})$$

To facilitate the calculation, the sines of  $\delta$  and  $P$  in eqs. (E) and (F), may be replaced by the arcs.



To obtain an expression for the parallax in declination in terms of the *true declination*, developpe  $\sin (d + \delta - y)$  in equation (F), which gives,

$$\sin \delta = \frac{\sin P \sin l}{\cos y} [\sin (d + \delta) \cos y - \sin y \cos (d + \delta)];$$

developping  $\sin (d + \delta)$  and  $\cos (d + \delta)$  and reducing, we have,

$$\sin \delta = \frac{\sin P \sin l}{\cos y} [\sin (d - y) \cos \delta + \cos (d - y) \sin \delta],$$

dividing by  $\cos \delta$ ,

$$\tan \delta = \frac{\sin P \sin l}{\cos y} [\sin (d - y) + \cos (d - y) \tan \delta],$$

$$\text{whence } \tan \delta = \frac{\frac{\sin P \sin l}{\cos y} \cdot \sin (d - y)}{1 - \frac{\sin P \sin l}{\cos y} \cdot \cos (d - y)}$$

$$= \frac{\sin P \sin l}{\cos y} \sin (d - y) + \left( \frac{\sin P \sin l}{\cos y} \right)^2 \sin (d - y) \cos (d - y);$$

or, replacing  $\tan \delta$  and  $\sin P$ , by  $\delta$  and  $P$ ; expressing these arcs in seconds, (For. 47, p. 343), and reducing by For. 13, p. 341,

$$\delta = \frac{P \sin l}{\cos y} \sin (d - y) + \left( \frac{P \sin l}{\cos y} \right)^2 \frac{\sin 1''}{2} \cdot \sin 2 (d - y). (G)$$

If the place of a body be referred to the ecliptic, similar formulæ will give the *parallax in longitude and latitude*, but as the ecliptic and its pole are continually in motion by virtue of the diurnal rotation of the heavens, it is necessary, in order to be able to determine the parallax in longitude at any given instant, to know the situation of the ecliptic at the same instant.

This is ascertained by finding the situation of the point of the ecliptic  $90^\circ$  distant from the points in which it cuts the horizon, and which are respectively just rising and setting, called the *Nonagesimal Degree*, or the *Nonagesimal*.

Let K (Fig. 81) be the pole of the ecliptic  $f b$ ,  $p$  the pole of the equator  $f a$ ;  $f$  is the vernal equinox, the origin of longitudes and of right ascensions;  $h b s$  is the eastern horizon,  $b$  the *horoscope*, or the point of the ecliptic which is just rising;  $p z = 90^\circ - l$  (the

latitude of given place);  $K p = \omega$  the obliquity of the ecliptic. The circle  $K z n v$  is at the same time perpendicular at  $n$  to the ecliptic  $f b$ , and at  $v$  to the horizon  $h b$ : it is a circle of latitude and a vertical circle, since it passes through the pole  $K$  and the zenith  $z$ ;  $b$  is  $90^\circ$  from all the points of the circle  $K n v$ ;  $z n$  is the latitude of the zenith,  $f n$  its longitude; the point  $n$  is the nonagesimal, since  $b n = 90^\circ$ ;  $n v$  is the altitude of this point, and the complement of  $z n$ ;  $n v$  measures the inclination of the ecliptic to the horizon at the given instant, or the angle  $b$ , so that  $b = n v = K z$ ; thus  $f n = N$  the longitude of the nonagesimal, and  $n v = h$  the altitude of the nonagesimal, designate the situation of this point, and consequently ascertain the position of the ecliptic and its pole at the moment of observation.

The points  $m$  and  $d$  are those of the equator and ecliptic which are on the meridian; the arc  $f m$ , in time, is the sidereal time  $s$ , which is known; the arc  $f i = 90^\circ$ , since the plane  $K p i$ , passing through the poles  $K$  and  $p$ , is at the same time perpendicular to the ecliptic and to the equator; the arc  $m i = f i - f m = 90^\circ - s$ ; then the angle  $z p K = 180^\circ - z p i = 180^\circ - m i = 90^\circ + s$ .

Now, in the spherical triangle  $p K z$  we know the sides  $K p = \omega$ ,  $z p = 90^\circ - l = H$ , and the included angle  $z p K = 90^\circ + s$ ; and may therefore find  $K z = h$  the altitude of the nonagesimal, and the angle  $p K z = n c = f c - f n = 90^\circ - N$ , complement of the longitude  $N$  of the nonagesimal. Let  $S$  = sum of the angles  $K z p$  and  $z K p$ , then (For. 86, page 346),

$$\tan \frac{1}{2} S = \frac{\cos \frac{1}{2} (H - \omega)}{\cos \frac{1}{2} (H + \omega)} \cdot \cot \frac{1}{2} (90^\circ + s),$$

$$\text{or,} \quad \tan \frac{1}{2} S = \frac{\cos \frac{1}{2} (H - \omega)}{\cos \frac{1}{2} (H + \omega)} \cdot \tan \frac{1}{2} (90^\circ - s);$$

$$\tan \frac{1}{2} S = - \tan (180^\circ - \frac{1}{2} S), \quad \tan \frac{1}{2} (90^\circ - s) = - \tan (s - 90^\circ);$$

substituting, and denoting  $(180^\circ - \frac{1}{2} S)$  by  $E$ , we have,

$$\tan E = \frac{\cos \frac{1}{2} (H - \omega)}{\cos \frac{1}{2} (H + \omega)} \cdot \tan \frac{1}{2} (s - 90^\circ) \dots (H).$$

Again, let  $D = Z K p - K z p$ , then (For. 87),

$$\tan \frac{1}{2} D = \frac{\sin \frac{1}{2} (H - \omega)}{\sin \frac{1}{2} (H + \omega)} \cdot \cot \frac{1}{2} (90^\circ + s);$$

whence, by transforming as above, and denoting  $(180^\circ - \frac{1}{2} D)$  by  $F$ , we have,

$$\tan F = \frac{\sin \frac{1}{2} (H - \omega)}{\sin \frac{1}{2} (H + \omega)} \cdot \tan \frac{1}{2} (s - 90^\circ) \dots (I).$$

Now,  $\frac{1}{2} S + \frac{1}{2} D = p K z = 90^\circ - N$ ;

whence,  $N = 90^\circ - (\frac{1}{2} S + \frac{1}{2} D)$ ,

or,

$$N = 360^\circ + 90^\circ - (\frac{1}{2} S + \frac{1}{2} D) = 180^\circ - \frac{1}{2} S + 180^\circ - \frac{1}{2} D + 90^\circ,$$

consequently,  $N = E + F + 90^\circ \dots (J)$ ,

rejecting  $360^\circ$ , when the sum exceeds that number.

Next, for the altitude of the nonagesimal we have, (For. 88),

$$\begin{aligned} \tan \frac{1}{2} h &= \frac{\cos \frac{1}{2} S}{\cos \frac{1}{2} D} \cdot \tan \frac{1}{2} (H + \omega), \\ &= \frac{\cos E}{\cos F} \cdot \tan \frac{1}{2} (H + \omega) \dots (K). \end{aligned}$$

$N$  and  $h$  being known, to obtain the *formulæ for the parallax in longitude and latitude*, we have only to replace in the formulæ for the parallax in right ascension and declination, the altitude  $l$  of the pole of the equator, by that  $90^\circ - h$  of the pole  $K$  of the ecliptic, and the distance  $im$  of the star  $s$  from the meridian by the distance  $nc$  to the vertical through the nonagesimal. Let us change then in formulæ (A), (B), (C), (D), (E), (F), and (G),  $l$  into  $90^\circ - h$ , and  $q$  into  $fc - fn = L - N$ ,  $L$  being the longitude  $fc$  of the star  $s$ . Besides  $d$  will become the distance  $s$   $K$  to the pole of the ecliptic, complement of the latitude  $\lambda = sc$ . Making these substitutions, and denoting the parallax in longitude by  $\Pi$ , and the parallax in latitude by  $\pi$ , we obtain in terms of the apparent longitude and latitude,

$$\sin \Pi = \frac{\sin P \sin h}{\sin d} \cdot \sin (L - N + \Pi) \dots (L),$$

$$\cot (d + \pi) = \frac{\sin (L - N + \Pi)}{\sin (L - N)} \left( \cot d - \frac{\sin P \cos h}{\sin d} \right) \dots (M),$$

$$\tan x = \frac{\sin P \cos h}{\sin d} \dots (N),$$

$$\cot (d + \pi) = \frac{\sin (L - N + \Pi) \cos (d + x)}{\sin (L - N) \sin d \cos x} \dots (O),$$

$$\sin \pi = \sin P \cos h \sin (d + \pi) - \frac{\cos (d + \pi) \cos (L - N + \frac{1}{2} \Pi) \sin P \sin h}{\cos \frac{1}{2} \Pi} \dots (P),$$

$$\tan y = \frac{\tan h \cos (L - N + \frac{1}{2} \Pi)}{\cos \frac{1}{2} \Pi} \dots (Q),$$

$$\sin \pi = \frac{\sin P \cos h}{\cos y} \sin (d + \pi - y) \dots (R),$$

and in terms of the true longitude and latitude,

$$\Pi = \frac{P \sin h}{\sin d} \cdot \sin (L - N) + \left( \frac{P \sin h}{\sin d} \right)^2$$

$$\sin (L - N) \cos (L - N) \sin 1'' \dots (S),$$

$$\pi = \frac{P \cos h}{\cos y} \cdot \sin (d - y) + \frac{1}{2} \left( \frac{P \cos h}{\cos y} \right)^2$$

$$\sin 2 (d - y) \sin 1'' \dots (T),$$

$$\tan y = \frac{\tan h \cos (L - N + \frac{1}{2} \Pi)}{\cos \frac{1}{2} \Pi}.$$

To facilitate the computation,  $\sin \Pi$ ,  $\sin \pi$ , and  $\sin P$ , in formulæ (L), (P), and (R), may be replaced by the arcs themselves.

The distance  $d$  from the pole of the ecliptic enters into these formulæ in place of the latitude  $\lambda$ .

To find the apparent distance  $d'$  we have,

$$d' = d + \pi;$$

for the apparent latitude  $\lambda'$ ,

$$\lambda' = \lambda - \pi;$$

for the apparent longitude  $L'$ ,

$$L' = L + \Pi.$$

The logarithmic formulæ given on page 292, were derived from equations (L), (O), and (P), and the logarithmic formula on page 294 from equa. (O).

To determine now the effect of parallax upon the apparent diameter of the moon.

Let  $ACB$ , (Fig. 51) represent the moon, and  $E$  the station of an observer; also let  $R$  = apparent semi-diameter of the moon, and  $D$  = its distance. The triangle  $AES$  gives

$$\sin AES = \frac{AS}{ES}, \text{ or, } \sin R = \frac{AS}{D}.$$

At any other distance  $D'$  we should have for the apparent semi-diameter  $R'$ ,

$$\sin R' = \frac{AS}{D'},$$

whence, 
$$\frac{\sin R'}{\sin R} = \frac{D}{D'}$$

Thus, if  $R'$  = moon's apparent semi-diameter to an observer at the earth's surface, as at  $O$  Fig. (20),  $R$  = the same as it would be seen from the centre  $C$ , and  $S$  represents the situation of the moon,

$$\frac{\sin R'}{\sin R} = \frac{CS}{OS} = \frac{\sin ZOS}{\sin ZCS} = \frac{\sin Z}{\sin z}$$

But we have, (see page 351.)

$$\frac{\sin Z}{\sin z} = \frac{\sin (d + \delta)}{\sin d} \cdot \frac{\sin (q + a)}{\sin q}$$

or, in terms of the apparent longitude and latitude, (see page 355),

$$\frac{\sin Z}{\sin z} = \frac{\sin (d + \pi)}{\sin d} \cdot \frac{\sin (L - N + \Pi)}{\sin L - N}.$$

Hence, 
$$\sin R' = \frac{\sin R \sin (d + \pi) \sin (L - N + \Pi)}{\sin d \sin (L - N)} \dots (U).$$

*Aberration in Longitude and Latitude, and in Right Ascension and Declination.* (Referred to from Art. 114, page 51.)

Aberration is caused by the motion of light in conjunction with the motion of the earth. Light comes to us from the sun in  $8' 13''.2$ , during which time the earth describes an arc  $a = 20''.36$ , of its orbit  $pbd$  in (Fig. S2,) supposed circular:  $p$  is the place of the earth. Let us take any plane whatsoever, which we will



call *relative*, passing through the star, and let  $d d'$  be the intersection of this plane and the ecliptic, with which it makes an angle  $k$ : let us seek the quantity  $\varphi$  by which the aberration displaces the star in the direction perpendicular to this plane. The question is to project perpendicularly to the relative plane, the small constant arc  $a$  which the earth describes, this being the quantity that the star is displaced from its line of direction, (which lies in the relative plane,) in a direction parallel to the line of the earth's motion, (see Art. 109 of the text): this projection is  $\varphi$ , variable according to the position of the *relative plane* in relation to which it is estimated. The velocity along the tangent at  $p$ , makes with  $p h$  an angle  $\theta = p c h =$  the arc  $p d'$ ;  $a \cos \theta$  is then the projection of this velocity on the line  $p h$ . The angle of our two planes being  $k$ , this projection will be reduced to  $a \cos \theta \sin k$ , when it is taken perpendicularly to the relative plane. Thus,

$$\varphi = a \sin k \cos \theta. \dots (V).$$

The aberration displaces the star from the relative plane by this quantity  $\varphi$ ,  $k$  designating the inclination of this plane to the ecliptic, and  $\theta$  the arc  $p d'$ , reckoned from  $p$  the place of the earth to  $d'$  the point of intersection of these two planes. Let us give to the relative plane the positions which are met with in applications.

Let us suppose at first that  $k = 90^\circ$ , or  $\sin k = 1$ ; the relative plane will then be perpendicular to the ecliptic. Let  $n$  be the vernal equinox; we have  $p d' = n p - n d'$ ;  $n p$  is the longitude of the earth, or  $180^\circ +$  that  $\odot$  of the sun;  $n d'$  is the longitude  $l$  of the star; whence

$$\varphi = -a \cos (\odot - l).$$

Now, let  $M$ , (Fig. 83), be the true place of the star,  $M'$  the star as displaced by aberration,  $K M$  is the circle of true latitude,  $K M'$  the circle of apparent latitude, and  $M M' = \varphi$ : this arc has its centre  $C$  on the axis which passes through the pole  $K$  of the ecliptic; the longitude of the star is then altered by the part  $O O'$  of the ecliptic comprised between these two planes; and since  $O O'$  is to the arc  $M M'$  as the radius 1 is to the radius  $C M = \sin K M = \cos$  latitude  $\lambda$  of the star, we have

$$\text{aberr. in long.} = -\frac{a}{\cos \lambda} \cos (\odot - l) \dots (W).$$

If the relative plane is  $k c$  (Fig. 84), perpendicular to the circle

of latitude  $K c d$ , the aberration  $\varphi$  perpendicularly to it, will be the aberration in latitude. Let  $k d$  be the ecliptic, and  $o$  the earth; the angle  $k$  is measured by the arc  $c d = \lambda$ ; the arc  $o k = \theta = \odot$  — long. of  $k$ ; and as  $k d = 90^\circ$ , long. of point  $k = l - 90^\circ$ ; substituting in equation (V) we find,

$$\text{aberr. in lat.} = -a \sin \lambda \sin (\odot - l) \dots (X).$$

These aberrations of the star produce a small apparent orbit, which is confounded with its projection on the tangent plane to the celestial sphere. Let us suppose the orbit to be referred to two co-ordinate axes passing through the true place of the star and lying in the tangent plane, of which one is parallel to the plane of the ecliptic, and the other perpendicular to this, or tangent to the circle of latitude at the star; and let  $\frac{x}{\cos \lambda} = \text{aberr.}$

in long., and  $y = \text{aberr. in lat.}$ ;  $y$  will be the ordinate, and  $x$  (the aberr. in long., reduced to the parallel through the star) the abscissa; we have,

$$\frac{x}{\cos \lambda} = -\frac{a}{\cos \lambda} \cdot \cos (\odot - l),$$

$$y = -a \sin \lambda \sin (\odot - l),$$

or, 
$$\frac{x}{a} = -\cos (\odot - l),$$

$$\frac{y}{a \sin \lambda} = -\sin (\odot - l).$$

Squaring the last two equations, and adding them together,  $\odot$  disappears, and we find

$$y^2 + x^2 \sin^2 \lambda = a^2 \sin^2 \lambda \dots (Y).$$

Whatever may be the place of the earth, such is the equation of the *apparent orbit*, which, as we perceive, is an ellipse of which the semi-axes are  $a$  and  $a \sin \lambda$ , and whose centre is the true place of the star. When the star is at the pole of the ecliptic,  $\lambda = 90^\circ$ , and the ellipse becomes a circle of which the radius is  $a$ . When  $\lambda = 0$ , this ellipse is reduced to an arc  $2 a$  of the ecliptic.

To find the aberration in right ascension, the relative plane must be perpendicular to the equator. Let  $k c$  be the equator (Fig. 84),  $p$  its pole,  $p s d$  the relative plane, which is the circle of declination of the star  $s$ ;  $k d$  the ecliptic,  $o$  the earth,  $k$  the

vernal equinox,  $k c = R$ ,  $s c = D$ . Aberration carries the star  $s$  out of the plane  $p c d$  a distance  $\varphi$ , which it is the question to determine. Equa. (V) is here

$$\begin{aligned}\varphi &= a \sin d \cos d o = a \sin d \cos (k d - k o) \\ &= a \sin d (\cos k d \cos k o + \sin k d \sin k o) \\ &= a \sin d \cos k d \cos k o + a \sin d \sin k d \sin k o,\end{aligned}$$

but  $k o = \text{long. of earth} = 180^\circ + \odot$ ; we have also the angle  $k$  = the obliquity  $\omega$  of the ecliptic, and the right angled spherical triangle  $k c d$  gives by Napier's rules,

$$\cot k d = \cot R \cos \omega, \quad \sin d \sin k d = \sin R.$$

The 1st equa. multiplied by the 2d, gives

$$\sin d \cos k d = \cos R \cos \omega,$$

whence,  $\varphi = -a (\cos R \cos \omega \cos \odot + \sin R \sin \odot)$ .

The displacement of  $M$  to  $M'$  (Fig. S3,) conducts, as before, to the division of  $\varphi$  by  $\cos D$ , to have the corresponding arc of the equator: thus the *aberration in right ascension* is,

$$d R = -a \sin R \sec D \sin \odot - a \cos \omega \cos R \sec D \cos \odot \text{ (Z)}.$$

Taking the relative plane perpendicular to the circle of declination, we find for the *aberration in declination*,

$$d D = -a \sin D \cos R \sin \odot - a \cos \omega (\tan. \omega \cos D. - \sin R \sin D) \cos \odot \dots (a).$$

These formulæ may easily be adapted to logarithmic computation:

In formula (Z) let  $a \sin R \sec D = A$ , and  $a \cos \omega \cos R \sec D = B$ ; then,

$$d R = -A (\sin \odot + \frac{B}{A} \cos \odot) \dots (Z').$$

$$\text{Put } \tan \varphi = \frac{B}{A} = \frac{a \cos \omega \cos R \sec D}{a \sin R \sec D} = \cos \omega \cot R \dots (b)$$

and we shall have

$$\begin{aligned}d R &= -A \left( \sin \odot + \frac{\sin \varphi}{\cos \varphi} \cos \odot \right) \\ &= -A \cdot \frac{\sin \odot \cos \varphi + \sin \varphi \cos \odot}{\cos \varphi}\end{aligned}$$

$$= -\frac{A}{\cos \varphi} \cdot \sin (\odot + \varphi).$$

Restoring the value of  $A$ , and taking  $\frac{1}{\cos D}$  for  $\sec D$ , we obtain,

$$dR = -\frac{a \sin R}{\cos D \cos \varphi} \cdot \sin (\odot + \varphi) \dots (c).$$

The auxiliary arc  $\varphi$  is given by equation (b); it must be substituted in equation (c), with its sign, and we then obtain  $dR \cdot \tan \varphi$ , and the co-efficient of  $\sin (\odot + \varphi)$  are constant, for the same star, for a long period of time, since these quantities vary very slowly with  $\omega$  and the precession. Moreover, the co-efficient of  $\sin (\odot + \varphi)$  is the maximum value of  $dR$ , since it answers to  $\sin (\odot + \varphi) = 1$ . Thus we shall be able to calculate in advance, for any designated star, the values of  $\varphi$  and of the *maximum* of the aberration in right ascension, or of the logarithm of this maximum.

The results of these calculations for 50 principal stars are given in Table XCI, columns headed  $M$  and  $\varphi$ .

If in equation (a), we make  $a \sin D \cos R = A'$ , and  $a \cos \omega (\tan \omega \cos D - \sin R \sin D) = B'$ , we shall have the equation,

$$dD = -A' (\sin \odot + \frac{B'}{A'} \cos \odot),$$

in which  $A'$  and  $B'$  are constants. This equation is of the same form with equa. (Z'). We therefore have in the same manner as for the right ascension,

$$\begin{aligned} \tan \theta &= \frac{B'}{A'} = \frac{a \cos \omega (\tan \omega \cos D - \sin R \sin D)}{a \sin D \cos R} = \\ &= \frac{a \sin \omega \cos D - a \cos \omega \sin R \sin D}{a \sin D \cos R} = \\ &= \frac{\sin \omega \cot D}{\cos R} - \cos \omega \tan R \dots (d), \end{aligned}$$

$$dD = -\frac{A'}{\cos \theta} \cdot \sin (\odot + \theta) = -\frac{a \sin D \cos R}{\cos \theta} \times \sin (\odot + \theta) \dots (e).$$

$\theta$  is given by equation (d), and being substituted in equation (e), we shall have  $dD$ .  $\theta$  and the co-efficient of  $\sin (\odot + \theta)$

are constant for the same star, and we can therefore calculate in advance the values of this  $\varphi$ , and of the co-efficient, which is the *maximum* of the aberration in declination. Columns headed  $\theta$  and N, Table XCI, contain the quantities  $\theta$  and the logarithms of the maxima of the aberration in declination for 50 principal stars.

For convenience in calculation, the angles  $\varphi$ ,  $\theta$ , and the maxima, M, N, in Table XCI, have been rendered positive in all cases. This has been accomplished by adding  $12^s$  to  $\varphi$  and  $\theta$  whenever the calculation conducted to a negative value, and by adding  $6^s$  to  $\odot + \varphi$ , or  $\odot + \theta$ , whenever the co-efficient had the sign — ; in this manner the sign of the two factors is changed, which does not alter the sign of the product.

*Formulae for the Nutation in Right Ascension and Declination.* (Referred to from Article 132, p. 56).

In deriving these formulæ, we must begin with borrowing certain results established by Physical Astronomy. It has been proved, in confirmation of Bradley's conjectures, that the phenomena of nutation are explicable on the hypothesis of the pole of the earth, describing, round its mean place, (that place which, see p. 53, it would hold in the small circle described round the pole of the ecliptic, were there no *inequality* of precession) an ellipse, in a period equal to the revolution of the moon's nodes. The major axis of this ellipse is situated in the solstitial colure and equal to  $18''.50$  ; it bears that proportion to the minor axis (such are the results of theory) which the cosine of the obliquity bears to the cosine of twice the obliquity : consequently, the minor axis will be  $13''.77$ .

Let  $C d A$  represent such an ellipse,  $P$  being the mean place of the pole,  $K$  the pole of the ecliptic.  $C D A O$  is a circle described with the centre  $P$  and radius  $C P$ .  $\varpi L$  is the ecliptic,  $\varpi w$  the equator,  $K P L$  the solstitial colure. In order to determine the true place of the pole, take the angle  $A P O$  equal to the retrogradation of the moon's ascending node from  $\varpi$  : draw  $O i$  perpendicular to  $P A$ , and the point in the ellipse, through which  $O i$  passes, is the true place of the pole. This construction being admitted, the *nutations* in right ascension and north



polar distance may,  $P p$  being very small, be thus easily computed.

*Nutation in North Polar Distance.*

$$\begin{aligned} \text{Nutation in } N D P &= P \sigma - p \sigma. = P r = P p. \cos p P \sigma, \text{ nearly,} \\ &= P p \cos (A P p + A P \sigma) \\ &= P p \cos (A P p + R - 90^\circ) \\ &= P p \sin (A P p + R) \end{aligned}$$

$R$  denoting the right ascension.

*Nutation in Right Ascension.*

The right ascension of a star is, by the effect of nutation, changed from  $\varphi w$  into  $\varphi' t s$ . Now,

$$\varphi' t s = \varphi' v + \varphi w + t s, \text{ nearly,}$$

$$\text{whence,} \quad \varphi w - \varphi' t s = - \varphi' v - t s$$

$$= - \varphi \varphi' \cos \varphi \varphi' v - P p \sin p P \sigma \cdot \frac{\sin \sigma s}{\sin P \sigma},$$

in which expression  $\varphi' v (= \varphi \varphi' \cos \varphi \varphi' v)$  is, as in the case of precession, common to all stars.

In order to reduce farther the above expression, we have

$$p P \sigma = A P p + A P \sigma = A P p + R - 90^\circ,$$

$$\text{and } \varphi \varphi' = L l = P p \cdot \frac{\sin A P p}{\sin P K};$$

$$\text{whence,} \quad - \varphi' v - t s = - P p \sin A P p \cotan \omega$$

$$- P p \sin (A P p + R - 90^\circ), \cot N. P. D$$

$$= - P p \sin A P p \cot \omega + P p \cos (A P p + R) \cot \delta,$$

$\delta$  representing the north polar distance, and  $\omega$  the obliquity of the ecliptic.

But these forms are not convenient for computation. In order to render them convenient, we must, from the properties of the ellipse, deduce the values of  $P p$ , and of the tangent of  $A P p$ , and then substitute such values in the above expressions: thus,

$$\frac{P p}{P O} = \frac{\sec A P p}{\sec A P O} = \frac{\cos A P O}{\cos A P p} = \frac{\cos (12^s - \Omega)}{\cos A P p}.$$

$= \frac{\cos \Omega}{\cos A P p}$ ,  $\Omega$  designating the longitude of the moon's ascending node.

Again,

$$\frac{\tan A P p}{\tan A P O} = \frac{p i}{O i} = \frac{P d}{P D} = \frac{P d}{P O};$$

$$\begin{aligned} \text{hence, } \tan A P p &= \frac{P d}{P O} \cdot \tan A P O = \frac{P d}{P O} \cdot \tan (12^{\circ} - \Omega) \\ &= - \frac{P d}{P O} \tan \Omega. \end{aligned}$$

Now substitute, and there will result

*The Nutation in North Polar Distance.*

$$\begin{aligned} &= \frac{P O \cos \Omega}{\cos A P p} (\sin A P p \cos R + \cos A P p \sin R) \\ &= P O (\tan A P p \cos R \cos \Omega + \cos \Omega \sin R), \\ &= - P d \cos R \sin \Omega + P O \cos \Omega \sin R, \\ &= - 6''.887 \cos R \sin \Omega + 9''.250 \cos \Omega \sin R \dots (f) \end{aligned}$$

which is the difference, as far as nutation is concerned, between the *mean* and *apparent* north polar distance. The *apparent* north polar distance, therefore, must be had by adding the preceding quantity, with its sign changed, to the mean.

Nutation in right ascension  $= P d \sin \Omega \cot \omega$

$$+ P O \cos \Omega \cos R \cot \delta + P d \sin \Omega \sin R \cot \delta,$$

which, as far as nutation is concerned, is the difference of the mean and apparent right ascensions: and, consequently, the above expression must be subtracted from the mean, in order to obtain the apparent right ascension; or, which is the same, must be added after a negative sign has been prefixed; in which case, we have, substituting for  $P O$ ,  $P d$  their numerical values.

*The Nutation in Right Ascension.*

$$\begin{aligned} &= - 6''.887 \sin \Omega \cot \omega \\ &- 9''.250 \cos \Omega \cos R \cot \delta - 6''.887 \sin \Omega \sin R \cot \delta \dots (g). \end{aligned}$$

Formulæ (f) and (g) are of the same form with (Z) and (a) for the aberration in right ascension and declination, and

therefore formulæ may be derived from them similar to (c) and (e), adapted to logarithmic computation. The quantities corresponding to  $\varphi$ ,  $M$ ,  $\theta$ ,  $N$ , have been calculated for the stars in the catalogue of Table XC, and inserted in Table XCI, in the columns headed  $\varphi'$ ,  $M'$ ,  $\theta'$ ,  $N'$ .

The *Solar Nutation* arises from like causes as the Lunar, and admits of similar formulæ. As an ellipse, made the locus of the true place of the pole, served to exhibit the effects of the lunar nutation, so an ellipse, of different, and much smaller dimensions, may be made to represent the path which the true pole of the equator would, by reason of the sun's inequality of force in causing precession, describe about the mean place of the pole. Thus, in Figure S6, the ellipse  $A d C$  will serve to represent the locus of the pole, when  $A P = 0''.500$ ,  $P d = 0''.545$ , and  $A P O$ , instead of being  $= \Omega$ , is equal to  $2 \odot$ , or twice the sun's longitude, according to the order of the signs; the equations, therefore, for the solar nutation in north polar distance, and right ascension, analogous to those of p. 364 will be

*The Solar Nutation in North Polar Distance.*

$$= - 0''.500 \cos R \sin 2 \odot + 0''.545 \sin R \cos 2 \odot. \dots (h).$$

*The Solar Nutation in Right Ascension.*

$$= - 0''.500 \sin 2 \odot \cot \omega \\ - 0''.545 \cos 2 \odot \cos R \cot \delta - 0''.500 \sin 2 \odot \sin R \cot \delta. \dots (i).$$

If the apparent place of a star should be required with great precision, it would be necessary to compute the solar nutations from these formulæ, and apply them as corrections to the mean right ascension and declination. The calculation would be performed after the same manner as for the lunar nutation; but it is much abridged by remarking that the form of the equations is the same as that of the equations for the lunar nutation, and that the co-efficients are very nearly the 0.075 of those of the latter equations. Thus we can make use of the same arcs  $\varphi'$ ,  $\theta'$ , and  $\log. maxima$ ,  $M'$ ,  $N'$ , repeat the calculation for the lunar nutation, taking  $2 \odot$  instead of  $\Omega$ , and multiply the nutations in right ascension and declination thus obtained by 0.075. The results will be the solar nutations required. (See Prob. XX).

*Formulae for computing the effects of the Oblateness of the Earth's Surface upon the Apparent Zenith Distance and Azimuth of a Star.* (Referred to from Article 148, page 64).

From the centre of the earth, an observer would see a star at *I* (Fig. 85), and would have *V* for his zenith: from the surface his zenith is *Z*, and he sees this star at *B*; *IB* = *p* is the parallax in altitude; the azimuth *VZI* is changed *VZB*. If for a given time, we wish to calculate the apparent zenith distance *BZ*, and the apparent azimuth *VZB*, we have first to resolve the spherical triangle *IZP*, in which we know the two sides *ZP* = co-latitude and *IP* = co-declination, and the included hour angle *P*; the azimuth *VZI* = *A*, and the arc *IZ* = *n* will thus be known. But from the earth's surface, the star is seen at *B*: the azimuth *VZB* = *A* +  $\alpha$ : the zenith distance *BZ* = *n* + *p*, since, *VZ* = *i* being very small, we have sensibly *IB* + *BZ* = *BZ*. By reason of the want of sphericity of the earth, parallax then increases the true azimuth and zenith distance of a star by small quantities,  $\alpha$  and *p*, which it is necessary to calculate. In the triangle *VIZ* we have,

$\cos IV = \cos i \cos n + \sin i \sin n \cos A = \cos n + k \sin n$ ;  
making  $\cos i = 1$ ,  $\sin i = i$ , and  $i \cos A = k$ . Now,  $k \angle i$ , and *a fortiori*  $\cos k = 1$ ,  $\sin k = k$ ; whence

$$\cos IV = \cos n \cos k + \sin n \sin k = \cos (n - k),$$

and  $IV = n - k = n - i \cos A$ .

Thus we correct the calculated arc *n* by the quantity  $-i \cos A$ , to have

$$IV = z = n - i \cos A \dots (j).$$

If this value of *z* be introduced into equation (12), page 44, we shall have *p*, and thence the apparent zenith distance *Z* = *n* + *p* = *BZ*.

Afterwards, to obtain *IZB* =  $\alpha$ , or the *parallax in azimuth*, the triangles *ZBV*, *ZBI* give,

$$\frac{\sin ZBV}{\sin i} = \frac{\sin (A + \alpha)}{\sin (z + p)}, \quad \frac{\sin ZBV}{\sin n} = \frac{\sin \alpha}{\sin p};$$

equating the values of  $\sin ZBV$ ,

$$\frac{\sin n \sin \alpha}{\sin p} = \frac{\sin i \sin (A + \alpha)}{\sin (z + p)}$$

substituting for  $\sin p$  its value  $\sin H \sin (z + p) = \sin H \sin Z$  (equa. 10, page 43), and reducing,

$$\frac{\sin \alpha}{\sin H \sin i} = \frac{\sin (A + \alpha)}{\sin n},$$

and as  $i$  is very small,  $\sin i \sin (A + \alpha)$  does not differ sensibly from  $i \sin A$ , and we thus have *in seconds* (For. 47, page 343),

$$\alpha = \frac{H i \sin A \sin 1''}{\sin n} \dots (k).$$

*Solution of Kepler's Problem, by which a Body's Place is found in an Elliptical Orbit.*

Let  $APB$  be an ellipse,  $E$  the focus occupied by the sun, round which  $P$  the earth or any other planet is supposed to revolve. Let the time and planet's motion be dated from the apside or aphelion  $A$ . The *condition given* is the time elapsed from the planet's quitting  $A$ ; the *result sought* is the place  $P$ ; to be determined either by finding the value of the angle  $AEP$ , or by cutting off, from the whole ellipse, an area  $AEP$  bearing the same proportion to the area of the ellipse which the given time bears to the periodic time.

There are some technical terms used in this problem which we will now explain.

Let a circle  $AMB$  be described on  $AB$  as its diameter, and suppose a point to describe this circle uniformly, and the whole of it, in the same time as the planet describes the ellipse; let also  $t$  denote the time elapsed during  $P$ 's motion from  $A$  to  $P$ ; then if  $AM = \frac{t}{\text{period}} \times 2AMB$ ,  $M$  will be the place of the point that moves uniformly, whilst  $P$  is that of the planet's; the angle  $ACM$  is called the *Mean Anomaly*, and the angle  $AEP$  is called the *True Anomaly*.

Hence, since the time ( $t$ ) being given, the angle  $ACM$  can always be immediately found (see Art. 243, p. 101), we may vary the enunciation of Kepler's problem, and state its object to be *the finding of the true anomaly in terms of the mean*.

Besides the mean and true anomalies, there is a third called the *Eccentric Anomaly*, which is expounded by the angle  $DCA$ ,



and which is always to be found (geometrically) by producing the ordinate  $NP$  of the ellipse to the circumference of the circle. This eccentric anomaly has been devised by mathematicians for the purposes of expediting calculation. It holds a mean place between the two other anomalies, and mathematically connects them. There is one equation by which the mean anomaly is expressed in terms of the eccentric; and another equation by which the true anomaly is expressed in terms of the eccentric.

We will now deduce the two equations by which the *eccentric* is expressed, respectively, in terms of the *true* and *mean* anomalies.

Let  $t$  = time of describing,  $AP$ ,

$\overline{P}$  = periodic time in the ellipse,

$a$  =  $CA$ ,

$ae$  =  $EC$ ,

$v$  =  $\angle PEA$ ,

$u$  =  $\angle DCA$ ; (whence,  $ET$ , perpendicular to  $DT$ , =  $EC$   
 $\times \sin u$ ),

$\rho$  =  $PE$ ,

$\pi$  = 3.14159, &c.;

then, by Kepler's law of the equable description of areas,

$$\begin{aligned} t &= \overline{P} \times \frac{\text{area } PEA}{\text{area of ellip.}} = \overline{P} \times \frac{\text{area } DEA}{\text{area } \odot} = \frac{\overline{P}}{\pi a^2} (DEC + DCA) \\ &= \frac{\overline{P}}{\pi a^2} \left( \frac{ET \cdot DC}{2} + \frac{AD \cdot DC}{2} \right) = \frac{\overline{P}a}{2\pi a^2} (EC \cdot \sin u + DC \cdot u) \\ &= \frac{\overline{P}}{2\pi} (e \sin u + u) : \text{ hence, if we put } \frac{\overline{P}}{2\pi} = \frac{1}{n}, \end{aligned}$$

we have,

$$nt = e \cdot \sin u + u \dots (l),$$

an equation connecting the mean anomaly  $nt$ , and the eccentric  $u$ .

In order to find the other equation, that subsists between the true and eccentric anomaly, we must investigate, and equate, two values of the radius vector  $\rho$ , or  $EP$ .

First value of  $\rho$ , in terms of  $v$  the true anomaly,

$$\rho = \frac{a(1 - e^2)}{1 - e \cos v} \dots (1).$$

Second, in terms of  $u$  the eccentric anomaly,

$$\rho = a(1 + e \cos u) \dots (2).$$

For,  $\rho^2 = \text{EN}^2 + \text{PN}^2$

$$= \text{EN}^2 + \text{DN}^2 \times (1 - e^2)$$

$$= (ae + a \cos u)^2 + a^2 \sin^2 u (1 - e^2)$$

$$= a^2 \{e^2 + 2e \cos u + \cos^2 u\} + a^2(1 - e^2) \sin^2 u$$

$$= a^2 \{1 + 2e \cos u + e^2 \cos^2 u\}.$$

Hence, extracting the square root,

$$\rho = a(1 + e \cos u).$$

Equating the expressions (1), (2), we have

$$(1 - e^2) = (1 - e \cos v)(1 + e \cos u), \text{ whence,}$$

$$\cos v = \frac{e + \cos u}{1 + e \cos u}, \text{ an expression for } v \text{ in terms of } u;$$

but, in order to obtain a formula fitted to logarithmic computation, we must find an expression for  $\tan \frac{v}{2}$ : now, (see For. 12, p. 341),

$$\begin{aligned} \tan \frac{v}{2} &= \sqrt{\left(\frac{1 - \cos v}{1 + \cos v}\right)} = \sqrt{\left(\frac{(1 - e)(1 - \cos u)}{(1 + e)(1 + \cos u)}\right)} \\ &= \sqrt{\left(\frac{1 - e}{1 + e}\right)} \tan \frac{u}{2} \dots (m). \end{aligned}$$

These two expressions (l) and (m), that is,

$$n t = e \cdot \sin u + u,$$

$$\tan \frac{v}{2} = \sqrt{\left(\frac{1 - e}{1 + e}\right)} \tan \frac{u}{2}$$

analytically resolve the problem, and, from such expressions, by certain formulæ belonging to the higher branches of analysis, may  $v$  be expressed in the terms of a series involving  $n t$ .

Instead, however, of this exact but operose and abstruse method of solution, we shall now give an approximate method of expressing the true anomaly in terms of the mean.

MO is drawn parallel to DC. (1.) Find the half difference of the angles at the base of the triangle ECM, from this expression,

$$\tan \frac{1}{2} (CEM - CME) = \tan \frac{1}{2} (CEM + CME) \times \frac{1-e}{1+e},$$

in which, CEM, + CME = ACM, the mean anomaly.

(2.) Find CEM by adding  $\frac{1}{2} (CEM + CME)$  and  $\frac{1}{2} (CEM - CME)$  and use this angle as an approximate value to the eccentric anomaly DCA, from which, however, it really differs by  $\angle EMO$ .

(3.) Use this approximate value of  $\angle DCA = \angle ECT$  in computing ET which equals the arc DM; for, since (see p. 368),

$$t = \frac{P}{\text{area } \odot} \times DEA, \text{ and (the body being supposed to revolve}$$

$$\text{in the circle ADM)} = \frac{P}{\text{area } \odot} \times ACM, \text{ area AED} = \text{area ACM, or, the area DEC} + \text{area ACD} = \text{area DCM} + \text{area ACD};$$

consequently the area DEC = the area DCM; and, expressing their values,

$$\frac{ET \times DC}{2} = \frac{DM \times DC}{2}, \text{ and thus, } ET = DM.$$

Having then computed ET = DM, find the sine of the resulting arc DM, which sine = OT; the difference of the arc and sine (ET - OT) gives EO.

(4.) Use EO in computing the angle EMO, the real difference, between the eccentric anomaly DCA, and the  $\angle MEC$ ; add the computed  $\angle EMO$  to  $\angle MEC$ , in order to obtain  $\angle DCA$ . The result, however, is not the exact value of  $\angle DCA$ , since  $\angle EMO$  has been computed only approximately; that is, by a process which commenced by assuming  $\angle MEC$ , for the value of the  $\angle DCA$ .

For the purpose of finding the eccentric anomaly, this is the entire description of the process; which, if greater accuracy be required, must be repeated; that is, from the last found value of  $\angle DCA = \angle ECT$ , ET, EO, and  $\angle EMO$  must be again computed.

## NOTE TO PROBLEM XIV, (Page 277.)

*Rules for finding the Moon's Longitude, Latitude, Hourly Motions, Equatorial Parallax, and Semi-diameter, for a given time, from the Nautical Almanac.*

Reduce the given time to mean time at Greenwich; then,

*For the Longitude.*

Take from the Nautical Almanac the calculated longitudes answering to the noon and midnight, or midnight and noon, next preceding and next following the given time. Commencing with the longitude answering to the first noon or midnight, subtract each longitude from the next following one: the three remainders will be the *first differences*. Also subtract each first difference from the following for the *second differences*, which will have the plus or minus sign, according as the first differences increase or decrease.

Find the quantity to be added to the second longitude by reason of the first differences, by the proportion,  $12^h$  : excess of given time above time of second longitude : : second first difference : *fourth term*.

With the given time from noon or midnight at the side, take from Table XCIII the quantities corresponding to the minutes, tens of seconds, and seconds of the mean or half sum of the two second differences, at the top: the sum of these will be the *correction for second differences*, which must have the same sign as the mean.

The sum of the second longitude, the fourth term, and the correction for second differences, will be the longitude required.

*For the Latitude.*

Prefix to *north* latitudes the *positive* sign, but to *south* latitudes the *negative* sign, and proceed according to the rules for the longitude, only that attention must now be paid to the signs of the first differences, which may either be plus or minus.

The sign of the resulting latitude will ascertain whether it is *north* or *south*.

*For the Hourly Motion in Longitude.*

Solve the proportion,  $12^h$  : given time from noon or midnight : : half sum of second differences : a fourth term.

Take the sum of the second first difference, half the mean of the second differences, with its sign changed, and this fourth term, and divide it by 12: the quotient will be the required hourly motion in longitude.

*For the Hourly Motion in Latitude.*

With the given time from noon or midnight, the second first difference of latitude, and the mean of the second differences, find the hourly motion in latitude in the same manner as directed for finding the hourly motion in longitude. When the hourly motion is *positive*, the moon is tending *north*; and when it is *negative*, she is tending *south*.

*For the Semi-diameter and Equatorial Parallax.*

The moon's semi-diameter and equatorial parallax may be taken from the Nautical Almanac, with sufficient accuracy, by simple proportion, the correction for second differences being too small to be taken into account, unless great precision is required.

*Corrections for Third and Fourth Differences.*

When the moon's longitude and latitude are required with great precision, corrections must also be applied for the third and fourth differences. To determine these, take from the Almanac the three longitudes or latitudes immediately preceding the given time, and the three immediately following it, and find the first, second, third, and fourth differences, subtracting always each number from the following one, and paying attention to the signs. With the given time from noon or midnight at the side, take from Table XCIV the quantities answering to the minutes and seconds of the middle third difference, at the top. Their sum will be the correction for third differences, which must have the same sign as the middle third difference when the given time from noon or midnight is less than 6 hours; the contrary sign, when the given time is more than 6 hours.



With the given time, and half sum of fourth differences, take from Table XCV the correction for fourth differences, giving it always the same sign as the half sum.

The sum of the third longitude or latitude, the proportional part of the middle first difference answering to the given time from noon or midnight, and the corrections for second, third, and fourth differences, having regard to the signs of all the quantities, will be the longitude or latitude required.



ASTRONOMICAL TABLES.



TABLE I.

*Latitudes and Longitudes from the Meridian of Greenwich, of some cities, and other conspicuous places.*

Names of Places.		Latitude.	Longitude in Degrees	Longitude in Time.
		° ' "	° ' "	h m s
Albany, <i>Capitol</i> ,	New York,	42 39 3 N	73 44 49 W	4 54 59.3
Altona, <i>Obs.</i>	Denmark,	53 32 45 N	9 56 39 E	0 39 46.6
Baltimore, <i>Bat. Mon't.</i> ,	Maryland,	39 17 13 N	76 37 50 W	5 6 31
Berlin, <i>Obs.</i>	Germany,	52 31 13 N	13 23 52 E	0 53 35.5
Boston, <i>State House</i> ,	Mass'ts.,	42 21 15 N	71 4 9 W	4 44 16.6
Bremen, <i>Obs.</i>	Germany,	53 4 36 N	8 48 58 E	0 35 15.9
Brunswick, <i>Bowdoin Coll.</i> ,	Maine,	43 53 0 N	69 58 51 W	4 39 55
Canton,	China,	23 8 9 N	113 16 54 E	7 33 8
Cape of Good Hope, <i>Obs.</i>	Africa,	33 56 3 S	18 28 45 E	1 13 55
Cape Horn,	S. America,	55 58 41 S	67 10 53 W	4 28 43
Charleston, <i>St. Mich's Ch.</i>	S. Carolina,	32 46 33 N	79 57 27 W	5 19 49.8
Charlottesville, <i>Univers.</i> ,	Virginia,	38 2 3 N	78 31 29 W	5 14 6
Cincinnati, <i>Fort Wash.</i> ,	Ohio,	39 5 54 N	84 24 0 W	5 37 36
Copenhagen, <i>Obs.</i>	Denmark,	55 40 53 N	12 34 57 E	0 50 19.8
Dorpat, <i>Obs.</i>	Russia,	58 22 47 N	26 43 45 E	1 46 55
Dublin, <i>Obs.</i>	Ireland,	53 23 13 N	6 20 30 W	0 25 22
Edinburgh, <i>Obs.</i>	Scotland,	55 57 20 N	3 10 54 W	0 12 43.6
Gotha, <i>Obs.</i>	Germany,	50 56 5 N	10 44 6 E	0 42 56.4
Göttingen, <i>Obs.</i>	Germany,	51 31 48 N	9 56 37 E	0 39 46.5
Greenwich, <i>Obs.</i>	England,	51 28 39 N	0 0 0	0 0 0
Königsberg, <i>Obs.</i>	Prussia,	54 42 50 N	20 30 7 E	1 22 0.5
London, <i>St. Paul's Ch.</i> ,	England,	51 30 49 N	0 5 48 W	0 0 23
Marseilles, <i>Obs.</i>	France,	43 17 50 N	5 22 15 E	0 21 29.0
Milan, <i>Obs.</i>	Italy,	45 28 1 N	9 11 48 E	0 36 47.2
Naples, <i>Obs.</i>	Italy,	40 51 47 N	14 15 4 E	0 57 0.3
New Haven, <i>College</i> ,	Connecticut,	41 17 58 N	72 57 46 W	4 51 51.1
New Orleans, <i>City Hall</i> ,	Louisiana,	29 57 45 N	90 6 49 W	6 0 27.3
New York, <i>City Hall</i> ,	New York,	40 42 40 N	74 1 8 W	4 56 4.5
Palermo, <i>Obs.</i>	Italy,	38 6 44 N	13 21 24 E	0 53 25.6
Paramatta, <i>Obs.</i>	New Hol.	33 48 50 S	151 1 34 E	10 4 6.3
Paris, <i>Obs.</i>	France,	48 50 13 N	2 20 24 E	0 9 21.6
Petersburgh, <i>Obs.</i>	Russia,	59 56 31 N	30 18 57 E	2 1 15.8
Philadelphia, <i>Ind'ce. H.</i> ,	Penn.,	39 56 59 N	75 10 59 W	5 0 44
Point Venus,	Otaheite,	17 29 21 S	149 28 55 W	9 57 56
Princeton, <i>College</i> ,	New Jersey,	40 22 N	74 35 W	4 58 20
Providence, <i>University</i> ,	Rhode Isl.	41 49 25 N	71 25 56 W	4 45 44
Quebec, <i>Cast'le</i> ,	L. Canada,	46 49 12 N	71 16 0 W	4 45 4
Richmond, <i>Capitol</i> ,	Virginia,	37 32 17 N	77 27 28 W	5 9 50
Rome, <i>St. Peter's Ch.</i> ,	Italy,	41 54 8 N	12 27 5 E	0 49 48
Savannah, <i>Exchange</i> ,	Georgia,	32 4 56 N	81 7 9 W	5 24 29
Schenectady,	New York,	42 48 N	73 55 W	4 55 40
Stockholm, <i>Obs.</i>	Sweden,	59 20 31 N	18 3 44 E	1 12 15
Turin, <i>Obs.</i>	Italy,	45 4 6 N	7 42 6 E	0 30 48.4
Vienna, <i>Obs.</i>	Austria,	48 12 35 N	16 23 0 E	1 5 32
Wardhus,	Lapland,	70 22 36 N	31 7 54 E	2 4 32
Washington, <i>Capitol</i> ,	Dist. Colum.	38 52 54 N	77 1 48 W	5 8 7.2



2      TABLE II. *Elements of the Planetary Orbits.*

Epoch for Vesta, Juno, Ceres, and Pallas, Jan. 1, 1820, mean noon at Greenwich; for the other planets, Jan. 1, 1801, mean noon at Greenwich.

Planet's name.	Inclination to the Ecliptic.	Sec. Var.	Longitude of ascending node.	Sec. Var.	Longitude of Perihelion.	Sec. Var.
	° ' "	"	° ' "	'	° ' "	'
Mercury	7 0 9.1	+ 18.2	45 57 30.9	+ 70.44	74 21 46.9	+ 93.22
Venus	3 23 28.5	— 4.6	74 54 12.9	+ 51.10	128 43 53.1	+ 78.30
Earth	"	"			99 31 9.9	+ 103.15
Mars	1 51 6.2	— 0.2	48 0 3.5	+ 41.67	332 23 56.6	+ 109.71
Vesta	7 8 9.0	— 12	103 13 18.2	+ 26	249 33 24.4	+ 157
Juno	13 4 9.7		171 7 40.4		53 33 46.0	
Ceres	10 37 26.2	— 44	80 41 24.0	+ 25	147 7 31.5	+ 202
Pallas	34 34 55.0		172 39 26.8		121 7 4.3	
Jupiter	1 18 51.3	— 22.6	98 26 18.9	+ 57.18	11 8 34.6	+ 94.59
Saturn	2 29 35.7	— 15.5	111 56 37.4	+ 51.12	89 9 29.8	+ 115.68
Uranus	0 46 28.4	+ 3.1	72 59 35.3	+ 23.58	167 31 16.1	+ 87.44

Planet's name.	Mean distance from Sun, or Semi-axis.	Mean distance from Sun in miles.	Eccentricity in Parts of the Semi-axis.	Sec. Variation.
Mercury	0.3870981	36814000	0.20551494	+ .000003866
Venus	0.7233316	68787000	0.00686074	— .000062711
Earth	1.0000000	95103000	0.01678357	— .000041630
Mars	1.5236923	144908000	0.09330700	+ .000090176
Vesta	2.3678700	225291000	0.08913000	+ .000004009
Juno	2.6690090	253829000	0.25784800	
Ceres	2.7672450	263165000	0.07843900	— .000005830
Pallas	2.7728860	263707000	0.24164800	
Jupiter	5.2027760	494797000	0.04816210	+ .000159350
Saturn	9.5387861	907162000	0.05615050	— .000312402
Uranus	19.1823900	1824290000	0.04661080	— .000025072

Planet's name.	Mean Long. at the Epoch.	Mean Sidereal Period in Mean Solar Days.	Motion in mean Longitude in 1 yr. of 365 days	Mean Daily Motion in Longitude.
	° ' "	d	° ' "	° ' "
Mercury	166 0 48.6	87.9692580	53 43 3.6	4 5 32.6
Venus	11 33 3.0	224.7007869	224 47 29.7	1 36 7.8
Earth	100 39 13.3	365.2563835	— 0 14 19.5	0 59 8.3
Mars	64 22 55.5	686.9796458	191 17 9.1	0 31 26.7
Vesta	278 30 0.4	1325.7431000	97 28 53	0 16 17.9
Juno	200 16 19.1	1592.6608000	82 25 8	0 13 32.9
Ceres	123 16 11.9	1681.3931000	78 8 8	0 12 50.9
Pallas	108 24 57.9	1686.5388000	77 54 26	0 12 48.4
Jupiter	112 15 23.0	4332.5848212	30 20 31.9	0 4 59.3
Saturn	135 20 6.5	10759.2198174	12 13 36.1	0 2 0.6
Uranus	177 48 23.0	30686.8208296	4 17 45.1	0 0 42.4

TABLE III. *Elements of Moon's Orbit. Epoch, Jan. 1, 1801.*

Mean inclination of orbit. . . . .	5 8 47.9
Mean longitude of node at epoch . . . . .	13 53 17.7
Mean longitude of perigee at epoch . . . . .	266 10 7.5
Mean longitude of moon at epoch . . . . .	118 17 8.3
Mean distance from earth or semi-axis . . . . .	59r.964350
Eccentricity in parts of semi-axis . . . . .	0.0548442
Mean sidereal revolution. . . . .	d h m s d
Mean tropical do. . . . .	27 7 43 11.5 = 27.32166142
Mean synodical do. . . . .	27 7 43 4.7 = 27.3. 158242
Mean anomalistic do. . . . .	29 12 44 2.9 = 29.53058872
Mean anomalous do. . . . .	27 13 18 37.4 = 27.55459950
Mean nodical do. . . . .	27 5 5 36.0 = 27.21222222
Mean revolution of nodes; sider. . . . .	= 6798d.279; trop. = 6788d.50982
Mean revolution of perigee; sider. . . . .	= 3232d.57534; trop. = 3231d.4751

TABLE IV.

3

*Diameters, Volumes, Masses, &c., of Sun, Moon, and Planets.*

	Apparent Diameter.			True Diameter.	Volume.
	Least.	at Mean Distance.	Greatest.		
Mercury	5.0	6.9	12.0	0.398	0.063
Venus	9.6	16.9	61.2	0.975	0.927
Earth				1.000	1.000
Mars	3.6	6.3	18.3	0.517	0.139
Jupiter	30.0	36.7	45.9	10.860	1280.900
Saturn		17.0		9.982	995.000
Uranus		4.0		4.332	80.490
	"	"	"		
Sun	31 31.0	32 1.8	32 35.6	111.454	1384472.000
Moon	29 21.9	31 7.0	33 31.1	0.275	0.020

	Mass.	Density.	Gravity.	Sidereal Rotation.	Light and Heat.
Mercury	$\frac{1}{2025810}$	2.782	1.03	$\begin{smallmatrix} h & m & s \\ 24 & 5 & 28 \end{smallmatrix}$	6.680
Venus	$\frac{1}{403871}$	0.9434	0.98	23 21 7	1.911
Earth	$\frac{1}{357300}$	1.	1.	24 0 0	1.000
Mars	$\frac{1}{2346320}$	0.931	0.33	24 39 21	.431
Jupiter	$\frac{1}{1048.7}$	0.2589	2.72	9 55 50	.037
Saturn	$\frac{1}{3500.2}$	0.1016	1.01	10 29 17	.011
Uranus	$\frac{1}{17918}$	0.2797	0.95	Unknown.	.003
Sun	$\frac{1}{26620200}$	0.2543	27.90	25 12 0	
Moon		0.615	0.16	27 7 43	

TABLE V.

*Elements of the Retrograde Motion of the Planets.*

Planets.	Arc of Retrogradation.	Duration of Retrogradation.		Elongation at the Stations.	Synodical Revolution.
	$\begin{smallmatrix} ^\circ & ' & ^\circ & ' \end{smallmatrix}$	$\begin{smallmatrix} d & h \end{smallmatrix}$	$\begin{smallmatrix} d & h \end{smallmatrix}$	$\begin{smallmatrix} ^\circ & ' & ^\circ & ' \end{smallmatrix}$	
Mercury	9 22 to 15 44	23 12 to	21 12	14 49 to 20 51	116 days.
Venus	14 35 to 17 12	40 21 to	43 12	27 40 to 29 41	584
Mars	10 6 to 19 35	60 18 to	80 15	128 44 to 146 37	780
Jupiter	9 51 to 9 59	116 18 to	122 12	113 35 to 116 42	399
Saturn	6 41 to 6 55	138 18 to	135 9	107 25 to 110 46	378
Uranus	3 36	151		103 30	370

*Elements of the Orbits of the Satellites.*

The distances are expressed in equatorial radii of the primaries. The epoch is Jan. 1, 1801. The periods, &c., are expressed in mean solar days.

I. *Satellites of Jupiter.*

Sat.	Mean Distance.	Sidereal Revolution.	Inclination of Orbit to that of Jupiter.	Mass; that of Jupiter being 1000000000
1	6.04853	<i>d h m</i> 1 18 23	3 5 30	17328
2	9.62347	3 13 14	Variable.	23235
3	15.35024	7 3 43	Variable.	88497
4	26.99835	16 16 32	2 58 48	42659

II. *Satellites of Saturn.*

Sat.	Mean Distance.	Sidereal Revolution.	Eccentricities and Inclinations.
		<i>d h m</i>	
1	3.351	0 22 38	The orbits of the six interior satellites are nearly circular, and very nearly in the plane of the ring. That of the seventh is considerably inclined to the rest and approaches nearer to coincidence with the ecliptic.
2	4.300	1 8 53	
3	5.284	1 21 18	
4	6.819	2 17 45	
5	9.524	4 12 25	
6	22.081	15 22 41	
7	64.359	79 7 55	

III. *Satellites of Uranus.*

Sat.	Mean Distance.	Sidereal Period.	Inclination to Ecliptic.
		<i>d h m s</i>	
1	13.120	5 21 25 0	Their orbits are inclined about 78° 58' to the ecliptic, and their motion is retrograde. The periods of the 2d and 4th require a trifling correction. The orbits appear to be nearly circles.
2	17.022	8 16 56 5	
3	19.845	10 23 4 0	
4	22.752	13 11 8 59	
5	45.507	38 1 48 0	
6	91.008	107 16 40 0	

TABLE VII. *Saturn's Ring.*

	Miles.
Exterior diameter of exterior ring . . . . .	= 176418
Interior ditto . . . . .	= 155272
Exterior diameter of interior ring . . . . .	= 151690
Interior ditto . . . . .	= 117339
Equatorial diameter of the body . . . . .	= 79160
Interval between the planet and interior ring . . . . .	= 19090
Interval of the rings . . . . .	= 1791
Thickness of the rings not exceeding . . . . .	= 100

TABLE X.

7

*Parallax of the Sun, on the first day of each Month: the mean horizontal Parallax being assumed = 8".60.*

Altitude.	Jan.	Feb. Dec.	March. Nov.	April. Oct.	May. Sept.	June. Aug.	July.
0	"	"	"	"	"	"	"
0	8.75	8.73	8.67	8.60	8.53	8.48	8.46
5	8.73	8.69	8.64	8.56	8.50	8.44	8.42
10	8.62	8.59	8.54	8.47	8.40	8.35	8.33
15	8.45	8.43	8.38	8.30	8.24	8.19	8.17
20	8.22	8.20	8.15	8.08	8.01	7.97	7.95
25	7.93	7.91	7.86	7.79	7.73	7.68	7.67
30	7.58	7.56	7.51	7.45	7.39	7.34	7.33
35	7.17	7.15	7.11	7.04	6.99	6.94	6.93
40	6.70	6.68	6.64	6.59	6.53	6.49	6.48
45	6.19	6.17	6.13	6.08	6.03	5.99	5.98
50	5.62	5.61	5.58	5.53	5.48	5.45	5.44
55	5.02	5.01	4.98	4.93	4.89	4.86	4.85
60	4.37	4.36	4.34	4.30	4.26	4.24	4.23
65	3.70	3.69	3.67	3.63	3.60	3.58	3.57
70	2.99	2.98	2.97	2.94	2.92	2.90	2.89
75	2.26	2.26	2.25	2.23	2.21	2.19	2.19
80	1.52	1.52	1.51	1.49	1.48	1.47	1.47
85	0.76	0.76	0.76	0.75	0.74	0.74	0.74
90	0.00	0.00	0.00	0.00	0.00	0.00	0.00

TABLE XI.

*Semi-diurnal Arcs.*

Lat.	Declination.						
	1°	5°	10°	15°	20°	25°	30°
5°	<i>h m</i>	<i>h m</i>	<i>h m</i>	<i>h m</i>	<i>h m</i>	<i>h m</i>	<i>h m</i>
5	6 0	6 2	6 4	6 5	6 7	6 9	6 12
10	6 1	6 4	6 7	6 11	6 15	6 19	6 24
15	6 1	6 5	6 11	6 16	6 22	6 29	6 36
20	6 1	6 7	6 15	6 22	6 30	6 39	6 49
25	6 2	6 9	6 19	6 29	6 39	6 50	7 2
30	6 2	6 12	6 23	6 36	6 49	7 2	7 18
35	6 3	6 14	6 28	6 43	6 59	7 16	7 35
40	6 3	6 17	6 34	6 52	7 11	7 32	7 56
45	6 4	6 20	6 41	7 2	7 25	7 51	8 21
50	6 5	6 24	6 49	7 14	7 43	8 15	8 54
55	6 6	6 29	6 58	7 30	8 5	8 47	9 42
60	6 7	6 35	7 11	7 51	8 36	9 35	12 0
65	6 9	6 43	7 29	8 20	9 25	12 0	

*Equation of Time, to convert Apparent Time into Mean Time.*  
Argument, Mean Longitude of the Sun.

	O <sup>s</sup>	I <sup>s</sup>	II <sup>s</sup>	III <sup>s</sup>	IV <sup>s</sup>	V <sup>s</sup>
°	<i>min. sec.</i>	<i>min. sec.</i>	<i>min. sec.</i>	<i>min. sec.</i>	<i>min. sec.</i>	<i>min. sec.</i>
0	+ 6 58.4	— 1 29.7	— 3 38.7	+ 1 27.0	+ 6 4.1	+ 2 49.7
1	6 39.7	1 42.0	3 34.2	1 40.1	6 6.3	2 34.5
2	6 20.9	1 53.8	3 29.1	1 53.1	6 8.0	2 18.9
3	6 2.1	2 5.2	3 23.5	2 6.0	6 9.1	2 2.8
4	5 43.3	2 15.9	3 17.3	2 18.9	6 9.5	1 46.4
5	5 24.5	2 26.1	3 10.7	2 31.7	6 9.3	1 29.5
6	5 5.7	2 35.9	3 3.5	2 44.3	6 8.5	1 12.3
7	4 46.9	2 45.0	2 56.0	2 56.7	6 7.2	0 54.6
8	4 28.2	2 53.6	2 47.9	3 8.9	6 5.2	0 36.6
9	4 9.6	3 1.8	2 39.5	3 20.8	6 2.5	+ 0 18.2
10	3 51.1	3 9.3	2 30.5	3 32.5	5 59.3	— 0 0.4
11	3 32.6	3 16.3	2 21.2	3 43.9	5 55.4	0 19.5
12	3 14.3	3 22.8	2 11.5	3 55.0	5 51.0	0 38.8
13	2 56.2	3 28.6	2 1.4	4 5.8	5 45.8	0 58.4
14	2 38.3	3 33.9	1 51.0	4 16.3	5 40.1	1 18.2
15	2 20.5	3 38.6	1 40.1	4 26.5	5 33.7	1 38.3
16	2 3.0	3 42.7	1 29.0	4 36.3	5 26.7	1 58.5
17	1 45.7	3 46.3	1 17.6	4 45.7	5 19.2	2 19.1
18	1 28.6	3 49.2	1 5.9	4 54.7	5 11.1	2 39.8
19	1 11.7	3 51.5	0 54.1	5 3.3	5 2.3	3 0.7
20	0 55.2	3 53.3	0 42.0	5 11.3	4 53.0	3 21.6
21	0 39.1	3 54.4	0 29.6	5 18.9	4 43.1	3 42.8
22	0 23.3	3 55.0	0 17.1	5 26.0	4 32.7	4 4.0
23	+ 0 7.8	3 55.0	— 0 4.4	5 32.6	4 21.6	4 25.3
24	— 0 7.3	3 54.5	+ 0 8.4	5 38.6	4 10.1	4 46.6
25	0 22.0	3 53.3	0 21.5	5 44.2	3 57.9	5 8.1
26	0 36.3	3 51.5	0 34.5	5 49.3	3 45.3	5 29.5
27	0 50.3	3 49.2	0 47.6	5 53.9	3 32.1	5 51.0
28	1 3.8	3 46.2	1 0.7	5 57.8	3 18.5	6 12.3
29	1 16.9	3 42.8	1 13.8	6 1.2	3 4.3	6 33.7
30	— 1 29.7	— 3 38.7	+ 1 27.0	+ 6 4.1	+ 2 49.7	— 6 54.9

TABLE XIII.

*Secular Variation of Equation of Time.*  
Argument, Sun's Mean Longitude.

	O <sup>s</sup>	I <sup>s</sup>	II <sup>s</sup>	III <sup>s</sup>	IV <sup>s</sup>	V <sup>s</sup>
<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>
0	— 3	+ 4	+ 11	+ 14	+ 13	+ 9
3	2	5	11	14	13	8
6	1	6	12	14	12	8
9	— 1	6	12	15	12	7
12	0	7	12	14	12	7
15	+ 1	8	13	14	11	6
18	2	8	13	14	11	6
21	2	9	14	14	10	5
24	3	9	14	14	10	5
27	4	10	14	14	9	4
30	+ 4	+ 11	+ 14	+ 13	+ 9	+ 4



*Equation of Time, to convert Apparent Time into Mean Time.*

Argument, Mean Longitude of the Sun.

	VI <sup>s</sup>	VII <sup>s</sup>	VIII <sup>s</sup>	IX <sup>s</sup>	X <sup>s</sup>	XI <sup>s</sup>
°	<i>min. sec.</i>	<i>min. sec.</i>	<i>min. sec.</i>	<i>min. sec.</i>	<i>min. sec.</i>	<i>min. sec.</i>
0	— 6 54.9	— 15 18.9	— 13 58.7	— 1 30.6	+ 11 30.0	+ 14 3.1
1	7 16.1	15 27.9	13 43.0	1 0.2	11 47.0	13 56.0
2	7 37.2	15 36.1	13 26.3	— 0 29.8	12 3.3	13 48.4
3	7 58.3	15 43.7	13 8.9	+ 0 0.6	12 18.7	13 40.1
4	8 19.1	15 50.5	12 50.5	0 31.0	12 33.4	13 31.1
5	8 39.8	15 56.5	12 31.4	1 1.3	12 47.2	13 21.6
6	9 0.2	16 1.8	12 11.6	1 31.4	13 0.1	13 11.4
7	9 20.5	16 6.3	11 51.1	2 1.3	13 12.2	13 0.7
8	9 40.6	16 9.9	11 29.9	2 31.0	13 23.5	12 49.4
9	10 0.3	16 12.9	11 7.9	3 0.5	13 33.9	12 37.4
10	10 19.8	16 15.1	10 45.4	3 29.7	13 43.6	12 25.0
11	10 38.9	16 16.5	10 22.0	3 58.6	13 52.3	12 12.2
12	10 57.8	16 17.0	9 58.1	4 27.1	14 0.2	11 58.9
13	11 16.2	16 16.6	9 33.5	4 55.2	14 7.3	11 45.1
14	11 34.4	16 15.4	9 8.4	5 22.9	14 13.5	11 30.9
15	11 52.1	16 13.4	8 42.6	5 50.2	14 18.9	11 16.3
16	12 9.5	16 10.4	8 16.4	6 17.1	14 23.4	11 1.1
17	12 23.5	16 6.7	7 49.6	6 43.5	14 27.2	10 45.6
18	12 42.9	16 2.1	7 22.5	7 9.3	14 30.0	10 20.7
19	12 58.9	15 56.6	6 54.9	7 34.6	14 32.1	10 13.5
20	13 14.4	15 50.1	6 27.0	7 59.3	14 33.3	9 56.9
21	13 29.5	15 42.9	5 58.5	8 23.4	14 33.7	9 40.1
22	13 44.1	15 34.8	5 29.7	8 45.9	14 33.3	9 23.0
23	13 58.0	15 25.8	5 0.5	9 9.8	14 32.2	9 5.7
24	14 11.4	15 16.0	4 31.0	9 32.0	14 30.2	8 48.0
25	14 24.1	15 5.2	4 1.4	9 53.5	14 27.5	8 30.2
26	14 36.3	14 53.6	3 31.6	10 14.3	14 24.0	8 12.2
27	14 47.9	14 41.1	3 1.5	10 34.4	14 19.9	7 54.0
28	14 58.8	14 27.7	2 31.3	10 53.8	14 15.0	7 35.5
29	15 9.2	14 13.6	2 1.0	11 12.3	14 9.4	7 17.0
30	— 15 18.9	— 13 58.7	— 1 30.6	+ 11 30.0	+ 14 3.1	+ 6 58.4

TABLE XIII.

*Secular Variation of Equation of Time.*

Argument, Sun's Mean Longitude.

	VI <sup>s</sup>	VII <sup>s</sup>	VIII <sup>s</sup>	IX <sup>s</sup>	X <sup>s</sup>	XI <sup>s</sup>
°	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>
0	+4	— 2	— 10	— 15	— 15	— 10
3	3	3	10	15	14	10
6	3	4	11	15	14	9
9	2	4	12	15	14	8
12	1	5	12	15	13	8
15	+1	6	13	15	13	7
18	0	7	13	15	12	6
21	0	7	14	15	12	5
24	— 1	8	14	15	11	5
27	2	9	15	15	11	4
30	— 2	— 10	— 15	— 15	— 10	— 3

*Perturbations of Equation of Time.*

## III.

II.	0	100	200	300	400	500	600	700	800	900	1000
	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>
0	1.4	0.8	1.0	1.7	1.7	1.2	0.7	0.4	0.6	1.4	1.4
100	1.2	1.4	1.1	1.0	1.6	1.8	1.1	0.7	0.6	0.7	1.2
200	0.9	1.0	1.2	1.2	1.2	1.5	1.7	1.1	0.5	0.7	0.9
300	0.7	1.1	1.1	0.9	1.2	1.4	1.5	1.6	1.2	0.5	0.7
400	0.5	0.6	1.2	1.2	0.8	1.0	1.6	1.7	1.5	1.2	0.5
500	1.0	0.5	0.6	1.2	1.4	0.8	0.8	1.5	1.9	1.5	1.0
600	1.7	1.0	0.4	0.5	1.2	1.4	0.9	0.6	1.3	2.0	1.7
700	1.9	1.8	1.1	0.4	0.4	1.1	1.6	1.1	0.7	1.2	1.9
800	1.2	1.8	1.8	1.2	0.4	0.3	1.0	1.6	1.2	0.7	1.2
900	0.7	1.1	1.7	1.8	1.2	0.6	0.2	0.8	1.6	1.3	0.7
1000	1.4	0.8	1.0	1.7	1.7	1.2	0.7	0.4	0.6	1.4	1.4

## II.

## IV.

	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>
0	0.6	0.7	0.5	0.3	0.2	0.6	0.7	0.5	0.2	0.1	0.6
100	0.2	0.7	0.6	0.5	0.2	0.3	0.6	0.9	0.5	0.2	0.2
200	0.2	0.4	0.6	0.5	0.4	0.3	0.4	0.6	0.5	0.5	0.2
300	0.4	0.2	0.5	0.5	0.5	0.4	0.4	0.4	0.5	0.5	0.4
400	0.5	0.4	0.4	0.4	0.4	0.4	0.5	0.5	0.4	0.4	0.5
500	0.4	0.5	0.5	0.5	0.4	0.4	0.3	0.4	0.5	0.3	0.4
600	0.3	0.3	0.5	0.6	0.4	0.4	0.3	0.5	0.7	0.4	0.3
700	0.4	0.2	0.3	0.6	0.6	0.4	0.2	0.2	0.7	0.7	0.4
800	0.6	0.3	0.2	0.3	0.7	0.6	0.3	0.2	0.3	0.8	0.6
900	0.8	0.5	0.3	0.1	0.4	0.7	0.5	0.3	0.1	0.5	0.8
1000	0.6	0.7	0.5	0.3	0.2	0.6	0.7	0.5	0.2	0.1	0.6

## II.

## V.

	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>
0	1.0	1.0	1.1	1.2	1.1	1.0	0.7	0.4	0.6	0.9	1.0
100	0.9	0.9	0.8	1.0	1.3	1.3	1.0	0.7	0.4	0.5	0.9
200	0.5	0.7	0.7	0.8	1.0	1.0	1.1	1.2	0.9	0.3	0.5
300	0.2	0.5	0.7	0.7	0.8	1.2	1.5	1.5	1.1	0.5	0.2
400	0.3	0.2	0.5	0.7	0.7	0.9	1.3	1.4	1.4	1.0	0.3
500	0.8	0.3	0.2	0.5	0.7	0.7	1.0	1.4	1.4	1.4	0.8
600	1.3	0.7	0.3	0.3	0.5	0.7	0.9	1.1	1.4	1.6	1.3
700	1.5	1.1	0.7	0.3	0.4	0.5	0.8	1.0	1.2	1.4	1.5
800	1.3	1.3	1.0	0.7	0.4	0.4	0.6	0.8	1.0	1.2	1.3
900	1.1	1.2	1.2	1.0	0.8	0.6	0.5	0.6	0.9	1.1	1.1
1000	1.0	1.0	1.1	1.2	1.1	1.0	0.7	0.4	0.6	0.9	1.0

*Moon and Nutation.*

I.	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>	<i>sec.</i>
N.	0.5	0.8	1.0	1.0	0.8	0.5	0.2	0.0	0.0	0.2	0.5
	0.1	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.1

Constant 33.0

*For converting any given day into the decimal part of a year of 365 days.*

Day	Jan.	Feb.	March	April	May	June
1	.000	.085	.162	.247	.329	.414
2	.003	.088	.164	.249	.331	.416
3	.006	.090	.167	.252	.334	.419
4	.008	.093	.170	.255	.337	.422
5	.011	.096	.173	.258	.340	.425
6	.014	.099	.175	.260	.342	.427
7	.016	.101	.178	.263	.345	.430
8	.019	.104	.181	.266	.348	.433
9	.022	.107	.184	.268	.351	.436
10	.025	.110	.186	.271	.353	.438
11	.027	.112	.189	.274	.356	.441
12	.030	.115	.192	.277	.359	.444
13	.033	.118	.195	.279	.362	.446
14	.036	.121	.197	.282	.364	.449
15	.038	.123	.200	.285	.367	.452
16	.041	.126	.203	.288	.370	.455
17	.044	.129	.205	.290	.373	.458
18	.046	.132	.208	.293	.375	.460
19	.049	.134	.211	.296	.378	.463
20	.052	.137	.214	.299	.381	.466
21	.055	.140	.216	.301	.384	.468
22	.058	.142	.219	.304	.386	.471
23	.060	.145	.222	.307	.389	.474
24	.063	.148	.225	.310	.392	.477
25	.066	.151	.227	.312	.395	.479
26	.068	.153	.230	.315	.397	.482
27	.071	.156	.233	.318	.400	.485
28	.074	.159	.236	.321	.403	.488
29	.077		.238	.323	.405	.490
30	.079		.241	.326	.408	.493
31	.082		.244		.411	

*For converting any given day into the decimal part of a year of 365 days.*

Day	July	August	Sept.	Oct.	Nov.	Dec.
1	.496	.581	.666	.748	.833	.915
2	.499	.584	.668	.751	.836	.918
3	.501	.586	.671	.753	.838	.921
4	.504	.589	.674	.756	.841	.923
5	.507	.592	.677	.759	.844	.926
6	.510	.595	.679	.762	.846	.929
7	.512	.597	.682	.764	.849	.931
8	.515	.600	.685	.767	.852	.934
9	.518	.603	.688	.770	.855	.937
10	.521	.605	.690	.773	.858	.940
11	.523	.608	.693	.775	.860	.942
12	.526	.611	.696	.778	.863	.945
13	.529	.614	.699	.781	.866	.948
14	.532	.616	.701	.784	.868	.951
15	.534	.619	.704	.786	.871	.953
16	.537	.622	.707	.789	.874	.956
17	.540	.625	.710	.792	.877	.959
18	.542	.627	.712	.795	.879	.962
19	.545	.630	.715	.797	.882	.964
20	.548	.633	.718	.800	.885	.967
21	.551	.636	.721	.803	.888	.970
22	.553	.638	.723	.805	.890	.973
23	.556	.641	.726	.808	.893	.975
24	.559	.644	.729	.811	.896	.978
25	.562	.647	.731	.814	.899	.981
26	.564	.649	.734	.816	.901	.984
27	.567	.652	.737	.819	.904	.986
28	.570	.655	.740	.822	.907	.989
29	.573	.658	.742	.825	.910	.992
30	.575	.660	.745	.827	.912	.995
31	.578	.663		.830		.997

*For converting time into decimal parts of a day.*

Hours		Minutes				Seconds			
h.		m.		m.		s.		s.	
1	.04167	1	.00039	31	.02153	1	.00001	31	.00036
2	.08333	2	.00139	32	.02222	2	.00002	32	.00037
3	.12500	3	.00203	33	.02292	3	.00003	33	.00033
4	.16667	4	.00273	34	.02361	4	.00005	34	.00030
5	.20833	5	.00347	35	.02430	5	.00006	35	.00040
6	.25000	6	.00417	36	.02500	6	.00007	36	.00042
7	.29167	7	.00436	37	.02569	7	.00008	37	.00043
8	.33333	8	.00556	38	.02639	8	.00009	38	.00044
9	.37500	9	.00625	39	.02708	9	.00010	39	.00045
10	.41667	10	.00694	40	.02778	10	.00012	40	.00046
11	.45833	11	.00764	41	.02847	11	.00013	41	.00047
12	.50000	12	.00833	42	.02917	12	.00014	42	.00049
13	.54167	13	.00903	43	.02983	13	.00015	43	.00050
14	.58333	14	.00972	44	.03056	14	.00016	44	.00051
15	.62500	15	.01042	45	.03125	15	.00017	45	.00052
16	.66667	16	.01111	46	.03194	16	.00018	46	.00053
17	.70833	17	.01180	47	.03264	17	.00020	47	.00054
18	.75000	18	.01250	48	.03333	18	.00021	48	.00056
19	.79167	19	.01319	49	.03403	19	.00022	49	.00057
20	.83333	20	.01389	50	.03472	20	.00023	50	.00058
21	.87500	21	.01458	51	.03542	21	.00024	51	.00059
22	.91667	22	.01528	52	.03611	22	.00025	52	.00060
23	.95833	23	.01597	53	.03680	23	.00027	53	.00061
24	1.00000	24	.01667	54	.03750	24	.00028	54	.00062
		25	.01736	55	.03819	25	.00029	55	.00064
		26	.01805	56	.03889	26	.00030	56	.00065
		27	.01875	57	.03958	27	.00031	57	.00066
		28	.01944	58	.04028	28	.00032	58	.00067
		29	.02014	59	.04097	29	.00034	59	.00068
		30	.02083	60	.04167	30	.00035	60	.00069



*For converting Minutes and Seconds of a degree, into the decimal division of the same.*

Minutes				Seconds			
'		'		"		"	
1	.01667	31	.51667	1	.00028	31	.00861
2	.03333	32	.53333	2	.00056	32	.00889
3	.05000	33	.55000	3	.00083	33	.00917
4	.06667	34	.56667	4	.00111	34	.00944
5	.08333	35	.58333	5	.00139	35	.00972
6	.10000	36	.60000	6	.00167	36	.01000
7	.11667	37	.61667	7	.00194	37	.01028
8	.13333	38	.63333	8	.00222	38	.01056
9	.15000	39	.65000	9	.00250	39	.01083
10	.16667	40	.66667	10	.00278	40	.01111
11	.18333	41	.68333	11	.00306	41	.01139
12	.20000	42	.70000	12	.00333	42	.01167
13	.21667	43	.71667	13	.00361	43	.01194
14	.23333	44	.73333	14	.00389	44	.01222
15	.25000	45	.75000	15	.00417	45	.01250
16	.26667	46	.76667	16	.00444	46	.01278
17	.28333	47	.78333	17	.00472	47	.01306
18	.30000	48	.80000	18	.00500	48	.01333
19	.31667	49	.81667	19	.00528	49	.01361
20	.33333	50	.83333	20	.00556	50	.01389
21	.35000	51	.85000	21	.00583	51	.01417
22	.36667	52	.86667	22	.00611	52	.01444
23	.38333	53	.88333	23	.00639	53	.01472
24	.40000	54	.90000	24	.00667	54	.01500
25	.41667	55	.91667	25	.00694	55	.01528
26	.43333	56	.93333	26	.00722	56	.01556
27	.45000	57	.95000	27	.00750	57	.01583
28	.46667	58	.96667	28	.00778	58	.01611
29	.48333	59	.98333	29	.00806	59	.01639
30	.50000	60	1.00000	30	.00833	60	.01667

TABLE XVIII.

*Sun's Epochs.*

Years.	M. Long.	Long.Peri.	I	II	III	IV	V	N	VI	VII
	s ° ' "	s ° ' "								
1830	9 10 37 46.9	9 10 0 54	228	279	169	598	758	519	989	362
1831	9 10 23 27.4	9 10 1 55	588	278	793	130	842	573	235	396
1832B.	9 10 9 7.9	9 10 2 57	948	278	418	661	926	627	482	430
1833	9 10 53 56.8	9 10 3 59	342	280	47	194	11	681	764	464
1834	9 10 39 37.3	9 10 5 0	702	279	671	725	95	734	11	498
1835	9 10 25 17.8	9 10 6 2	62	279	296	256	179	788	257	532
1836B.	9 10 10 58.4	9 10 7 3	422	278	920	788	261	842	504	566
1837	9 10 55 47.2	9 10 8 5	816	280	549	321	348	895	787	600
1838	9 10 41 27.8	9 10 9 6	176	279	173	852	432	949	33	624
1839	9 10 27 8.3	9 10 10 8	536	279	798	383	517	3	279	668
1840B.	9 10 12 48.8	9 10 11 9	896	278	422	915	601	56	526	702
1841	9 10 57 37.7	9 10 12 11	290	280	51	447	685	110	809	736
1842	9 10 43 18.2	9 10 13 12	650	279	676	979	770	164	55	770
1843	9 10 28 58.8	9 10 14 14	10	279	300	510	554	218	301	804
1844B.	9 10 14 39.3	9 10 15 15	370	278	924	41	938	272	548	838
1845	9 10 59 28.2	9 10 16 17	764	280	553	574	23	325	831	872
1846	9 10 45 8.7	9 10 17 19	124	280	177	106	107	379	77	906
1847	9 10 30 49.2	9 10 18 20	484	279	802	637	191	433	324	940
1848B.	9 10 16 29.8	9 10 19 22	844	278	427	168	276	487	570	900
1849	9 11 1 18.6	9 10 20 23	238	280	55	700	360	540	853	8
1850	9 10 46 59.2	9 10 21 25	598	280	680	231	414	54	99	41
1851	9 10 32 39.7	9 10 22 26	958	279	304	762	529	608	346	75
1852B.	9 10 18 20.2	9 10 23 28	319	278	929	294	613	701	592	109
1853	9 11 3 9.1	9 10 24 29	713	280	557	827	697	755	875	143
1854	9 10 48 49.6	9 10 25 31	73	280	182	358	782	809	121	177
1855	9 10 34 30.2	9 10 26 32	483	279	806	889	866	863	268	211
1856B.	9 10 20 10.7	9 10 27 34	793	279	430	421	950	916	614	245
1857	9 11 4 59.6	9 10 28 35	187	281	60	953	35	970	897	279
1858	9 10 59 40.1	9 10 29 37	547	280	684	485	119	24	144	313
1859	9 10 36 20.7	9 10 30 39	907	279	308	16	203	78	390	347
1860B.	9 10 22 1.2	9 10 31 40	267	279	933	547	288	131	636	381
1861	9 11 6 50.1	9 10 32 42	661	281	562	80	372	185	919	415
1862	9 10 52 30.6	9 10 33 43	21	280	186	612	456	239	166	449
1863	9 10 38 11.1	9 10 34 45	381	280	810	143	541	292	412	483
1864B.	9 10 23 51.7	9 10 35 46	741	279	435	674	625	346	659	517
1865	9 11 8 40.5	9 10 36 48	135	281	64	207	709	400	941	551
1866	9 10 54 21.1	9 10 37 49	495	280	688	738	794	453	188	585
1867	9 10 40 1.6	9 10 38 51	855	280	313	270	878	507	434	619
1868B.	9 10 25 42.2	9 10 39 52	215	279	937	801	962	561	681	653
1869	9 11 10 31.0	9 10 40 54	609	281	566	334	47	615	963	687
1870	9 10 56 11.6	9 10 41 56	969	280	190	865	131	668	210	721

*Sun's Motions for Months.*

Months		M.	Long.	Per.	I	II	III	IV	V	N	VI	VII
		s	o	'	"	"						
January		0	0	0	0.0	0	0	0	0	0	0	0
February		1	0	33	18.2	5	47	85	138	45	7	5
March	{ Com.	1	23	9	11.4	10	933	102	273	86	14	9
	{ Bis.	1	29	8	19.8	10	27	164	277	87	14	9
April	{ Com.	2	23	42	20.7	15	42	245	401	131	21	13
	{ Bis.	2	29	41	38.0	15	70	249	465	132	21	13
May	{ Com.	3	23	16	29.6	20	59	329	534	175	23	18
	{ Bis.	3	29	15	47.9	20	52	331	538	176	23	18
June	{ Com.	4	28	49	57.9	26	110	414	672	220	35	22
	{ Bis.	4	29	49	6.2	26	144	416	676	221	35	23
July	{ Com.	5	28	24	7.8	31	129	496	806	263	41	27
	{ Bis.	5	29	23	16.1	31	163	499	810	265	42	27
Aug.	{ Com.	6	28	57	26.1	36	182	580	943	309	49	31
	{ Bis.	6	29	56	34.4	36	216	583	948	310	49	31
Sep.	{ Com.	7	29	30	44.2	41	233	665	81	354	56	36
	{ Bis.	8	0	29	52.6	41	268	668	86	355	56	36
Oct.	{ Com.	8	29	4	54.1	46	250	748	215	397	63	40
	{ Bis.	9	0	4	2.5	46	284	750	219	399	63	40
Nov.	{ Com.	9	29	38	12.5	51	300	832	353	443	70	45
	{ Bis.	10	0	37	20.7	51	333	835	357	444	70	45
Dec.	{ Com.	10	29	12	22.3	56	313	915	486	486	77	49
	{ Bis.	11	0	11	30.6	56	347	917	491	488	77	49

TABLE XX.

*Sun's Motions for Days and Hours.*

Days	M.	Long.	Per.	I	II	III	IV	V	N	VI	VII	Hrs.	Long.	I	II	III
	o	'	"	"									"			
1	0	0	0.0	0	0	0	0	0	0	0	0	1	2 27.8	1	0	0
2	0	59	8.3	0	34	3	4	1	0	0	36	0	4 55.7	3	0	0
3	1	58	16.7	0	68	5	9	3	0	0	73	0	7 23.5	4	0	1
4	2	57	25.0	0	101	8	13	4	1	0	109	0	9 51.4	6	0	1
5	3	56	33.3	1	135	11	18	6	1	1	145	0	12 19.2	7	1	1
6	4	55	41.6	1	169	14	22	7	1	1	181	0	14 47.1	8	1	1
7	5	54	50.0	1	203	16	27	9	1	1	218	1	17 14.9	10	1	1
8	6	53	58.3	1	236	19	31	10	2	1	254	1	19 42.8	11	1	1
9	7	53	6.6	1	270	22	36	12	2	1	290	1	22 10.6	13	1	2
10	8	52	15.0	1	304	25	40	13	2	1	327	1	24 38.5	14	1	2
11	9	51	23.3	2	338	27	44	15	2	1	363	1	27 6.3	16	1	2
12	10	50	31.6	2	371	30	49	16	2	2	399	1	29 34.2	17	1	2
13	11	49	40.0	2	405	33	53	17	3	2	435	1	32 2.0	18	1	2
14	12	48	48.3	2	439	36	58	19	3	2	472	1	34 29.9	20	2	3
15	13	47	56.6	2	473	38	62	20	3	2	508	2	36 57.7	21	2	3
16	14	47	4.9	2	506	41	67	22	3	2	544	2	39 25.6	23	2	3
17	15	46	13.3	3	540	44	71	23	4	2	581	2	41 53.4	24	2	3
18	16	45	21.6	3	574	47	76	25	4	2	617	2	44 21.2	25	2	3
19	17	44	29.9	3	608	49	80	26	4	3	653	2	46 49.1	27	2	4
20	18	43	38.3	3	641	52	85	28	4	3	690	2	49 16.9	28	2	4
21	19	42	46.6	3	675	55	89	29	5	3	726	2	51 44.8	30	2	4
22	20	41	54.9	4	709	58	93	31	5	3	762	2	54 12.6	31	2	4
23	21	41	3.3	4	743	60	98	32	5	3	798	2	56 40.5	32	3	4
24	22	40	11.6	4	777	63	102	33	5	3	835	2	59 8.3	34	3	4
25	23	39	19.9	4	810	66	107	35	5	4	871	2				
26	24	38	28.2	4	844	68	111	36	6	4	907	2				
27	25	37	36.6	4	878	71	116	38	6	4	943	2				
28	26	36	44.9	5	912	74	120	39	6	4	980	2				
29	27	35	53.2	5	945	77	125	41	6	4	16	3				
30	28	35	1.6	5	979	79	129	42	7	4	52	3				
31	29	34	9.9	5	13	82	134	44	7	4	89	3				

TABLE XXI.

*Sun's Motions for Minutes and Seconds.*

Min.	Long.	Min.	Long.	Sec.	Lon.	Sec.	Lon.
	' "		' "		"		"
1	0 2.5	31	1 16.4	1	0.0	31	1.3
2	4.9	32	1 18.8	2	0.1	32	1.3
3	7.4	33	1 21.3	3	0.1	33	1.4
4	9.9	34	1 23.8	4	0.2	34	1.4
5	12.3	35	1 26.2	5	0.2	35	1.4
6	14.8	36	1 28.7	6	0.2	36	1.5
7	17.2	37	1 31.2	7	0.3	37	1.5
8	19.7	38	1 33.6	8	0.3	38	1.6
9	22.2	39	1 36.1	9	0.4	39	1.6
10	24.6	40	1 38.6	10	0.4	40	1.6
11	27.1	41	1 41.0	11	0.5	41	1.7
12	29.6	42	1 43.5	12	0.5	42	1.7
13	32.0	43	1 46.0	13	0.5	43	1.8
14	34.5	44	1 48.4	14	0.6	44	1.8
15	37.0	45	1 50.9	15	0.6	45	1.8
16	39.4	46	1 53.3	16	0.7	46	1.9
17	41.9	47	1 55.8	17	0.7	47	1.9
18	44.4	48	1 58.3	18	0.7	48	2.0
19	46.8	49	2 0.7	19	0.8	49	2.0
20	49.3	50	2 3.2	20	0.8	50	2.0
21	51.7	51	2 5.7	21	0.9	51	2.1
22	54.2	52	2 8.1	22	0.9	52	2.1
23	56.7	53	2 10.6	23	0.9	53	2.2
24	59.1	54	2 13.1	24	1.0	54	2.2
25	1 1.6	55	2 15.5	25	1.0	55	2.3
26	1 4.1	56	2 18.0	26	1.1	56	2.3
27	1 6.5	57	2 20.5	27	1.1	57	2.3
28	1 9.0	58	2 22.9	28	1.1	58	2.4
29	1 11.5	59	2 25.4	29	1.2	59	2.4
30	1 13.9	60	2 27.8	30	1.2	60	2.5

TABLE XXII. 17

*Mean Obliquity of the Ecliptic.*

Years	23 27'
	"
1835	38 80
1836	38.35
1837	37.89
1838	37.43
1839	36.98
1840	36.52
1841	36.06
1842	35.61
1843	35.15
1844	34.69
1845	34.23
1846	33.78
1847	33.32
1848	32.86
1849	32.41
1850	31.95
1851	31.49
1852	31.04
1853	30.58
1854	30.12
1855	29.66
1856	29.21
1857	28.75
1858	28.29
1859	27.84
1860	27.38
1861	26.92
1862	26.47
1863	26.01
1864	25.55

TABLE XXIII.

*Sun's Hourly Motion.*

Argument. Sun's Mean Anomaly.

	Os	Is	IIs	IIIs	IVs	Vs	
°	' "	' "	' "	' "	' "	' "	°
0	2 32.92	2 32.20	2 30.23	2 27.74	2 25.32	2 23.60	30
10	2 32.84	2 31.67	2 29.46	2 26.89	2 24.64	2 23.26	20
20	2 32.59	2 31.02	2 28.61	2 26.07	2 24.06	2 23.05	10
30	2 32.20	2 30.28	2 27.74	2 25.32	2 23.60	2 22.99	0
	XIs	Xs	IXs	VIIIs	VIIs	VIs	

TABLE XXIV.

*Sun's Semi-diameter.*

Argument. Sun's Mean Anomaly.

	Os	Is	IIs	IIIs	IVs	Vs	
°	' "	' "	' "	' "	' "	' "	°
0	16 17.3	16 15.0	16 8.8	16 0.6	15 52.7	15 47.0	30
10	16 17.0	16 13.3	16 6.2	15 57.8	15 50.5	15 45.9	20
20	16 16.2	16 11.2	16 3.4	15 55.1	15 48.6	15 45.2	10
30	16 15.0	16 8.8	16 0.6	15 52.7	15 47.0	15 45.0	0
	XIs	Xs	IXs	VIIIs	VIIs	VIs	

TABLE XXV.  
*Equation of the Sun's Centre.*  
 Argument. Sun's Mean Anomaly.

	Os				Is				IIs				IIIs				IVs				Vs			
°	'	''	'''	''''	°	'	''	'''	°	'	''	'''	°	'	''	'''	°	'	''	'''	°	'	''	'''
0	11	29	59	13.9	0	57	58.5		1	40	10.7		1	54	34.1		1	38	4.8		0	55	52.6	
1	0	0	1	17.3	0	59	43.9		1	41	8.9		1	54	30.5		1	37	2.4		0	54	8.7	
2		0	3	20.6	1	1	23.0		1	42	5.1		1	54	24.8		1	35	53.1		0	52	24.0	
3		0	5	23.9	1	3	10.9		1	42	59.3		1	54	17.0		1	34	52.2		0	50	33.2	
4		0	7	27.0	1	4	52.6		1	43	51.8		1	54	7.1		1	33	44.6		0	48	51.6	
5		0	9	30.0	1	6	33.0		1	44	42.1		1	53	55.2		1	32	35.4		0	47	4.2	
6		0	11	32.8	1	8	12.3		1	45	30.4		1	53	41.0		1	31	24.4		0	45	16.0	
7		0	13	35.4	1	9	50.1		1	46	16.8		1	53	24.9		1	30	11.9		0	43	26.9	
8		0	15	37.7	1	11	26.5		1	47	1.2		1	53	6.7		1	28	57.7		0	41	37.0	
9		0	17	39.6	1	13	1.7		1	47	43.5		1	52	46.5		1	27	42.0		0	39	46.5	
10		0	19	41.2	1	14	35.3		1	48	23.9		1	52	24.2		1	26	24.8		0	37	55.3	
11		0	21	42.4	1	16	7.5		1	49	2.2		1	51	59.8		1	25	5.9		0	36	3.3	
12		0	23	43.1	1	17	38.2		1	49	38.4		1	51	33.4		1	23	45.7		0	34	10.8	
13		0	25	43.4	1	19	7.5		1	50	12.6		1	51	5.0		1	22	23.8		0	32	17.7	
14		0	27	43.2	1	20	35.2		1	50	44.7		1	50	34.5		1	21	0.6		0	30	23.8	
15		0	29	42.3	1	22	1.5		1	51	14.9		1	50	2.2		1	19	36.0		0	28	29.6	
16		0	31	40.9	1	23	26.0		1	51	42.9		1	49	27.7		1	18	9.9		0	26	34.8	
17		0	33	38.9	1	24	43.9		1	52	8.7		1	48	51.3		1	16	42.4		0	24	39.6	
18		0	35	36.2	1	26	10.3		1	52	32.5		1	48	13.0		1	15	13.7		0	22	43.9	
19		0	37	32.9	1	27	30.0		1	52	54.3		1	47	32.7		1	13	43.5		0	20	47.9	
20		0	39	28.8	1	28	48.0		1	53	13.9		1	46	50.4		1	12	12.1		0	18	51.4	
21		0	41	23.9	1	30	4.2		1	53	31.4		1	46	6.3		1	10	39.3		0	16	54.6	
22		0	43	18.1	1	31	18.8		1	53	46.8		1	45	20.3		1	9	5.4		0	14	57.5	
23		0	45	11.5	1	32	31.7		1	54	0.1		1	44	32.2		1	7	30.3		0	13	0.1	
24		0	47	4.0	1	33	42.7		1	54	11.2		1	43	42.4		1	5	54.0		0	11	2.6	
25		0	48	55.6	1	34	52.0		1	54	20.4		1	42	50.7		1	4	16.5		0	9	4.8	
26		0	50	46.3	1	35	59.4		1	54	27.2		1	41	57.1		1	2	37.8		0	7	6.9	
27		0	52	36.0	1	37	5.1		1	54	32.1		1	41	1.7		1	0	58.0		0	5	8.7	
28		0	54	24.6	1	38	8.8		1	54	34.9		1	40	4.5		0	59	17.3		0	3	10.5	
29		0	56	12.1	1	39	10.8		1	54	35.4		1	39	5.6		0	57	35.4		0	1	12.2	
30		0	57	58.5	1	40	10.7		1	54	34.1		1	38	4.8		0	55	52.6					

TABLE XXVI.  
*Secular Variation of Equation of Sun's Centre.*  
 Argument. Sun's Mean Anomaly.

	Os	Is	IIs	IIIs	IVs	Vs
°	''	''	''	''	''	''
0	— 0	— 9	— 15	— 17	— 15	— 8
2	1	9	15	17	14	8
4	1	10	16	17	14	7
6	2	10	16	17	14	7
8	2	11	16	17	13	6
10	3	11	16	17	13	6
12	4	12	17	17	12	5
14	4	12	17	16	12	5
16	5	13	17	16	12	4
18	5	13	17	16	11	3
20	6	13	17	16	11	3
22	7	14	17	16	10	2
24	7	14	17	15	10	2
26	8	15	17	15	9	1
28	8	15	17	15	9	1
30	— 9	— 15	— 17	— 15	— 8	— 0



TABLE XXV.

19

*Equation of the Sun's Centre.*

Argument. Sun's Mean Anomaly.

	VI <sup>s</sup>	VII <sup>s</sup>	VIII <sup>s</sup>	IX <sup>s</sup>	X <sup>s</sup>	XI <sub>s</sub>
	11 <sup>s</sup>	11 <sup>s</sup>	11 <sup>s</sup>	11 <sup>s</sup>	11 <sup>s</sup>	11 <sub>s</sub>
°	° ' "	° ' "	° ' "	° ' "	° ' "	° ' "
0	29 59 13.9	29 2 35.2	28 20 23.0	28 3 53.7	28 18 17.1	29 0 29.3
1	29 57 15.6	29 0 52.4	28 19 22.2	28 3 52.3	28 19 17.0	29 2 15.7
2	29 55 17.3	28 59 10.5	28 18 23.3	28 3 52.8	28 20 19.0	29 4 3.2
3	29 53 19.1	28 57 29.8	28 17 26.1	28 3 55.6	28 21 22.7	29 5 51.8
4	29 51 20.9	28 55 50.0	28 16 30.7	28 4 0.5	28 22 28.4	29 7 41.5
5	29 49 23.0	28 54 11.4	28 15 37.1	28 4 7.4	28 23 35.8	29 9 32.2
6	29 47 25.2	28 52 33.8	28 14 45.4	28 4 16.6	28 24 45.1	29 11 23.8
7	29 45 27.7	28 50 57.5	28 13 55.6	28 4 27.7	28 25 56.1	29 13 16.3
8	29 43 30.3	28 49 22.4	28 13 7.5	28 4 41.0	28 27 9.0	29 15 9.7
9	29 41 33.2	28 47 48.5	28 12 21.5	28 4 56.4	28 28 23.6	29 17 3.9
10	29 39 36.4	28 46 15.7	28 11 37.4	28 5 13.9	28 29 39.8	29 18 59.0
11	29 37 39.9	28 44 44.3	28 10 55.1	28 5 33.5	28 30 57.8	29 20 54.9
12	29 35 43.9	28 43 14.1	28 10 14.8	28 5 55.3	28 32 17.5	29 22 51.6
13	29 33 48.2	28 41 45.4	28 9 36.5	28 6 19.1	28 33 38.9	29 24 48.9
14	29 31 53.0	28 40 17.9	28 9 0.0	28 6 44.9	28 35 1.8	29 26 46.9
15	29 29 58.2	28 38 51.8	28 8 25.6	28 7 12.9	28 36 26.3	29 28 45.5
16	29 28 4.0	28 37 27.2	28 7 53.2	28 7 43.1	28 37 52.6	29 30 44.6
17	29 26 10.1	28 36 4.0	28 7 22.8	28 8 15.2	28 39 20.3	29 32 44.4
18	29 24 17.0	28 34 42.1	28 6 54.4	28 8 49.4	28 40 49.6	29 34 44.7
19	29 22 24.5	28 33 21.9	28 6 28.0	28 9 25.6	28 42 20.3	29 36 45.4
20	29 20 32.5	28 32 3.0	28 6 3.6	28 10 3.9	28 43 52.5	29 38 46.6
21	29 18 41.3	28 30 45.8	28 5 41.4	28 10 44.3	28 45 26.1	29 40 48.2
22	29 16 50.8	28 29 30.1	28 5 21.1	28 11 26.6	28 47 1.3	29 42 50.1
23	29 15 0.9	28 28 15.9	28 5 2.9	28 12 11.0	28 48 37.7	29 44 52.5
24	29 13 11.8	28 27 3.4	28 4 46.8	28 12 57.4	28 50 15.5	29 46 55.0
25	29 11 23.6	28 25 52.4	28 4 32.6	28 13 45.7	28 51 54.8	29 48 57.8
26	29 9 36.2	28 24 43.2	28 4 20.7	28 14 36.0	28 53 35.2	29 51 0.8
27	29 7 49.5	28 23 35.6	28 4 10.8	28 15 28.5	28 55 16.9	29 53 3.9
28	29 6 3.8	28 22 29.7	28 4 3.0	28 16 22.7	28 56 59.8	29 55 7.2
29	29 4 19.1	28 21 25.4	28 3 57.3	28 17 18.9	28 58 43.9	29 57 10.5
30	29 2 35.2	28 20 23.0	28 3 53.7	28 18 17.1	29 0 29.3	29 59 13.9

TABLE XXVI.

*Secular Variation of Equation of Sun's Centre.*

Argument. Sun's Mean Anomaly.

	VI <sup>s</sup>	VII <sup>s</sup>	VIII <sup>s</sup>	IX <sup>s</sup>	X <sup>s</sup>	XI <sup>s</sup>
°	"	"	"	"	"	'
0	+ 0	+ 8	+ 15	+ 17	+ 15	+ 9
2	1	9	15	17	15	8
4	1	9	15	17	15	8
6	2	10	15	17	14	7
8	2	10	16	17	14	7
10	3	11	16	17	14	6
12	3	11	16	17	13	6
14	4	12	16	17	13	5
16	5	12	16	17	12	4
18	5	12	17	17	12	4
20	6	13	17	16	11	3
22	6	13	17	16	11	2
24	7	14	17	16	10	2
26	7	14	17	16	10	1
28	8	14	17	15	9	1
30	+ 8	+ 15	+ 17	+ 15	+ 9	+ 0

*Nutations.*

Argument. Supplement of the Node, or N.

*Solar Nutation.*

N.	Long.	R. Asc.	Obliq.	N.	Long.	R. Asc.	Obliq.		Long.	Obliq.		
0	+	0.0	+0.0	+	9.2	509	— 0.0	— 0.0	— 9.3	Jan.	—	—
10	+	1.0	1.0	9.1	510	1.1	1.9	9.3	1	+ 0.5	— 0.5	
20	+	2.1	2.1	9.1	520	2.2	2.0	9.3	11	0.8	0.4	
30	+	3.2	3.0	9.0	530	3.3	2.9	9.2	21	1.1	0.2	
40	+	4.2	4.0	8.9	540	4.4	3.9	9.0	31	1.2	— 0.1	
50	+	5.2	+ 4.9	+ 8.7	550	— 5.5	— 4.8	— 8.9	Feb.	—	—	
60	+	6.2	6.0	8.5	560	6.5	5.7	8.7	10	1.2	+ 0.1	
70	+	7.2	6.9	8.3	570	7.5	6.6	8.4	20	1.0	0.3	
80	+	8.2	7.8	8.1	580	8.5	7.5	8.1	March.	—	—	
90	+	9.1	8.7	7.8	590	9.5	8.4	7.8	2	0.7	0.4	
100	+	10.0	+ 9.4	+ 7.5	600	— 10.4	— 9.1	— 7.5	12	+ 0.3	0.5	
110	+	10.8	10.3	7.1	610	11.2	9.9	7.1	22	— 0.1	0.5	
120	+	11.6	11.1	6.7	620	12.0	10.6	6.7	April.	—	—	
130	+	12.4	11.7	6.3	630	12.8	11.4	6.3	1	0.5	0.5	
140	+	13.1	12.4	5.9	640	13.5	12.0	5.9	11	0.8	0.2	
150	+	13.8	+ 13.0	+ 5.5	650	— 14.2	— 12.6	— 5.4	21	1.1	0.2	
160	+	14.4	13.6	5.0	660	14.8	13.2	4.9	May.	—	—	
170	+	15.0	14.1	4.5	670	15.3	13.8	4.4	1	1.2	+ 0.1	
180	+	15.5	14.5	4.0	680	15.8	14.2	3.9	11	1.2	— 0.1	
190	+	15.9	14.8	3.5	690	16.2	14.7	3.3	21	1.1	0.3	
200	+	16.3	+ 15.1	+ 2.9	700	— 16.6	— 15.0	— 2.8	31	0.8	0.4	
210	+	16.6	15.4	2.4	710	16.9	15.3	2.2	June.	—	—	
220	+	16.9	15.6	1.8	720	17.1	15.4	1.6	10	0.4	0.5	
230	+	17.1	15.7	1.2	730	17.2	15.7	1.1	20	— 0.0	0.5	
240	+	17.2	15.9	0.7	740	17.3	15.9	— 0.5	30	+ 0.4	0.5	
250	+	17.3	+ 15.9	+ 0.1	750	— 17.3	— 15.9	+ 0.1	July.	—	—	
260	+	17.3	15.9	— 0.5	760	17.2	15.9	0.7	10	0.7	0.4	
270	+	17.2	15.7	1.1	770	17.1	15.7	1.2	20	1.0	0.3	
280	+	17.1	15.6	1.6	780	16.9	15.4	1.8	30	1.2	— 0.1	
290	+	16.9	15.4	2.2	790	16.6	15.3	2.4	Aug.	—	—	
300	+	16.6	+ 15.1	— 2.8	800	— 16.3	— 15.0	+ 2.9	9	1.3	+ 0.0	
310	+	16.2	14.8	3.3	810	15.9	14.7	3.5	19	1.2	0.4	
320	+	15.8	14.5	3.9	820	15.5	14.2	4.0	29	0.9	0.4	
330	+	15.3	14.1	4.4	830	15.0	13.8	4.5	Sept.	—	—	
340	+	14.8	13.6	4.9	840	14.4	13.2	5.0	8	0.6	0.5	
350	+	14.2	+ 13.0	— 5.4	850	— 13.8	— 12.6	+ 5.5	18	+ 0.2	0.5	
360	+	13.5	12.4	5.9	860	13.1	12.0	5.9	28	— 0.2	0.5	
370	+	12.8	11.7	6.3	870	12.4	11.4	6.3	Oct.	—	—	
380	+	12.0	11.1	6.7	880	11.6	10.6	6.7	8	0.6	0.5	
390	+	11.2	10.3	7.1	890	10.8	9.9	7.1	18	1.0	0.3	
400	+	10.4	+ 9.4	— 7.5	900	— 10.0	— 9.1	+ 7.5	28	1.2	0.2	
410	+	9.5	8.7	7.8	910	9.1	8.4	7.8	Nov.	—	—	
420	+	8.5	7.8	8.1	920	8.2	7.5	8.1	7	1.2	+ 0.0	
430	+	7.5	6.9	8.4	930	7.2	6.6	8.3	17	1.2	0.2	
440	+	6.5	6.0	8.7	940	6.2	5.7	8.5	27	1.0	0.4	
450	+	5.5	+ 4.9	— 8.9	950	— 5.2	— 4.8	+ 8.7	Dec.	—	—	
460	+	4.4	4.0	9.0	960	4.2	3.9	8.9	7	0.6	0.5	
470	+	3.3	3.0	9.2	970	3.2	2.9	9.0	17	— 0.2	0.5	
480	+	2.2	2.1	9.3	980	2.1	2.0	9.1	27	+ 0.3	0.5	
490	+	1.1	1.0	9.3	990	1.0	1.0	9.1	37	+ 0.6	— 0.5	
500	+	0.0	+ 0.0	— 9.3	1000	— 0.0	— 0.0	+ 9.2				



*Perturbations produced by Venus.*

Arguments II and III.

III.

II.	0	10	20	30	40	50	60	70	80	90	100	110	120
	"	"	"	"	"	"	"	"	"	"	"	"	"
0	21.6	20.8	19.8	19.0	17.9	16.8	15.9	14.7	14.0	13.2	12.8	12.5	12.3
20	23.1	22.7	21.6	21.0	20.1	19.3	18.4	17.4	16.4	15.5	14.5	13.8	13.4
40	23.5	23.2	22.9	22.7	22.0	21.1	20.4	19.5	18.7	17.9	16.9	16.1	15.3
60	22.2	22.5	23.1	22.7	22.8	22.5	21.9	21.3	20.5	19.9	19.1	18.2	17.4
80	20.0	20.7	21.4	21.7	22.1	22.3	22.2	22.2	21.7	21.3	20.7	19.9	19.3
100	17.6	18.6	19.2	19.9	20.5	21.0	21.6	21.7	21.6	21.6	21.5	21.1	20.5
120	15.3	16.0	16.9	17.7	18.4	19.2	19.8	20.2	20.7	20.8	21.1	21.1	20.8
140	13.6	14.2	14.8	15.5	16.2	17.0	17.6	18.3	19.0	19.4	20.0	20.0	20.4
160	12.7	13.2	13.6	14.1	14.6	15.0	15.7	16.4	17.0	17.3	18.1	18.7	19.2
180	12.7	12.9	13.1	13.5	13.9	14.0	14.5	14.8	15.0	15.8	16.4	16.8	17.2
200	13.2	13.2	13.2	13.4	13.7	13.8	14.1	14.2	14.5	14.5	14.8	15.2	16.0
220	13.5	13.6	13.9	14.1	14.1	14.1	14.2	14.3	14.5	14.6	14.6	14.7	14.8
240	13.6	13.8	14.1	14.4	14.6	14.8	14.8	14.9	15.1	15.1	15.1	14.9	14.8
260	12.8	13.3	13.8	14.2	14.6	15.0	15.3	15.6	15.5	15.5	15.6	15.6	15.6
280	11.5	12.3	13.0	13.4	14.0	14.6	15.1	15.4	16.0	16.2	16.2	16.3	16.2
300	10.1	10.9	11.3	12.1	12.9	13.7	14.2	14.9	15.4	16.0	16.4	16.5	16.7
320	8.2	8.8	9.6	10.6	11.3	12.0	12.9	13.7	14.3	15.0	15.8	16.3	16.8
340	6.9	7.5	8.1	8.4	9.4	10.1	11.1	11.9	12.7	13.6	14.4	15.2	16.0
360	6.5	6.5	6.8	7.4	8.0	8.4	9.1	9.9	10.8	11.5	12.6	13.4	14.4
380	6.8	6.5	6.3	6.4	6.7	7.0	7.6	8.2	8.9	9.6	10.6	11.4	12.4
400	7.5	7.1	6.7	6.4	6.2	6.4	6.5	6.9	7.5	7.9	8.7	9.4	10.3
420	9.1	8.4	7.6	7.1	6.7	6.5	6.3	6.2	6.7	6.8	7.2	7.8	8.4
440	10.6	9.8	9.0	8.6	7.9	7.2	6.7	6.4	6.4	6.4	6.6	6.8	7.1
460	12.1	11.5	10.5	9.6	9.0	8.5	8.0	7.3	6.8	6.6	6.5	6.4	6.5
480	13.6	12.8	11.9	11.0	10.4	9.6	8.8	8.2	7.7	7.2	6.8	6.4	6.5
500	15.1	14.4	13.4	12.4	11.6	10.8	10.1	9.3	8.6	8.1	7.5	7.1	6.8
520	16.5	15.6	14.8	13.9	13.1	12.3	11.3	10.5	9.7	9.1	8.6	7.9	7.4
540	18.1	17.5	16.4	15.5	14.5	13.7	12.8	11.8	11.1	10.4	9.7	8.9	8.2
560	20.4	19.3	18.2	17.6	16.5	15.4	14.4	13.4	12.7	11.6	10.8	10.2	9.2
580	22.8	21.7	20.7	19.7	18.4	17.6	16.6	15.5	14.3	13.4	12.5	11.6	10.6
600	25.2	24.1	23.1	22.2	21.2	19.9	18.6	17.8	16.6	15.6	14.5	13.4	12.6
620	27.3	26.5	25.6	24.7	23.5	22.5	21.6	20.4	19.0	18.1	16.8	15.7	14.7
640	29.0	28.5	27.7	26.9	26.2	25.1	24.1	22.9	21.8	20.8	19.6	18.4	17.2
660	29.8	29.6	29.2	28.5	28.1	27.4	26.5	25.6	24.5	23.4	22.5	21.2	19.8
680	29.7	29.6	29.5	29.5	29.1	28.8	28.2	27.6	27.0	26.0	25.0	23.8	22.8
700	28.8	29.2	29.3	29.5	29.5	29.5	29.2	28.8	28.4	27.8	27.2	26.4	25.2
720	26.9	27.6	28.3	29.0	29.2	29.4	29.4	29.3	29.1	28.9	28.4	27.9	27.3
740	24.7	25.7	26.6	27.3	27.9	28.5	29.1	29.0	29.2	29.3	29.1	28.8	28.4
760	22.2	23.5	24.3	25.3	26.2	27.0	27.6	28.3	28.6	28.7	28.9	29.1	29.0
780	19.6	21.0	22.0	23.2	24.2	25.1	25.9	26.7	27.3	27.8	28.4	28.5	28.7
800	17.2	18.5	19.3	20.9	21.8	22.9	23.9	25.0	25.8	26.4	26.9	27.6	28.1
820	15.2	15.9	17.0	18.4	18.9	20.7	21.7	22.8	23.8	24.8	25.6	26.2	26.6
840	13.2	14.0	15.0	16.0	17.0	18.2	18.8	20.3	21.7	22.7	23.6	24.5	25.3
860	11.5	12.2	13.0	13.9	14.9	15.9	17.1	18.0	18.9	20.3	21.4	22.6	23.5
880	11.0	11.2	11.5	12.2	13.0	13.7	14.8	15.7	16.8	18.1	19.1	20.2	21.1
900	11.2	10.2	10.9	11.5	12.5	12.1	12.8	13.7	14.5	15.5	16.6	17.9	18.5
920	12.1	11.6	11.5	11.1	11.2	11.3	11.7	12.1	12.7	13.4	14.4	15.2	16.4
940	14.0	13.3	12.6	12.3	11.6	11.5	11.3	11.4	11.6	12.0	12.8	13.3	14.2
960	16.7	15.6	14.6	13.7	13.1	12.5	11.9	11.7	11.6	11.4	11.7	12.1	12.6
980	19.5	18.3	17.3	16.4	15.2	14.2	13.4	12.7	12.2	12.0	11.9	11.8	11.8
1000	21.6	20.8	19.8	19.0	17.9	16.8	15.9	14.7	14.0	13.2	12.8	12.5	12.2
	0	10	20	30	40	50	60	70	80	90	00	110	120



*Perturbations produced by Venus.*

Arguments II and III.

III.

II.	120	130	140	150	160	170	180	190	200	210	220	230	240
0	12.2	12.2	12.3	12.4	12.8	13.3	13.9	14.7	15.6	16.5	17.7	18.8	20.1
20	13.4	12.9	12.6	12.3	12.2	12.4	12.9	13.3	14.0	14.6	15.5	16.4	17.3
40	15.3	14.4	14.0	13.5	13.0	12.9	12.6	12.6	13.1	13.5	14.0	14.4	15.4
60	17.4	16.7	16.0	15.2	14.5	14.0	13.6	13.3	13.2	13.2	13.4	13.5	14.1
80	19.3	18.7	17.7	17.1	16.4	15.9	15.4	14.6	14.3	13.9	13.8	13.7	13.6
100	20.5	20.2	19.5	18.9	18.2	17.5	17.1	16.3	15.9	15.4	14.8	14.6	14.3
120	20.8	20.7	20.4	20.0	19.7	19.2	18.5	18.0	17.3	16.9	16.5	16.2	15.6
140	20.4	20.4	20.2	20.0	20.1	19.7	19.5	19.3	18.8	18.2	17.7	17.4	17.0
160	19.2	19.1	19.4	19.7	19.5	19.6	19.3	19.6	19.2	19.0	18.7	18.4	18.1
180	17.2	17.7	18.5	18.5	18.5	18.8	18.4	18.8	19.0	19.0	18.9	18.6	18.5
200	16.0	16.2	16.6	16.8	17.5	17.6	17.7	17.9	18.1	18.2	18.3	18.3	18.3
220	14.8	15.0	15.3	15.7	16.1	16.2	16.6	16.8	17.1	17.5	17.1	17.4	17.5
240	14.8	14.7	14.8	15.0	15.1	15.4	15.7	15.8	16.0	16.1	16.1	16.3	16.4
260	15.6	15.7	15.3	14.8	15.0	15.0	15.1	15.0	15.1	15.2	15.2	15.1	15.3
280	16.2	16.2	16.2	15.9	15.8	15.8	15.5	15.4	15.1	14.9	14.8	14.7	15.0
300	16.7	17.0	17.1	16.9	16.9	16.6	16.5	16.3	15.9	15.7	15.2	14.9	14.8
320	16.8	17.3	17.5	17.6	17.7	17.6	17.5	17.2	17.0	16.8	16.5	16.1	15.6
340	16.0	16.4	17.2	17.8	17.9	18.1	18.3	18.2	18.2	17.9	17.5	17.3	16.8
360	14.4	15.2	16.0	16.7	17.4	18.1	18.4	18.6	18.8	18.8	18.8	18.7	18.4
380	12.4	13.4	14.3	15.3	16.1	16.9	17.5	18.1	18.6	19.1	19.3	19.5	19.5
400	10.3	11.2	12.3	13.2	14.2	15.1	16.0	16.8	17.8	18.4	18.8	19.3	19.8
420	8.4	9.2	10.0	11.0	12.2	13.0	14.1	15.0	15.9	16.9	17.7	18.5	19.0
440	7.1	7.6	8.4	9.0	9.9	10.9	11.8	12.9	13.8	14.9	16.0	16.7	17.8
460	6.5	6.8	7.2	7.4	8.1	9.0	9.7	10.6	11.7	12.6	13.8	14.6	15.9
480	6.5	6.5	6.4	6.6	7.0	7.5	8.2	8.8	9.6	10.4	11.5	12.5	13.5
500	6.8	6.7	6.5	6.3	6.5	6.6	7.0	7.4	8.2	8.6	9.4	10.4	11.3
520	7.4	7.0	6.8	6.5	6.3	6.1	6.3	6.6	7.0	7.5	8.0	8.8	9.3
540	8.2	7.6	7.2	6.8	6.5	6.3	6.2	6.0	6.2	6.5	6.9	7.4	7.9
560	9.2	8.6	7.9	7.5	6.8	6.6	6.3	6.1	6.0	6.1	6.2	6.5	6.9
580	10.6	9.8	9.1	8.4	7.7	7.3	6.6	6.3	6.1	5.9	5.7	5.9	6.0
600	12.6	11.4	10.5	9.5	8.7	8.1	7.4	7.0	6.4	6.1	5.8	5.5	5.6
620	14.7	13.5	12.4	11.4	10.4	9.5	8.7	7.9	7.3	6.7	6.2	5.6	5.2
640	17.2	16.2	14.9	13.7	12.5	11.4	10.4	9.5	8.7	7.8	7.0	6.5	5.9
660	19.8	19.0	17.6	16.5	15.1	13.9	12.8	11.5	10.5	9.6	8.6	7.7	6.9
680	22.8	21.7	20.4	19.3	18.1	16.8	15.7	14.2	13.0	11.9	10.7	9.6	8.6
700	25.2	24.3	23.3	22.1	20.7	19.7	18.5	17.3	16.0	14.3	13.4	12.1	11.0
720	27.3	26.4	25.7	24.5	23.7	22.5	21.1	20.2	18.8	17.7	16.4	15.3	13.9
740	28.4	27.7	27.4	26.6	25.9	24.9	24.0	22.8	21.5	20.6	19.2	18.1	16.8
760	29.0	28.7	28.3	27.8	27.3	26.8	25.9	25.2	24.3	23.0	21.7	20.7	19.7
780	28.7	28.7	28.8	28.7	28.3	28.0	27.2	26.1	26.1	25.2	24.3	23.3	22.2
800	28.1	28.3	28.4	28.5	28.5	28.4	28.2	27.3	27.3	26.7	25.9	25.1	24.4
820	26.6	27.3	27.8	28.1	28.3	28.1	28.1	28.0	27.9	27.7	27.2	26.5	25.9
840	25.3	26.2	26.7	27.2	27.5	27.9	28.1	28.1	27.9	27.9	27.6	27.3	27.2
860	23.5	24.5	25.1	25.9	26.6	27.1	27.4	27.7	27.9	28.0	27.9	27.7	27.5
880	21.1	22.4	23.3	24.2	25.1	25.8	26.5	27.0	27.3	27.5	27.8	28.0	27.7
900	18.5	20.1	21.3	22.1	23.1	24.7	25.0	25.7	26.3	26.9	27.3	27.5	27.6
920	16.4	17.7	18.4	20.0	21.0	22.2	23.0	23.9	24.9	25.7	26.2	26.9	27.3
940	14.2	14.9	16.1	17.5	18.2	19.6	20.8	21.9	23.0	23.9	24.7	25.7	26.1
960	12.6	13.3	14.1	14.4	15.9	17.2	17.9	19.5	20.5	21.7	22.7	23.9	24.7
980	11.8	12.1	12.7	13.3	14.1	14.8	15.6	16.8	17.6	19.3	20.2	21.4	22.6
1000	12.2	12.2	12.3	12.4	12.8	13.3	13.9	14.7	15.6	16.5	17.6	18.8	20.1
	120	130	140	150	160	170	180	190	200	210	220	230	240



*Perturbations produced by Venus.*

Arguments II. and III.

III.

II.	240	250	260	270	280	290	300	310	320	330	340	350	360
	"	"	"	"	"	"	"	"	"	"	"	"	"
0	20.1	21.1	22.2	23.4	24.3	25.2	25.8	26.6	27.2	27.6	27.7	27.6	27.6
20	17.3	18.6	19.7	20.9	21.9	23.0	24.2	24.9	25.8	26.6	27.0	27.4	27.7
40	15.4	16.5	17.3	18.3	19.4	20.5	21.6	22.7	23.7	24.9	25.5	26.3	26.9
60	14.1	14.6	15.2	16.3	17.2	18.1	18.9	20.3	21.2	22.3	23.4	24.5	25.3
80	13.6	14.0	14.5	14.9	15.5	16.3	17.3	18.2	19.0	20.0	21.1	22.0	23.1
100	14.3	14.3	14.3	14.4	14.6	15.0	15.5	16.2	16.9	17.7	18.9	19.8	20.8
120	15.6	15.2	14.8	14.8	15.0	14.9	15.0	15.2	15.9	16.3	17.0	17.7	18.5
140	17.0	16.6	16.4	15.8	15.5	15.4	15.6	15.6	15.5	15.6	16.1	16.7	17.1
160	18.1	17.7	17.5	17.3	16.9	16.6	16.3	15.9	16.1	16.3	16.3	16.2	16.5
180	18.5	18.5	18.3	18.1	17.9	17.6	17.5	17.3	17.0	16.9	16.7	16.8	16.9
200	18.3	18.4	18.2	18.2	18.2	18.2	18.1	18.1	17.8	17.7	17.6	17.5	17.7
220	17.5	17.6	17.8	17.8	18.0	18.0	18.2	18.1	18.1	18.3	18.4	18.3	18.3
240	16.4	16.5	16.7	16.9	17.1	17.3	17.3	17.7	17.5	18.0	18.3	18.4	18.6
260	15.3	15.5	15.5	15.6	15.8	16.1	16.4	16.6	16.8	16.9	17.4	17.7	18.2
280	15.0	14.9	14.9	14.9	14.9	14.7	15.0	15.3	15.5	15.9	16.1	16.4	16.8
300	14.8	14.6	14.6	14.2	14.0	14.0	13.9	13.9	14.2	14.5	14.8	15.0	15.5
320	15.6	15.3	14.7	14.5	14.4	13.1	13.6	13.4	13.3	13.1	13.4	13.6	13.8
340	16.8	16.6	16.0	15.5	15.2	14.5	14.3	13.7	13.1	13.0	12.7	12.6	12.6
360	18.4	17.9	17.5	17.0	16.5	15.9	15.4	14.9	14.3	13.7	13.0	12.6	12.3
380	19.5	19.2	18.9	18.5	17.9	17.7	16.9	16.4	15.8	15.0	14.5	13.6	13.1
400	19.8	19.8	20.1	19.7	19.4	19.1	18.6	18.1	17.5	17.0	16.1	15.2	14.8
420	19.0	19.5	20.0	20.3	20.3	20.3	20.1	19.4	19.0	18.9	18.1	17.3	16.5
440	17.8	18.7	19.2	19.7	20.1	20.4	20.7	20.7	20.5	20.2	19.8	19.5	18.6
460	15.9	16.8	17.6	18.6	19.2	19.9	20.3	20.6	21.0	20.9	20.9	20.8	20.3
480	13.5	14.6	15.5	16.6	17.7	18.5	19.3	19.9	20.5	20.8	21.1	21.2	21.2
500	11.3	12.4	13.4	14.4	15.5	15.5	17.7	18.6	19.1	19.9	20.7	21.0	21.4
520	9.3	10.2	11.2	12.2	13.3	14.2	15.4	16.4	17.6	18.4	19.2	19.8	20.6
540	7.9	8.6	9.4	10.1	11.1	12.1	13.1	14.2	15.3	16.3	17.4	18.3	19.2
560	6.9	7.2	7.8	8.4	9.2	10.1	11.0	11.9	13.1	14.1	15.2	16.2	17.2
580	6.0	6.3	6.6	7.0	7.6	8.4	9.1	9.9	10.9	11.9	12.9	14.1	15.0
600	5.6	5.6	5.8	6.1	6.5	6.8	7.4	8.1	8.8	9.9	10.7	11.8	12.8
620	5.2	5.4	5.3	5.3	5.5	5.9	6.3	6.6	7.2	8.0	8.7	9.5	10.6
640	5.9	5.6	5.2	4.9	5.0	5.0	5.2	5.5	5.8	6.4	7.0	7.6	8.5
660	6.9	6.3	5.7	5.4	5.0	4.8	4.5	4.7	4.9	5.1	5.5	6.0	6.8
680	8.6	7.6	6.9	6.2	5.6	5.1	4.8	4.6	4.2	4.2	4.5	4.6	5.1
700	11.0	10.0	8.7	7.8	6.8	6.3	5.6	5.0	4.6	4.2	4.2	4.0	4.2
720	13.9	12.5	11.2	10.3	9.1	7.9	7.1	6.2	5.6	4.8	4.5	4.2	3.8
740	16.8	15.5	14.4	13.0	11.7	10.5	9.4	8.4	7.2	6.5	5.6	5.0	4.3
760	19.7	18.5	17.2	15.9	14.7	13.5	12.2	10.8	9.8	8.9	7.6	6.7	5.9
780	22.2	21.2	20.1	19.0	17.6	16.3	15.1	14.0	12.6	11.6	10.2	9.2	8.1
800	24.4	23.4	22.2	21.3	20.3	19.2	18.0	16.7	15.4	14.3	13.2	11.9	10.8
820	25.9	25.1	24.4	23.3	22.3	21.6	20.4	19.4	18.2	17.2	15.9	14.6	13.6
840	27.2	26.6	25.8	25.0	24.3	23.5	22.4	21.6	20.5	19.4	18.4	17.3	16.4
860	27.5	27.1	26.8	26.4	25.5	24.8	24.3	23.3	22.2	21.5	20.5	19.6	18.4
880	27.7	27.5	27.2	27.0	26.5	26.0	25.5	24.7	24.1	23.2	22.0	21.4	20.4
900	27.6	27.8	27.9	27.6	27.1	26.7	26.5	25.7	25.3	24.6	23.9	23.0	22.0
920	27.3	27.5	27.5	27.6	27.7	27.5	27.2	26.7	26.3	25.7	25.1	24.3	23.6
940	26.1	26.7	27.2	27.4	27.7	27.7	27.6	27.5	27.1	26.6	26.2	25.6	25.5
960	24.7	25.4	26.2	26.6	27.2	27.5	27.7	27.7	27.6	27.4	27.1	27.0	26.2
980	22.6	23.7	24.6	25.3	25.9	26.8	27.2	27.5	27.7	27.8	27.6	27.5	27.1
1000	20.1	21.1	22.2	23.4	24.3	25.2	25.8	26.6	27.2	27.6	27.7	27.6	27.6
	240	250	260	270	280	290	300	310	320	330	340	350	360

*Perturbations produced by Venus.*

## Arguments II. and III.

## III.

II.	360	370	380	390	400	410	420	430	440	450	460	470	480
	"	"	"	"	"	"	"	"	"	"	"	"	"
0	27.6	27.7	27.3	26.7	26.2	25.5	24.7	23.8	23.1	22.3	21.3	20.2	19.3
20	27.7	27.8	27.8	27.6	27.4	26.8	26.2	25.6	24.8	24.0	23.1	22.0	20.9
40	26.9	27.3	27.6	27.9	27.9	27.7	27.5	27.1	26.3	25.6	24.9	24.0	23.2
60	25.3	26.0	26.8	27.1	27.5	27.9	27.8	27.7	27.3	27.1	26.7	25.9	25.0
80	23.1	24.0	25.1	25.9	26.5	27.3	27.5	27.9	28.2	28.0	27.6	27.5	27.2
100	20.8	21.8	22.6	23.6	24.6	25.5	26.2	26.7	27.2	27.5	27.6	27.8	27.4
120	18.5	19.6	20.6	21.5	22.4	23.2	24.1	25.1	25.8	26.4	26.9	27.3	27.5
140	17.1	17.9	18.6	19.3	20.3	21.3	22.0	22.9	23.7	24.7	25.5	26.0	26.7
160	16.5	17.1	17.4	18.1	18.8	19.3	20.1	21.0	21.9	22.6	23.5	24.2	25.1
180	16.9	17.0	17.1	17.4	18.0	18.4	18.9	19.4	20.1	20.7	21.2	22.2	23.0
200	17.7	17.5	17.7	17.7	17.6	18.1	18.3	18.7	19.2	19.7	20.1	20.8	21.5
220	18.3	18.2	18.3	18.3	18.3	18.3	18.6	18.7	18.9	19.3	19.5	20.0	20.4
240	18.6	18.8	18.9	18.9	18.9	19.0	19.2	19.1	19.2	19.5	19.6	19.7	19.9
260	18.2	18.5	18.7	18.8	19.0	19.3	19.5	19.6	19.9	19.9	20.0	20.1	20.2
280	16.8	17.4	17.9	18.3	18.7	19.1	19.3	19.8	20.0	20.2	20.4	20.6	20.8
300	15.5	15.8	16.2	16.6	17.6	18.1	18.5	19.2	19.4	19.9	20.6	20.8	20.9
320	13.8	14.2	14.6	15.1	15.6	16.2	16.8	17.7	18.3	18.9	19.5	20.1	20.8
340	12.6	12.9	13.0	13.3	13.7	14.4	14.9	15.5	16.2	17.1	18.0	18.6	19.4
360	12.3	12.1	11.9	12.0	12.3	12.5	13.0	13.4	14.2	14.9	15.7	16.5	17.3
380	13.1	12.5	11.9	11.6	11.5	11.4	11.6	11.7	12.3	12.7	13.3	14.0	15.0
400	14.8	13.9	13.1	12.5	11.7	11.2	11.1	10.9	11.0	11.1	11.4	12.0	12.6
420	16.5	15.7	15.1	14.3	13.4	12.5	11.7	11.1	10.8	10.8	10.5	10.6	10.7
440	18.6	17.9	17.1	16.1	15.6	14.4	13.5	12.8	11.9	11.1	10.6	10.3	10.3
460	20.3	19.8	19.3	18.5	17.6	16.8	15.9	14.7	13.7	12.9	12.0	11.1	10.9
480	21.2	21.1	20.8	20.3	19.7	19.1	18.3	17.4	16.4	15.0	14.1	13.2	12.2
500	21.4	21.4	21.4	21.3	21.1	20.8	20.0	19.5	18.8	17.8	17.0	15.7	14.4
520	20.6	21.2	21.7	21.7	21.5	21.5	21.4	21.1	20.5	19.8	19.1	18.2	17.6
540	19.2	20.0	20.7	21.1	21.8	22.0	21.8	21.7	21.5	21.2	20.9	20.3	19.6
560	17.2	18.4	19.0	20.0	20.8	21.1	22.7	21.9	22.2	22.1	21.9	21.7	21.1
580	15.0	16.0	17.3	18.2	19.1	19.9	20.8	21.1	21.7	22.0	22.2	22.3	22.1
600	12.8	13.9	15.1	15.9	17.2	18.0	19.0	19.9	20.6	21.3	21.8	22.0	22.4
620	10.6	11.5	12.7	13.7	14.9	16.0	17.1	18.3	19.1	19.9	20.8	21.3	22.0
640	8.5	9.5	10.4	11.3	12.3	13.7	14.9	16.0	17.1	18.1	19.0	19.9	20.7
660	6.8	7.4	8.2	9.1	10.1	11.1	12.2	13.6	14.6	15.8	17.1	18.1	19.0
680	5.1	5.7	6.4	7.1	7.9	8.7	9.7	11.0	12.1	13.1	14.1	15.7	16.8
700	4.2	4.4	4.7	5.1	5.8	6.7	7.4	8.4	9.4	10.6	11.5	13.0	14.1
720	3.8	3.8	3.8	4.0	4.4	4.8	5.4	5.9	6.9	8.0	9.1	10.1	11.5
740	4.3	3.9	3.8	3.7	3.6	3.8	3.9	4.4	4.9	5.7	6.4	7.4	8.9
760	5.9	5.1	4.4	4.0	3.6	3.4	3.4	3.5	3.9	4.3	4.7	5.2	5.9
780	8.1	7.1	6.1	5.3	4.6	4.1	3.7	3.3	3.3	3.1	3.4	3.6	4.1
800	10.8	9.7	8.5	7.5	6.5	5.6	4.9	4.2	3.8	3.4	3.2	3.1	3.1
820	13.6	12.5	11.2	10.1	9.0	8.0	6.9	6.1	5.3	4.7	3.9	3.7	3.1
840	16.4	15.1	13.7	12.9	11.7	10.6	9.5	8.6	7.5	6.6	5.7	4.9	4.4
860	18.4	17.5	16.6	15.4	14.3	13.1	12.1	11.1	10.0	9.1	7.9	7.0	6.3
880	20.4	19.6	18.7	17.5	16.6	15.6	14.5	13.6	12.5	11.5	10.4	9.5	8.6
900	22.0	21.1	20.2	19.4	18.7	17.7	16.5	15.7	14.7	13.8	12.5	11.9	10.9
920	23.6	22.7	21.7	21.1	20.1	19.4	18.4	17.5	16.7	15.6	14.8	13.9	13.1
940	25.5	24.1	23.4	22.4	21.4	20.6	19.9	19.0	18.2	17.3	16.6	15.7	14.8
960	26.2	25.6	24.7	24.1	23.3	22.3	21.3	20.6	19.0	18.9	17.9	17.1	16.3
980	27.1	26.7	26.3	25.5	24.9	23.8	23.4	22.2	21.0	20.4	19.4	18.6	17.7
1000	27.6	27.7	27.3	26.7	26.2	25.5	24.7	23.8	23.1	22.3	21.3	20.2	19.3
	360	370	380	390	400	410	420	430	440	450	460	470	480

*Perturbations produced by Venus.*

## Arguments II and III.

## III.

II.	480	490	500	510	520	530	540	550	560	570	580	590	600
	"	"	"	"	"	"	"	"	"	"	"	"	"
0	19.3	18.3	17.4	16.6	15.7	15.0	14.2	13.6	13.1	12.3	11.7	11.3	10.8
20	20.9	20.2	19.1	18.2	17.1	16.2	15.5	14.7	14.1	13.3	12.7	12.2	11.5
40	23.2	22.0	20.8	20.1	18.9	17.9	17.1	15.9	15.1	14.4	13.7	13.0	12.2
60	25.0	24.0	23.2	22.0	20.7	19.9	18.9	17.7	16.8	15.8	14.9	14.0	13.2
80	27.2	26.4	25.6	24.1	23.2	22.1	20.8	20.0	18.7	17.9	16.6	15.6	14.8
100	27.4	27.2	26.8	26.3	25.4	24.5	23.5	22.2	20.9	20.0	18.6	17.6	16.6
120	27.5	27.5	27.6	27.1	26.8	26.3	25.4	24.6	23.7	22.4	21.0	20.1	18.8
140	26.7	27.0	27.2	27.4	27.3	27.4	26.9	26.2	25.4	24.6	23.9	22.6	21.1
160	25.1	25.6	26.1	26.7	26.9	27.3	27.1	27.0	26.9	26.4	25.5	24.7	23.9
180	23.0	23.8	24.5	25.0	25.7	26.3	26.7	26.8	27.0	26.8	26.6	26.2	25.6
200	21.5	22.2	22.8	23.5	24.1	24.7	25.5	25.8	26.3	26.6	26.6	26.6	26.4
220	20.4	21.0	21.5	22.0	22.6	23.2	23.8	24.5	25.0	25.4	25.8	26.0	26.2
240	19.9	20.4	20.8	21.2	21.6	21.8	22.2	22.6	23.1	23.3	23.9	24.2	24.6
260	20.2	20.3	20.6	21.2	21.4	21.7	21.9	22.2	22.3	22.7	23.1	23.3	23.6
280	20.8	20.8	21.0	21.1	21.3	21.4	21.5	21.8	22.0	22.2	22.7	23.0	23.3
300	20.9	21.0	21.5	21.7	21.7	22.0	22.0	22.1	22.1	22.2	22.4	22.6	22.8
320	20.8	21.2	21.5	21.6	22.0	22.3	22.5	22.5	22.6	22.7	22.8	22.8	22.9
340	19.4	20.2	20.8	21.5	21.9	22.1	22.6	23.0	23.2	23.4	23.3	23.4	23.5
360	17.3	18.4	19.5	20.0	20.6	21.5	22.2	22.7	23.0	23.7	23.7	24.0	24.2
380	15.0	15.9	16.9	17.8	18.6	19.6	20.6	21.5	22.3	22.9	23.5	23.9	24.5
400	12.6	13.2	14.2	15.4	16.2	17.3	18.1	19.2	20.3	21.4	22.4	23.0	23.7
420	10.7	11.2	12.0	12.5	13.5	14.5	15.6	16.7	17.7	18.7	20.1	21.0	22.0
440	10.3	10.2	10.3	10.5	11.3	12.0	12.9	13.6	14.7	16.0	17.0	18.3	19.5
460	10.9	10.1	9.9	9.9	9.9	10.1	10.7	11.3	12.2	13.0	14.0	15.1	16.5
480	12.2	11.4	10.7	10.1	9.7	9.5	9.7	9.9	10.2	10.7	11.7	12.5	13.4
500	14.4	13.6	12.5	11.6	10.9	10.2	9.8	9.4	9.3	9.6	9.8	10.2	11.1
520	17.6	16.2	15.1	13.9	12.9	11.9	10.9	10.3	9.8	9.5	9.2	9.2	9.6
540	19.6	18.6	18.0	16.7	15.4	14.5	12.2	12.3	11.3	10.5	10.1	9.5	9.3
560	21.1	20.4	19.8	19.0	18.2	17.2	16.0	14.8	13.7	12.7	11.7	10.9	10.2
580	22.1	21.8	21.5	20.9	20.3	19.3	18.6	17.3	16.5	15.4	14.0	12.9	12.2
600	22.4	22.4	22.2	22.2	21.5	21.2	20.6	19.5	19.1	17.7	16.8	15.8	14.4
620	22.0	22.3	22.4	22.4	22.3	22.3	21.9	21.5	20.9	20.0	19.3	18.0	16.9
640	20.7	21.7	22.0	22.3	22.6	22.5	22.6	22.4	22.0	21.6	21.1	20.3	19.6
660	19.0	20.0	20.8	21.3	22.1	22.3	22.6	22.8	22.7	22.6	22.2	21.8	21.3
680	16.8	18.0	19.0	19.9	20.8	21.5	22.1	22.6	22.7	23.0	23.0	22.8	22.4
700	14.1	15.2	16.8	17.9	18.8	20.0	22.1	21.5	22.2	22.6	22.9	23.0	23.2
720	11.5	12.7	13.9	15.0	16.4	17.9	18.6	19.7	20.8	21.6	22.3	22.7	23.0
740	8.9	9.8	10.9	12.2	13.6	14.8	16.2	17.5	18.7	19.5	20.6	21.6	22.3
760	5.9	6.8	8.0	9.3	10.3	11.8	13.2	14.5	15.9	17.4	18.2	19.5	20.5
780	4.1	4.9	5.6	6.4	7.5	8.6	9.9	11.1	12.6	14.0	15.6	16.8	18.1
800	3.1	3.3	4.4	4.8	5.5	6.1	6.9	7.9	9.4	10.7	12.1	13.4	14.9
820	3.1	3.1	3.2	3.1	3.6	3.9	4.8	5.7	6.5	7.5	8.7	10.0	11.5
840	4.4	3.7	3.5	3.2	3.2	3.1	3.4	3.7	4.1	5.0	6.2	7.0	8.2
860	6.3	5.5	4.6	4.1	3.6	3.4	3.3	3.2	3.4	3.4	4.0	4.5	5.6
880	8.6	7.6	6.7	5.9	5.2	4.5	4.1	3.8	3.5	3.4	3.4	3.6	3.9
900	10.9	10.0	9.1	8.3	7.2	6.5	5.8	5.1	4.4	4.2	3.8	3.6	3.6
920	13.1	12.1	11.2	10.3	9.6	8.7	7.7	6.9	6.3	5.8	5.1	4.6	4.2
940	14.8	14.1	13.1	12.4	11.5	10.8	9.8	9.1	8.3	7.6	6.8	6.5	5.9
960	16.3	15.4	14.6	14.0	13.2	12.6	11.7	11.0	10.1	9.6	8.8	8.1	7.5
980	17.7	16.8	16.2	15.2	14.5	13.9	13.1	12.5	11.8	11.2	10.5	9.7	9.3
1000	19.3	18.3	17.4	16.6	15.7	15.0	14.2	13.6	13.1	12.3	11.7	11.3	10.8
	480	490	500	510	520	530	540	550	560	570	580	590	600

*Perturbations produced by Venus.*

## Arguments II. and III.

## III.

II.	600	610	620	630	640	650	660	670	680	690	700	710	720
	"	"	"	"	"	"	"	"	"	"	"	"	"
0	10.8	10.2	9.5	9.1	8.4	7.9	7.4	7.0	6.6	6.3	5.9	5.5	5.4
20	11.5	11.3	10.7	10.4	9.8	9.4	8.9	8.5	7.9	7.7	7.3	6.7	6.6
40	12.3	12.0	11.5	11.0	10.7	10.3	10.0	9.6	9.3	8.9	8.5	8.1	7.8
60	13.3	12.7	12.1	11.6	11.2	10.9	10.5	10.2	10.0	9.8	9.5	9.2	8.9
80	14.3	13.6	12.9	12.4	11.8	11.3	10.9	10.7	10.3	9.9	9.8	9.8	9.6
100	16.6	15.4	14.4	13.4	12.6	12.1	11.5	11.0	10.6	10.2	10.0	9.9	9.6
120	18.8	17.7	16.4	15.3	14.3	13.2	12.4	11.6	11.2	10.6	10.1	10.1	9.6
140	21.1	20.1	18.9	17.7	16.5	15.2	14.2	13.0	12.3	11.6	11.1	10.3	9.9
160	23.9	22.9	21.5	20.4	19.2	17.9	16.6	15.3	14.1	13.1	12.0	11.2	10.5
180	25.6	24.8	23.9	22.9	21.6	20.6	19.1	18.0	16.7	15.5	14.3	12.9	12.0
200	26.4	26.0	25.6	24.9	24.0	22.9	21.7	20.8	19.3	18.1	16.9	15.5	14.4
220	26.2	26.3	26.1	25.8	25.3	24.9	24.1	23.1	21.2	20.9	19.7	18.3	17.1
240	24.6	25.1	25.1	25.3	25.2	25.1	24.7	24.3	24.0	23.0	21.9	21.3	20.2
260	23.6	23.9	24.2	24.5	24.7	24.8	24.9	24.6	24.3	23.8	23.4	22.9	21.6
280	23.3	23.6	23.9	24.2	24.7	24.8	25.0	24.9	24.9	24.8	24.4	24.0	23.5
300	22.8	23.0	23.3	23.4	23.8	24.0	24.1	24.5	24.5	24.6	24.5	24.4	24.0
320	22.9	23.0	23.1	23.2	23.4	23.3	23.6	23.8	24.0	23.9	24.2	24.2	24.2
340	23.5	23.5	23.5	23.4	23.5	23.6	23.6	23.5	23.5	23.6	23.9	23.8	23.8
360	24.2	24.2	24.3	24.2	24.2	24.0	23.7	23.9	24.0	23.7	23.7	23.6	23.6
380	24.5	24.6	24.8	25.1	24.8	24.9	25.0	24.9	24.6	24.5	24.5	24.3	24.0
400	23.7	24.3	24.7	25.0	25.4	25.7	25.7	25.5	25.5	25.4	25.2	24.8	24.6
420	22.0	23.0	23.7	24.6	25.0	25.7	26.1	26.2	26.3	26.5	26.2	26.0	25.9
440	19.5	20.8	21.7	22.7	23.7	24.6	25.4	26.0	26.5	26.7	26.9	27.0	26.9
460	16.5	17.8	19.0	20.1	21.4	22.3	23.5	24.8	25.4	26.1	26.7	27.1	27.3
480	13.4	14.5	15.6	17.0	18.5	19.7	20.9	22.1	23.2	24.4	25.4	26.2	26.8
500	11.1	12.0	13.0	13.8	14.9	16.3	17.9	19.1	20.5	21.6	22.9	24.2	25.1
520	9.6	9.8	10.5	11.5	12.4	13.4	14.4	15.5	17.1	18.4	19.9	21.2	22.3
540	9.3	9.0	9.2	9.6	10.3	11.0	11.9	12.8	13.9	15.1	16.5	17.9	19.4
560	10.2	9.7	9.3	9.1	9.1	9.4	10.0	10.6	11.5	12.4	13.3	14.5	16.0
580	12.2	11.3	10.4	9.9	9.4	9.0	9.2	9.3	9.7	10.4	11.0	12.0	12.7
600	14.4	13.3	12.5	11.6	10.8	10.1	9.6	9.4	9.1	9.3	9.9	10.0	10.8
620	16.9	16.1	14.9	13.7	12.7	12.0	11.1	10.4	9.8	9.5	9.5	9.3	9.7
640	19.6	18.4	17.4	16.3	15.2	14.2	13.1	12.1	11.3	10.6	10.1	9.6	9.5
660	21.3	20.6	19.9	18.7	17.8	16.7	15.6	14.4	13.4	12.4	11.7	11.0	10.2
680	22.4	22.0	21.5	20.8	20.2	19.0	18.1	17.0	15.8	14.7	13.7	12.8	12.0
700	23.2	23.2	22.6	22.2	21.7	21.0	20.5	19.3	18.3	17.3	16.0	15.0	14.1
720	23.0	23.3	23.2	23.4	23.1	22.4	21.9	21.3	20.8	19.5	18.5	17.6	16.4
740	22.3	22.8	23.2	23.4	23.6	23.6	23.3	22.8	22.2	21.6	21.1	19.9	18.8
760	20.5	21.4	22.5	22.8	23.3	23.7	23.6	23.8	23.5	23.3	22.7	21.8	21.3
780	18.1	19.2	20.4	21.3	22.3	23.0	23.3	23.7	23.8	24.0	23.8	23.5	23.0
800	14.9	16.4	17.7	19.1	20.1	21.2	21.1	22.9	23.4	23.8	24.1	24.2	23.9
820	11.5	12.9	14.3	15.8	17.8	18.7	20.0	20.9	22.0	22.7	23.5	23.9	24.0
840	8.2	9.5	10.8	12.2	13.8	15.2	16.6	18.1	19.5	20.6	21.7	22.6	23.3
860	5.6	6.8	7.7	8.8	10.2	11.5	13.2	14.7	16.0	17.4	19.0	20.2	21.3
880	3.9	4.4	5.2	6.1	7.2	8.2	9.7	10.9	12.5	14.1	15.4	16.8	18.2
900	3.6	3.6	3.9	4.2	5.0	5.7	6.6	7.8	9.1	10.3	11.8	13.4	14.8
920	4.2	3.8	3.9	3.9	4.0	4.3	4.7	5.4	6.4	7.3	8.6	9.8	11.2
940	5.9	5.1	4.6	4.4	4.2	4.3	4.3	4.3	4.9	5.3	6.3	7.0	8.0
960	7.5	6.9	6.3	5.8	5.3	4.7	4.7	4.6	4.6	4.6	4.9	5.4	6.0
980	9.3	8.7	7.9	7.4	6.8	6.4	6.0	5.6	5.2	5.0	4.9	5.1	5.1
1000	10.8	10.2	9.5	9.1	8.4	7.9	7.4	7.0	6.6	6.3	5.9	5.5	5.4
	600	610	620	630	640	650	660	670	680	690	700	710	720



*Perturbations produced by Venus.*

Arguments II. and III.

III.

II.	720	730	740	750	760	770	780	790	800	810	820	830	840
	"	"	"	"	"	"	"	"	"	"	"	"	"
0	5.4	5.5	5.8	6.0	6.3	6.8	7.6	8.4	9.3	10.4	11.7	12.9	14.3
20	6.6	6.3	6.0	6.1	6.1	6.2	6.5	6.9	7.7	8.3	9.4	10.2	11.2
40	7.8	7.4	7.1	7.0	6.7	6.6	6.8	6.8	6.9	7.2	7.7	8.5	9.3
60	8.9	8.8	8.3	8.1	7.8	7.6	7.4	7.4	7.3	7.4	7.4	7.7	8.3
80	9.6	9.5	9.1	9.1	9.0	8.8	8.4	8.2	8.1	8.1	8.0	8.1	8.2
100	9.6	9.5	9.6	9.5	9.5	9.3	9.3	9.2	9.2	9.0	8.7	8.7	8.7
120	9.6	9.6	9.5	9.3	9.4	9.6	9.6	9.5	9.5	9.6	9.6	9.6	9.6
140	9.9	9.5	9.6	9.4	9.3	9.3	9.0	9.3	9.5	9.8	9.7	9.8	10.0
160	10.5	9.9	9.5	9.1	8.9	9.0	8.9	9.0	9.0	9.0	9.5	9.6	9.9
180	12.0	11.0	10.1	9.7	9.1	8.8	8.7	8.3	8.5	8.7	8.3	9.0	9.1
200	14.4	13.3	12.0	11.0	10.1	9.4	8.9	8.5	8.2	8.0	8.0	8.3	8.5
220	17.1	15.7	14.6	13.2	12.0	10.9	10.2	9.2	8.7	8.3	7.9	7.7	7.7
240	20.2	19.1	17.8	16.5	14.5	13.4	12.2	11.1	10.0	9.4	8.4	8.0	7.7
260	21.6	21.1	20.1	19.2	17.3	15.9	14.6	13.4	12.4	11.3	10.1	9.1	8.6
280	23.5	22.7	21.6	21.0	19.8	18.8	17.3	16.1	15.0	13.5	12.5	11.5	10.2
300	24.0	23.4	23.2	22.4	21.4	20.5	19.8	18.7	17.5	16.1	15.0	13.7	12.4
320	24.2	23.9	23.5	23.1	22.7	22.2	21.2	20.6	19.6	18.6	17.5	16.3	15.1
340	23.8	23.9	23.7	23.5	23.2	22.8	22.3	21.4	20.9	20.5	19.2	18.6	17.4
360	23.6	23.6	23.6	23.3	23.3	23.1	22.9	22.4	22.0	21.4	20.4	19.9	18.9
380	24.0	24.0	23.7	23.5	23.3	23.1	23.1	22.7	22.4	22.2	21.6	20.8	20.0
400	24.6	24.4	24.4	24.0	23.8	23.4	23.2	23.0	22.8	22.4	22.1	21.6	21.3
420	25.9	25.6	25.2	24.8	24.7	24.3	23.9	23.6	23.3	22.9	22.7	22.3	21.7
440	26.9	26.6	26.4	26.2	25.9	25.5	25.2	24.9	24.5	23.8	23.4	23.0	22.8
460	27.3	27.6	27.6	27.4	27.0	26.9	26.5	26.1	25.6	25.0	24.6	24.2	23.7
480	26.8	27.4	27.6	28.0	28.1	28.2	27.7	27.4	27.3	26.6	26.2	25.7	25.1
500	25.1	26.1	26.8	27.5	28.1	28.2	28.6	28.5	28.4	28.3	27.6	27.2	26.7
520	22.3	23.9	24.8	25.9	26.8	27.5	28.1	28.5	28.7	29.0	28.8	28.6	28.4
540	19.4	20.7	22.1	23.4	24.6	25.6	26.5	27.4	28.0	28.7	28.9	29.1	29.2
560	16.0	17.3	18.6	19.9	21.4	22.9	24.1	25.5	26.4	27.3	28.2	28.6	29.2
580	12.7	14.1	15.5	16.8	18.0	19.3	20.9	22.2	23.5	24.9	26.1	27.0	27.8
600	10.8	11.6	12.7	13.6	14.9	16.2	17.5	18.7	20.2	21.8	23.0	24.4	25.5
620	9.7	10.0	10.5	10.7	12.2	13.2	14.4	15.6	17.0	18.3	19.6	21.2	22.6
640	9.5	9.4	9.6	10.1	10.4	11.1	12.0	13.0	14.0	15.2	16.5	17.9	19.2
660	10.2	10.0	9.7	9.5	9.5	9.9	10.4	11.0	11.7	12.7	13.8	14.9	16.2
680	12.0	11.2	10.5	10.0	9.7	9.5	9.6	10.0	10.4	11.0	11.6	12.5	13.8
700	14.1	13.1	12.3	11.3	10.7	10.1	9.7	9.7	9.9	9.9	10.4	10.9	11.5
720	16.4	15.3	14.4	13.3	12.2	11.6	10.9	10.2	10.1	9.9	10.0	10.1	10.4
740	18.8	17.7	16.7	15.6	14.4	13.5	12.4	11.5	11.1	10.7	10.1	10.0	10.3
760	21.3	20.1	19.2	18.1	16.6	15.6	14.7	13.6	12.8	11.9	11.3	10.7	10.3
780	23.0	22.3	21.5	20.5	19.4	18.4	17.2	15.8	14.9	14.0	13.0	12.2	11.3
800	23.9	23.9	23.4	22.6	21.9	20.7	19.8	18.8	17.5	16.2	15.1	14.2	13.4
820	24.0	24.5	24.2	23.9	23.3	22.6	22.3	21.3	20.3	19.4	18.3	17.3	16.2
840	23.3	24.0	24.3	24.5	24.4	24.3	23.8	23.4	22.7	21.7	20.8	19.6	18.3
860	21.3	22.3	23.3	23.9	24.2	24.7	24.5	24.5	24.3	23.6	23.1	21.9	21.0
880	18.2	19.7	20.9	22.0	22.8	23.8	24.1	24.6	24.8	24.7	24.5	24.0	23.5
900	14.8	16.1	17.6	19.0	20.6	21.5	22.5	23.2	24.1	24.5	24.2	24.8	24.5
920	11.2	12.6	14.0	15.5	17.0	18.4	19.9	21.0	22.0	22.9	23.5	24.5	24.5
940	8.0	9.3	10.7	12.0	13.3	14.8	16.4	17.6	19.1	20.4	21.4	22.4	23.2
960	6.0	6.9	7.8	8.6	10.2	11.5	12.7	14.1	15.6	16.9	18.5	19.5	20.7
980	5.1	5.5	6.0	6.7	7.7	8.5	9.7	10.9	12.2	13.6	14.8	16.1	17.6
1000	5.4	5.5	5.8	5.8	6.3	6.8	7.6	8.4	9.3	10.5	11.7	12.9	14.3
	720	730	740	750	760	770	780	790	800	810	820	830	840



*Perturbations produced by Venus.*

## Arguments II. and III.

## III.

II.	810	850	860	870	880	890	900	910	920	930	940	950	960
	"	"	"	"	"	"	"	"	"	"	"	"	"
0	14.3	15.5	16.9	18.2	19.2	20.2	21.4	22.5	23.0	23.5	24.0	24.2	24.2
20	11.2	12.4	13.6	14.9	16.2	17.3	18.6	19.6	20.5	21.5	22.4	23.1	23.6
40	9.3	10.2	10.9	11.8	13.3	14.2	15.5	16.6	17.8	18.8	19.7	20.7	21.6
60	8.3	8.7	9.5	10.1	10.8	11.6	12.7	13.8	14.9	15.9	17.0	18.1	19.1
80	8.2	8.3	8.6	8.9	9.6	10.3	10.7	11.6	12.5	13.3	14.5	15.2	16.2
100	8.7	8.7	8.9	9.0	9.1	9.4	9.9	10.4	11.0	11.7	12.4	12.9	14.0
120	9.6	9.5	9.3	9.6	9.6	9.7	9.9	9.8	10.4	10.9	11.3	11.8	12.3
40	10.0	10.2	10.1	10.2	10.1	10.3	10.4	10.5	10.5	10.6	10.9	11.4	11.5
160	9.9	10.0	10.2	10.4	10.6	11.0	11.0	10.9	11.0	11.3	11.3	11.3	11.6
180	9.1	9.6	9.9	10.1	10.4	10.7	11.0	11.3	11.5	11.7	11.7	11.9	12.2
200	8.5	8.8	9.1	9.5	9.7	10.0	10.5	11.0	11.2	11.6	12.0	12.2	12.4
220	7.7	7.7	8.1	8.4	8.8	9.2	9.7	10.1	10.6	11.0	11.4	11.8	12.3
240	7.7	7.3	7.4	7.4	7.7	8.0	8.4	9.0	9.6	10.0	10.5	11.0	11.5
260	8.6	7.9	7.4	7.2	7.1	7.1	7.3	7.6	8.1	8.5	9.3	10.0	10.4
280	10.2	9.2	8.3	7.9	7.4	7.1	7.0	6.9	7.0	7.3	7.7	8.5	8.9
300	12.4	11.4	10.4	9.3	8.5	7.8	7.4	6.9	6.7	6.8	6.8	7.0	7.5
320	15.1	13.9	12.5	11.4	10.5	9.7	8.6	7.8	7.4	7.0	6.6	6.5	6.7
340	17.4	16.4	15.2	13.9	12.7	11.6	10.6	9.7	8.7	8.0	7.3	6.8	6.6
360	18.9	18.1	17.4	16.3	15.1	13.8	12.8	11.7	10.6	9.8	8.8	8.0	7.4
380	20.0	19.6	18.8	17.7	16.9	13.0	15.1	13.9	12.7	11.8	10.8	9.8	8.9
400	21.3	20.6	19.6	19.4	18.4	17.6	16.5	15.7	14.8	13.7	12.8	11.8	10.9
420	21.7	21.1	20.8	20.3	19.3	18.9	18.2	17.2	16.3	15.3	14.5	13.7	12.6
440	22.8	22.1	21.6	20.8	20.6	19.7	19.0	18.6	17.7	16.6	15.9	15.1	14.2
460	23.7	23.3	22.7	22.0	21.6	20.9	20.2	19.5	18.5	18.1	17.3	16.7	15.7
480	25.1	24.4	23.9	23.3	22.8	22.0	21.4	20.9	20.2	19.3	18.3	17.7	16.9
500	26.7	26.3	25.7	24.9	24.3	23.6	23.0	22.3	21.4	20.7	20.3	19.1	18.1
520	28.4	27.8	27.3	26.8	26.3	25.6	24.7	23.9	23.3	22.6	21.8	20.8	20.1
540	29.2	29.2	28.9	28.5	27.8	27.4	26.8	26.1	25.3	24.4	23.7	23.0	22.0
560	29.2	29.3	29.5	29.6	29.3	29.1	28.8	28.0	27.4	26.9	26.1	25.1	24.3
580	27.8	28.6	29.0	29.4	29.6	29.8	29.8	29.3	28.0	28.7	27.9	27.3	26.6
600	25.5	26.7	27.6	28.4	28.9	29.2	29.6	29.9	29.9	29.8	29.3	29.0	28.5
620	22.6	23.8	25.0	26.2	27.1	27.9	28.8	29.3	29.6	29.8	30.1	29.8	29.6
640	19.2	20.6	21.6	23.3	24.6	25.2	26.6	27.8	28.3	28.9	29.4	29.7	29.9
660	16.2	17.5	18.8	20.2	21.1	22.9	24.0	25.1	26.2	27.1	28.2	28.8	29.2
680	13.8	14.7	15.8	16.9	18.4	19.9	20.6	22.3	23.6	24.9	25.8	26.7	27.5
700	11.5	12.3	13.4	14.6	15.6	16.7	18.0	19.5	20.7	22.0	23.1	24.2	25.1
720	10.4	11.0	11.4	12.3	13.3	14.3	15.6	16.4	17.7	19.3	19.9	21.6	22.6
740	10.3	10.4	10.5	11.0	11.4	12.2	13.3	14.2	15.3	16.5	17.4	18.8	19.5
760	10.3	10.0	10.2	10.3	10.7	11.0	11.5	12.2	13.1	14.2	15.1	16.0	17.3
780	11.3	10.8	10.6	10.2	10.2	10.5	10.7	11.1	11.5	12.3	13.2	14.0	15.0
800	13.4	12.5	11.7	11.0	10.6	10.3	10.3	10.4	10.7	11.0	11.6	11.3	12.2
820	16.2	15.2	14.4	13.5	13.5	11.9	11.4	11.0	10.9	10.8	10.8	11.2	11.4
840	18.3	17.1	16.2	14.9	14.1	13.0	12.4	11.7	11.2	10.7	10.6	11.1	11.2
860	21.0	20.2	18.7	17.7	16.6	15.4	14.3	13.3	12.5	11.9	11.4	11.0	10.9
880	23.5	22.4	21.3	20.4	19.3	18.0	17.0	15.9	14.8	13.7	12.8	12.0	12.6
900	24.5	24.2	23.8	22.7	21.9	19.9	19.7	18.6	17.2	16.4	15.3	14.1	13.3
920	24.5	24.8	24.7	24.3	24.1	23.2	22.3	21.3	20.0	19.3	18.0	16.7	15.7
940	23.2	24.0	24.5	24.6	24.5	24.5	24.2	23.5	22.7	21.8	20.6	19.5	18.4
960	20.7	21.9	22.8	23.6	24.0	24.5	24.5	24.2	24.3	23.7	22.9	22.1	21.0
980	17.6	18.7	20.1	21.2	22.2	23.1	23.6	24.0	24.3	24.3	24.3	23.7	23.0
1000	14.3	15.5	16.9	18.2	19.2	20.2	21.4	22.5	23.0	23.5	24.0	24.2	24.2
	840	850	860	870	880	890	900	910	920	930	940	950	960

*Perturbations by Venus.**Perturbations by Mars.*

Arguments II and III.

Arguments II and IV.

III.

IV.

II.	960	970	980	990	1000	0	10	20	30	40	50	60	70
	"	"	"	"	"	"	"	"	"	"	"	"	"
0	24.2	23.7	23.1	22.5	21.6	9.5	10.2	10.8	11.2	11.5	11.7	11.8	11.5
20	23.6	23.7	24.0	23.4	23.1	8.3	9.1	9.8	10.5	10.9	11.2	11.5	11.6
40	21.6	22.4	22.9	23.5	23.5	7.1	7.9	8.8	9.4	10.0	10.6	10.8	11.2
60	19.1	20.1	20.7	21.5	22.2	5.8	6.7	7.6	8.4	9.1	9.8	10.3	10.5
80	16.2	17.3	18.4	19.7	20.0	4.3	5.3	6.4	7.2	8.0	8.9	9.3	9.9
100	14.0	14.8	15.6	16.5	17.6	3.3	4.2	5.0	5.9	6.8	7.6	8.4	9.1
120	12.3	12.9	13.7	14.3	15.3	2.4	3.1	3.9	4.8	5.6	6.4	7.3	8.0
140	11.5	12.0	12.6	12.8	13.6	2.1	2.4	2.9	3.8	4.6	5.5	6.3	7.0
160	11.6	11.8	12.1	12.3	12.7	2.0	2.2	2.4	2.7	3.5	4.4	5.1	5.9
180	12.2	12.2	12.3	12.5	12.7	1.9	2.0	2.3	2.6	2.9	3.4	3.9	4.9
200	12.4	12.7	12.8	13.1	13.2	2.3	2.2	2.2	2.4	2.7	3.0	3.4	3.8
220	12.3	12.7	13.0	13.3	13.5	3.0	2.6	2.5	2.4	2.5	2.7	3.1	3.5
240	11.5	12.1	12.4	13.1	13.6	3.7	3.3	3.0	2.9	2.7	2.8	2.9	3.2
260	10.4	11.0	11.5	12.2	12.8	4.8	4.1	3.7	3.5	3.1	3.1	3.0	3.1
280	8.8	9.6	10.4	10.7	11.5	5.5	5.1	4.6	4.1	3.8	3.5	3.5	3.4
300	7.5	7.9	8.6	9.0	10.1	6.2	5.8	5.6	5.0	4.8	4.2	3.9	3.8
320	6.7	6.8	7.3	7.8	8.3	6.9	6.6	6.1	5.9	5.4	5.1	4.7	4.3
340	6.6	6.4	6.6	6.7	6.2	7.2	7.1	6.9	6.5	6.2	5.8	5.5	5.1
360	7.4	6.9	6.5	6.5	6.5	7.5	7.4	7.1	7.0	6.8	6.4	6.2	5.8
380	8.9	8.2	7.5	6.9	6.8	7.5	7.6	7.3	7.3	7.2	7.1	6.7	5.5
400	10.9	10.0	9.0	8.3	7.5	7.3	7.3	7.5	7.4	7.4	7.4	7.1	7.0
420	12.6	11.6	10.7	9.9	9.1	6.9	7.0	7.3	7.4	7.4	7.4	7.3	7.5
440	14.2	13.3	12.5	11.6	10.6	6.5	6.8	6.8	7.1	7.2	7.3	7.3	7.4
460	15.7	14.8	13.9	13.0	12.1	6.2	6.2	6.5	6.7	6.8	7.1	7.1	7.3
480	16.9	16.3	15.5	14.5	13.6	5.8	5.9	6.0	6.2	6.4	6.5	7.0	3.9
500	18.1	17.6	16.6	15.8	15.1	5.3	5.4	5.7	5.8	6.0	6.0	6.3	6.6
520	20.1	19.2	18.1	17.4	16.5	5.1	5.1	5.1	5.3	5.4	5.6	5.8	6.0
540	22.0	21.0	20.2	19.2	18.1	4.7	4.8	4.8	4.8	5.0	5.1	5.4	5.5
560	24.3	23.5	22.6	21.5	20.6	4.4	4.5	4.6	4.6	4.7	4.8	4.8	5.0
580	26.6	25.7	24.9	23.8	23.0	4.2	4.3	4.4	4.3	4.5	4.4	4.4	4.5
600	28.5	27.8	27.0	26.3	25.4	4.0	4.2	4.3	4.2	4.2	4.2	4.2	4.3
620	29.6	29.2	28.8	28.2	27.4	4.2	4.0	4.1	4.0	4.0	4.0	4.0	3.9
640	29.9	30.0	29.9	29.5	29.5	4.3	4.2	4.1	4.0	4.1	4.0	3.9	3.9
660	29.2	29.5	29.7	29.8	29.9	4.6	4.4	4.3	4.1	4.1	4.1	4.0	3.8
680	27.5	28.6	28.9	29.2	29.7	4.8	4.6	4.5	4.3	4.2	4.1	4.0	3.9
700	25.1	26.4	27.3	27.8	28.7	5.3	5.0	4.8	4.5	4.6	4.0	4.1	4.1
720	22.6	23.9	25.0	26.1	26.8	5.8	5.5	5.1	5.0	4.7	4.5	4.1	4.1
740	19.5	21.3	22.5	23.6	24.6	6.5	6.1	5.7	5.4	5.2	4.9	4.6	4.3
760	17.3	18.6	19.4	21.0	22.1	7.4	6.7	6.4	6.0	5.6	5.3	5.1	5.0
780	15.0	15.8	17.1	18.5	19.3	8.2	7.6	6.9	6.5	6.4	5.8	5.6	5.3
800	12.2	14.1	14.8	15.9	17.0	9.2	8.5	8.0	7.3	6.8	6.5	6.1	5.8
820	11.4	12.0	12.5	13.4	15.4	10.1	9.6	8.8	8.2	7.6	7.1	6.7	6.5
840	11.2	11.3	11.7	12.2	13.2	10.9	10.4	9.8	9.1	8.4	7.9	7.5	6.9
860	10.9	10.8	10.9	11.2	11.5	11.7	11.0	10.4	10.0	9.4	8.7	8.2	7.7
880	12.6	11.3	11.1	10.8	11.0	12.3	11.9	11.3	10.6	10.2	9.7	8.9	8.4
900	13.3	12.3	12.9	11.3	11.2	12.4	12.2	11.8	11.6	10.8	10.3	9.7	9.3
920	15.7	14.6	13.7	12.8	12.1	12.3	12.3	12.2	11.9	11.6	11.0	10.5	9.9
940	13.4	17.3	16.2	14.5	14.0	12.1	12.1	12.2	12.2	11.8	11.4	11.0	10.6
960	21.0	20.0	18.9	17.9	16.7	11.4	11.9	11.9	12.0	12.0	11.7	11.1	11.0
980	23.0	22.4	21.4	20.3	19.5	10.6	11.1	11.6	11.8	11.9	11.9	11.7	11.4
1000	21.2	23.7	23.1	22.5	21.6	9.5	10.2	10.8	11.2	11.5	11.7	11.8	11.5
	960	970	980	990	1000	0	10	20	30	40	50	60	70

*Perturbations produced by Mars.*

Arguments II and IV.

IV.

II.	70	80	90	100	110	120	130	140	150	160	170	180	190	200
	"	"	"	"	"	"	"	"	"	"	"	"	"	"
0	11.5	11.2	11.0	10.6	10.1	9.9	9.5	9.0	8.6	8.2	8.1	7.8	7.6	7.4
20	11.6	11.4	11.0	10.9	10.6	10.2	9.7	9.1	9.1	8.8	8.4	8.1	7.9	7.8
40	11.2	11.3	11.2	11.0	10.8	10.5	10.3	9.8	9.4	9.3	9.1	8.7	8.4	8.2
60	10.5	10.9	11.1	10.9	11.0	10.9	10.4	10.0	9.7	9.5	9.2	8.8	8.7	8.4
80	9.9	10.0	10.5	10.9	10.8	10.7	10.4	10.3	10.0	9.7	9.3	9.0	8.8	8.6
100	9.1	9.5	9.8	10.1	10.6	10.5	10.4	10.3	10.1	9.9	9.6	9.3	9.0	8.8
120	8.0	8.8	9.3	9.5	9.9	10.2	10.2	10.1	10.0	9.8	9.6	9.4	9.1	8.9
140	7.0	7.9	8.4	9.0	9.3	9.6	9.9	9.9	9.9	9.7	9.7	9.4	9.3	8.9
160	5.9	6.5	7.2	8.0	8.5	8.9	9.2	9.6	9.5	9.6	9.5	9.5	9.3	9.1
180	4.9	5.6	6.4	6.9	7.7	8.3	8.6	8.9	9.4	9.3	9.3	9.3	9.2	9.1
200	3.8	4.6	5.3	6.0	6.7	7.4	7.9	8.3	8.0	8.9	9.1	9.0	9.0	8.9
220	3.5	3.9	4.4	5.1	5.8	6.4	7.1	7.6	7.9	8.4	8.6	8.8	8.8	8.7
240	3.2	3.6	4.0	4.4	5.0	5.5	6.2	6.8	7.4	7.6	8.1	8.4	8.4	8.5
260	3.1	3.2	3.8	4.1	4.5	4.9	5.4	5.9	6.6	7.1	7.5	7.7	8.0	8.2
280	3.4	3.4	3.5	3.8	4.2	4.5	4.9	5.5	6.2	6.8	7.1	7.5	7.8	8.1
300	3.8	3.7	3.7	3.7	3.9	4.4	4.7	4.9	5.4	5.7	6.0	6.6	6.9	7.3
320	4.3	4.2	4.1	4.0	4.1	4.2	4.4	4.7	5.0	5.4	5.8	6.0	6.4	6.6
340	5.1	4.9	4.6	4.4	4.4	4.3	4.5	4.5	5.0	5.2	5.5	5.8	6.0	6.3
360	5.8	5.6	5.3	5.0	4.8	4.8	4.7	4.8	4.9	5.1	5.4	5.5	5.9	6.1
380	6.5	6.4	5.9	5.7	5.5	5.4	5.1	5.1	5.1	5.1	5.4	5.5	5.7	5.8
400	7.0	6.7	6.7	6.3	6.1	5.9	5.7	5.6	5.5	5.5	5.5	5.6	5.7	5.9
420	7.4	7.2	6.9	7.1	6.7	6.4	6.3	6.1	6.0	5.9	5.9	5.8	5.8	6.1
440	7.5	7.4	7.4	7.0	7.1	7.4	6.8	6.7	6.5	6.3	6.3	6.4	6.2	6.3
460	7.3	7.4	7.4	7.5	7.4	7.3	7.3	7.2	7.1	7.1	6.7	6.7	6.7	6.7
480	6.9	7.1	7.3	7.4	7.5	7.3	7.6	7.5	7.4	7.5	7.4	7.2	7.1	7.1
500	6.6	6.8	6.9	7.2	7.3	7.5	7.5	7.6	7.8	7.7	7.8	7.7	7.6	7.4
520	6.0	6.3	6.5	6.7	7.1	7.2	7.5	7.5	7.7	7.8	7.9	7.6	7.9	7.9
540	5.5	5.7	6.0	6.3	6.6	6.9	7.1	7.3	7.4	7.7	7.9	8.0	8.2	8.3
560	5.0	5.2	5.4	5.8	5.9	6.2	6.6	6.9	7.1	7.4	7.7	7.8	8.1	8.2
580	4.5	4.7	4.9	5.0	5.3	5.7	6.0	6.6	6.8	7.1	7.2	7.5	7.9	8.2
600	4.3	4.3	4.4	4.6	4.6	5.0	5.3	5.6	5.9	6.5	6.9	7.0	7.4	7.7
620	3.9	4.0	4.0	4.1	4.3	4.4	4.6	4.9	5.3	5.4	6.1	6.6	6.9	7.4
640	3.9	3.8	3.8	3.8	3.9	3.9	4.1	4.3	4.5	5.0	5.2	5.8	6.3	6.7
660	3.8	3.7	3.7	3.6	3.6	3.7	3.8	3.9	4.1	4.2	4.5	5.0	5.3	6.0
680	3.9	3.8	3.6	3.4	3.5	3.4	3.5	3.5	3.6	3.7	3.8	4.2	4.6	4.9
700	4.1	3.9	3.8	3.6	3.5	3.3	3.3	3.2	3.2	3.2	3.5	3.6	3.8	4.2
720	4.1	4.1	4.0	3.8	3.6	3.5	3.3	3.2	3.3	3.2	3.0	3.2	3.4	3.6
740	4.3	4.3	4.2	4.0	3.8	3.7	3.5	3.2	3.0	3.0	2.9	2.8	2.9	3.1
760	5.0	4.7	4.4	4.3	4.1	3.8	3.7	3.4	3.1	3.0	2.9	2.7	2.7	2.8
780	5.3	5.1	4.7	4.6	4.4	4.4	4.0	3.8	3.4	3.2	2.9	2.8	2.7	2.5
800	5.8	5.5	5.4	4.8	4.7	4.7	4.5	4.2	3.9	3.5	3.3	2.9	2.8	2.7
820	6.5	6.1	5.8	5.6	5.0	5.0	4.9	4.6	4.3	4.1	3.6	3.3	3.0	2.9
840	6.9	6.7	6.3	6.1	5.8	5.3	5.2	4.9	4.9	4.5	4.2	3.9	3.5	3.1
860	7.7	7.4	6.9	6.6	6.2	6.2	5.5	5.4	5.2	5.0	4.8	4.4	4.1	3.6
880	8.4	7.9	7.6	7.1	6.9	6.4	6.4	5.8	5.7	5.4	5.2	5.0	4.6	4.3
900	9.3	8.7	8.3	7.7	7.4	7.1	6.7	6.6	6.1	6.0	5.6	5.4	5.2	4.9
920	9.9	9.3	8.8	8.4	7.9	7.7	7.3	6.9	6.6	6.3	6.2	6.1	5.6	5.4
940	10.6	10.1	9.5	8.9	8.7	8.2	7.8	7.6	7.2	7.1	6.5	6.5	6.3	5.9
960	11.0	10.7	10.3	9.7	9.1	8.7	8.4	8.0	7.8	7.4	7.2	6.9	6.7	6.5
980	11.4	11.0	10.6	10.2	9.8	9.2	8.9	8.4	8.1	8.0	7.6	7.3	7.2	6.9
1000	11.5	11.2	11.0	10.6	10.0	9.9	9.5	9.0	8.6	8.2	8.1	7.4	7.6	7.4
	70	80	90	100	110	120	130	14	150	160	170	180	190	200

*Perturbations produced by Mars.*

Arguments II. and IV.

## IV.

II.	200	210	220	230	240	250	260	270	280	290	300	310	320
0	"	"	"	"	"	"	"	"	"	"	"	"	"
20	7.4	7.2	7.0	6.6	6.4	6.2	5.7	5.3	4.9	4.7	4.1	3.8	3.4
40	7.8	7.2	7.3	7.2	7.0	6.6	6.3	6.0	5.7	5.3	5.0	4.4	3.9
60	8.2	8.1	7.6	7.5	7.3	7.2	6.8	6.6	6.2	5.9	5.6	5.2	4.7
80	8.4	8.0	7.9	7.8	7.6	7.5	7.3	7.1	6.8	6.4	6.1	5.8	5.4
100	8.6	8.5	8.2	8.0	7.6	7.7	7.6	7.4	7.1	7.0	6.7	6.3	6.0
120	8.8	8.5	8.6	8.4	8.2	7.6	7.7	7.8	7.6	7.3	7.2	6.9	6.6
140	8.9	8.7	8.4	8.4	8.3	8.3	8.0	7.9	7.7	7.6	7.5	7.3	7.0
160	8.9	8.7	8.4	8.3	8.2	8.1	8.3	8.0	7.9	7.8	7.7	7.5	7.4
180	9.1	8.9	8.7	8.4	8.3	8.3	8.2	8.1	8.0	7.9	7.9	7.7	7.6
200	9.1	8.8	8.7	8.5	8.4	8.2	8.0	8.0	8.1	7.9	7.8	8.0	7.8
220	8.9	8.8	8.6	8.4	8.4	8.3	8.1	8.0	7.9	7.8	7.8	7.9	7.9
240	8.7	8.7	8.6	8.4	8.2	8.1	8.0	7.9	7.8	7.7	7.7	7.6	7.7
260	8.5	8.4	8.5	8.3	8.1	8.0	7.8	7.8	7.8	7.8	7.8	7.8	7.6
280	8.2	8.2	8.1	8.1	8.1	7.8	7.8	7.7	7.6	7.6	7.6	7.5	7.4
300	7.8	7.8	8.0	7.8	7.9	7.9	7.7	7.5	7.5	7.3	7.3	7.4	7.3
320	7.3	7.6	7.5	7.6	7.7	7.6	7.6	7.6	7.4	7.3	7.1	7.0	7.1
340	6.6	7.1	7.3	7.4	7.4	7.3	7.4	7.4	7.3	7.1	7.0	7.0	6.8
360	6.3	6.4	6.7	7.2	7.1	7.2	7.2	7.1	7.1	7.0	6.9	6.8	6.8
380	6.1	6.2	6.4	6.5	6.9	6.9	7.0	7.0	6.9	6.8	6.7	6.6	6.5
400	5.8	6.1	6.3	6.4	6.6	6.7	6.6	6.6	6.7	6.8	6.7	6.6	6.5
420	5.9	6.0	6.2	6.3	6.4	6.5	6.6	6.6	6.5	6.6	6.6	6.5	6.4
440	6.1	6.3	6.2	6.4	6.3	6.4	6.5	6.6	6.5	6.5	6.5	6.5	6.4
460	6.3	6.4	6.4	6.6	6.5	6.6	6.5	6.5	6.5	6.5	6.3	6.3	6.2
480	6.7	6.5	6.5	6.6	6.7	6.9	6.7	6.6	6.6	6.6	6.5	6.3	6.2
500	7.1	7.1	7.0	6.9	6.9	6.9	7.0	7.0	6.8	6.7	6.6	6.5	6.3
520	7.4	7.5	7.4	7.4	7.3	7.2	7.3	7.2	7.1	6.9	6.8	6.8	6.6
540	7.9	7.8	7.8	7.8	7.8	7.6	7.6	7.5	7.5	7.4	7.1	7.0	6.9
560	8.3	8.3	8.3	8.2	8.2	8.1	8.0	7.9	7.9	7.8	7.6	7.5	7.2
580	8.2	8.6	8.4	8.6	8.7	8.5	8.5	8.4	8.2	8.3	8.2	8.0	7.6
600	8.2	8.3	8.6	8.8	8.8	9.0	8.9	8.9	8.7	8.7	8.6	8.4	8.4
620	7.7	8.1	8.5	8.6	8.9	9.1	9.1	9.2	9.2	9.1	9.0	8.8	8.7
640	7.4	7.6	8.0	8.5	8.7	9.0	9.2	9.5	9.5	9.5	9.4	9.3	9.2
660	6.7	7.2	7.5	7.9	8.3	8.7	9.0	9.3	9.5	9.8	9.8	9.7	9.7
680	6.0	6.3	7.0	7.3	7.7	8.2	8.7	9.0	9.4	9.7	9.8	10.1	10.0
700	4.9	5.6	6.0	6.6	7.1	7.7	8.1	8.5	9.0	9.3	9.8	10.0	10.2
720	4.2	4.5	5.2	5.8	6.4	6.8	7.4	8.0	8.5	8.9	9.2	9.8	10.1
740	3.6	3.9	4.3	4.7	5.3	5.9	6.6	7.0	7.8	8.3	8.8	9.1	9.7
760	3.1	3.3	3.6	3.9	4.4	4.8	5.6	6.2	6.9	7.5	8.0	8.7	9.2
780	2.8	2.8	3.0	3.3	3.6	4.0	4.4	5.1	5.8	6.5	7.2	7.8	8.4
800	2.5	2.6	2.5	2.7	3.1	3.3	3.7	4.1	4.8	5.4	6.1	6.9	7.6
820	2.7	2.5	2.5	2.5	2.5	2.7	3.0	3.4	3.8	4.4	5.0	5.6	6.6
840	2.9	2.6	2.4	2.3	2.2	2.3	2.6	2.8	3.1	3.4	4.1	4.7	5.4
860	3.1	2.8	2.6	2.4	2.3	2.2	2.3	2.4	2.6	2.8	3.2	3.8	4.3
880	3.6	3.3	3.0	2.7	2.4	2.3	2.1	2.2	2.3	2.5	2.7	3.0	3.4
900	4.3	3.8	3.6	3.2	2.8	2.5	2.3	2.1	2.0	2.2	2.3	2.5	2.6
920	4.9	4.6	4.2	3.6	3.4	2.9	2.6	2.3	2.2	2.2	2.1	2.2	2.4
940	5.4	5.1	4.6	4.5	3.9	3.5	3.2	2.9	2.6	2.2	2.0	2.1	2.2
960	5.9	5.7	5.3	4.9	4.7	4.3	3.8	3.4	3.0	2.7	2.4	2.1	2.0
980	6.5	6.2	5.9	5.5	5.1	4.9	4.5	4.0	3.4	3.1	2.8	2.4	2.3
1000	6.9	6.8	6.4	6.1	5.8	5.4	5.1	4.8	4.3	3.9	3.5	3.0	2.7
1000	7.4	7.2	7.0	6.6	6.4	6.2	5.7	5.3	4.9	4.7	4.1	3.8	3.4
	200	210	220	230	240	250	260	270	280	290	300	310	320



*Perturbations produced by Mars.*

Arguments II. and IV.

IV.

II.	320	330	340	350	360	370	380	390	400	410	420	430	440
	"	"	"	"	"	"	"	"	"	"	"	"	"
0	3.4	2.8	2.6	2.4	2.2	2.3	2.3	2.5	2.7	2.9	3.4	4.0	4.5
20	3.9	3.5	3.1	2.7	2.6	2.4	2.4	2.3	2.5	2.7	3.0	3.3	3.8
40	4.7	4.2	3.9	3.5	3.0	2.8	2.7	2.6	2.5	2.6	2.8	2.9	3.2
60	5.4	5.0	4.6	4.2	3.8	3.4	3.1	2.8	2.8	2.7	2.7	2.7	3.0
80	6.0	5.7	5.4	4.8	4.4	4.0	3.6	3.4	3.1	2.9	2.9	2.9	2.9
100	6.6	6.3	5.9	5.6	5.2	4.8	4.3	4.0	3.7	3.5	3.2	3.0	3.0
120	7.0	6.9	6.4	6.1	5.8	5.3	5.2	4.6	4.3	4.0	3.8	3.6	3.4
140	7.4	7.2	6.9	6.6	6.5	6.1	5.6	5.4	5.0	4.6	4.3	4.0	3.9
160	7.6	7.5	7.3	7.0	6.8	6.6	6.2	5.9	5.5	5.3	4.9	4.6	4.4
180	7.8	7.7	7.5	7.4	7.3	6.9	6.7	6.5	6.2	5.8	5.6	5.3	5.0
200	7.9	7.8	7.7	7.6	7.5	7.3	7.1	6.9	6.6	6.4	6.1	5.6	5.5
220	7.7	7.7	7.7	7.8	7.7	7.5	7.3	7.2	7.0	6.7	6.5	6.2	5.9
240	7.6	7.6	7.6	7.6	7.7	7.6	7.5	7.3	7.2	7.1	6.9	6.6	6.4
260	7.4	7.3	7.5	7.5	7.5	7.6	7.6	7.5	7.5	7.3	7.1	7.0	6.7
280	7.3	7.4	7.3	7.3	7.4	7.4	7.3	7.4	7.3	7.5	7.2	7.1	6.9
300	7.1	7.1	7.1	7.0	7.2	7.3	7.3	7.3	7.2	7.2	7.3	7.2	7.1
320	6.8	6.8	6.9	6.9	6.8	7.0	7.1	7.1	7.1	7.1	7.1	7.0	7.2
340	6.8	6.7	6.6	6.6	6.6	6.8	6.9	6.9	7.0	7.0	6.9	6.9	6.9
360	6.5	6.5	6.4	6.3	6.4	6.5	6.6	6.7	6.8	6.8	6.8	6.8	6.9
380	6.5	6.3	6.3	6.2	6.2	6.2	6.3	6.3	6.4	6.5	6.6	6.7	6.7
400	6.4	6.2	6.2	6.0	6.1	6.0	6.0	6.0	6.0	6.1	6.2	6.3	6.4
420	6.4	6.2	6.1	6.0	5.9	5.8	5.9	5.9	5.9	5.9	5.9	6.0	6.0
440	6.2	6.1	6.0	5.8	5.8	5.7	5.6	5.6	5.6	5.7	5.7	5.8	5.9
460	6.2	6.0	5.9	5.8	5.7	5.5	5.5	5.4	5.5	5.4	5.5	5.3	5.4
480	6.3	6.2	6.0	5.7	5.6	5.5	5.4	5.3	5.2	5.2	5.2	5.3	5.3
500	6.6	6.4	6.2	6.0	5.7	5.4	5.3	5.2	5.1	5.1	5.1	5.0	5.0
520	6.9	6.7	6.4	6.1	6.1	5.7	5.5	5.1	5.1	5.0	4.9	5.0	4.9
540	7.2	7.1	6.7	6.5	6.2	6.1	5.8	5.5	5.2	5.0	4.9	4.8	4.8
560	7.6	7.4	7.3	7.0	6.6	6.3	6.0	5.8	5.4	5.3	5.0	4.7	4.7
580	8.4	8.0	7.8	7.5	7.0	6.8	6.3	6.2	5.9	5.5	5.3	5.0	4.9
600	8.7	8.6	8.3	8.0	7.8	7.4	7.0	6.6	6.3	6.0	5.6	5.3	5.1
620	9.2	9.1	8.9	8.6	8.4	8.1	7.6	7.2	6.8	6.5	6.1	5.7	5.3
640	9.7	9.6	9.4	9.3	9.0	8.7	8.2	7.8	7.4	7.0	6.6	6.3	5.8
660	10.0	10.0	9.9	9.8	9.6	9.3	8.9	8.5	8.2	7.7	7.2	6.8	6.4
680	10.2	10.4	10.3	10.2	10.1	9.9	9.6	9.3	9.0	8.5	8.1	7.5	7.1
700	10.1	10.3	10.5	10.6	10.4	10.3	10.1	9.8	9.6	9.3	8.9	8.3	7.8
720	9.7	10.1	10.3	10.6	10.7	10.6	10.5	10.5	10.2	10.0	9.6	9.2	8.6
740	9.2	9.6	10.0	10.3	10.6	10.7	10.8	10.9	10.6	10.5	10.2	9.9	9.4
760	8.4	9.0	9.5	9.8	10.2	10.6	10.9	11.0	11.0	11.0	10.7	10.5	10.3
780	7.6	8.2	8.9	9.4	9.9	10.3	10.6	10.9	11.1	11.2	11.0	10.8	10.7
800	6.6	7.3	7.9	8.5	9.2	9.8	10.1	10.6	10.8	11.1	11.3	11.1	11.0
820	5.4	6.0	7.0	7.6	8.2	8.9	9.6	10.0	10.5	10.8	11.0	11.3	11.3
840	4.3	5.0	5.6	6.5	7.2	7.9	9.8	9.2	9.9	10.3	10.7	10.9	11.2
860	3.4	4.0	4.6	5.3	6.1	6.9	7.5	8.4	9.1	9.6	10.1	10.7	10.9
880	2.6	3.1	3.7	4.3	5.0	5.7	6.6	7.1	8.1	8.7	9.4	9.8	10.4
900	2.4	2.7	3.0	3.4	4.0	4.6	5.4	6.1	6.9	7.6	8.4	9.1	9.7
920	2.2	2.3	2.3	2.8	3.3	3.7	4.3	5.0	5.8	6.5	7.2	8.0	8.7
940	2.0	2.1	2.3	2.3	2.7	2.9	3.4	4.1	4.7	5.5	6.1	7.0	7.7
960	2.3	2.2	2.2	2.3	2.3	2.5	2.8	3.2	3.9	4.5	5.1	5.7	6.5
980	2.7	2.4	2.2	2.3	2.3	2.4	2.5	2.8	3.0	3.6	4.1	4.7	5.5
1000	3.4	2.8	2.6	2.4	2.2	2.3	2.3	2.5	2.7	2.9	3.4	4.0	4.5
	320	330	340	350	360	370	380	390	400	410	420	430	440



*Perturbations produced by Mars.*

## Arguments II and IV.

## IV.

II.	440	450	460	470	480	490	500	510	520	530	540	550	560
0	4.5	5.2	5.9	6.6	7.3	8.0	8.5	9.0	9.5	10.0	10.4	10.7	10.9
20	3.8	4.3	4.9	5.6	6.2	6.9	7.6	8.2	8.8	9.3	9.7	10.0	11.4
40	3.2	3.7	4.2	4.8	5.4	5.9	6.6	7.3	7.9	8.4	8.9	9.4	9.8
60	3.0	3.2	3.6	4.0	4.5	5.1	5.7	6.3	6.9	7.5	8.0	8.6	9.1
80	2.9	3.1	3.3	3.5	3.9	4.4	4.9	5.4	5.9	6.5	7.1	7.7	8.2
100	3.0	3.1	3.2	3.5	3.6	3.8	4.2	4.8	5.3	5.9	6.4	6.9	7.4
120	3.4	3.3	3.3	3.4	3.5	3.6	3.9	4.2	4.7	5.1	5.6	6.0	6.6
140	3.9	3.8	3.6	3.6	3.6	3.7	4.0	4.0	4.2	4.6	5.0	5.4	5.9
160	4.4	4.2	3.9	4.1	3.8	3.7	4.0	4.1	4.2	4.5	4.6	4.9	5.3
180	5.0	4.8	4.4	4.2	4.2	4.2	4.0	4.1	4.3	4.4	4.4	4.7	5.0
200	5.5	5.2	5.1	4.8	4.6	4.5	4.5	4.4	4.5	4.5	4.7	4.6	4.8
220	5.9	5.7	5.5	5.3	5.1	4.9	4.9	4.8	4.7	4.8	4.8	4.9	5.0
240	6.4	6.2	5.9	5.8	5.6	5.4	5.3	5.2	5.1	5.1	5.1	5.2	5.2
260	6.7	6.6	6.4	6.1	6.0	5.9	5.8	5.7	5.6	5.5	5.4	5.4	5.4
280	6.9	6.8	6.7	6.5	6.3	6.2	6.1	6.0	5.9	5.9	5.9	5.8	5.8
300	7.1	7.0	6.8	6.8	6.6	6.5	6.4	6.3	6.2	6.2	6.2	6.2	6.2
320	7.2	7.1	6.9	6.8	6.8	6.7	6.6	6.5	6.5	6.5	6.5	6.6	6.6
340	6.9	6.9	7.0	6.9	6.9	6.8	6.7	6.8	6.7	6.6	6.7	6.8	6.9
360	6.9	6.8	6.8	6.8	6.8	6.7	6.7	6.6	6.6	6.8	6.8	6.8	6.9
380	6.7	6.5	6.5	6.6	6.7	6.6	6.6	6.7	6.7	6.7	6.8	6.9	6.9
400	6.4	6.4	6.3	6.3	6.4	6.5	6.5	6.5	6.6	6.7	6.7	6.8	6.8
420	6.0	6.2	6.3	6.3	6.2	6.2	6.3	6.3	6.3	6.3	6.5	6.6	6.7
440	5.9	5.9	6.0	6.0	6.0	6.0	6.0	6.1	6.0	6.1	6.2	6.2	6.4
460	5.4	5.5	5.7	5.8	5.8	5.8	5.8	5.8	5.8	5.8	5.9	6.0	6.1
480	5.3	5.3	5.5	5.5	5.5	5.6	5.5	5.6	5.4	5.6	5.7	5.5	5.8
500	5.0	5.0	5.1	5.2	5.3	5.3	5.3	5.2	5.2	5.2	5.3	5.4	5.4
520	4.9	4.9	4.9	4.8	5.0	5.1	5.1	5.1	5.1	5.1	5.0	5.0	5.1
540	4.8	4.8	4.7	4.8	4.8	4.9	4.9	5.0	4.9	4.8	4.8	4.9	4.8
560	4.7	4.6	4.6	4.7	4.7	4.6	4.7	4.7	4.7	4.7	4.6	4.6	4.6
580	4.9	4.6	4.5	4.5	4.6	4.5	4.4	4.4	4.5	4.5	4.5	4.4	4.4
600	5.1	4.9	4.6	4.5	4.4	4.4	4.4	4.3	4.3	4.3	4.3	4.3	4.3
620	5.3	5.1	4.9	4.7	4.6	4.4	4.3	4.1	4.2	4.2	4.2	4.2	4.1
640	5.8	5.4	5.2	5.0	4.7	4.6	4.4	4.1	4.1	4.1	4.2	4.2	4.0
660	6.4	6.0	5.7	5.4	5.0	4.8	4.7	4.5	4.3	4.2	4.2	4.1	4.0
680	7.1	6.6	6.2	5.7	5.4	5.1	4.9	4.7	4.5	4.4	4.3	4.0	3.9
700	7.8	7.2	6.8	6.4	6.0	5.6	5.3	5.0	4.7	4.6	4.6	4.3	4.1
720	8.6	8.0	7.6	7.1	6.6	6.2	5.7	5.5	5.2	4.9	4.6	4.6	4.3
740	9.4	9.0	8.4	8.0	7.4	6.9	6.3	6.0	5.6	5.3	5.0	4.7	4.5
760	10.3	9.7	9.3	8.6	8.1	7.6	7.2	6.5	6.2	5.8	5.5	5.2	4.9
780	10.7	10.5	9.9	9.6	9.0	8.5	7.8	7.4	7.0	6.4	6.1	5.7	5.5
800	11.0	11.0	10.6	10.2	9.9	9.3	8.8	8.1	7.7	7.3	6.7	6.3	5.8
820	11.3	11.1	10.9	10.6	10.3	10.0	9.6	9.1	8.5	7.9	7.4	7.0	6.6
840	11.2	11.3	11.2	11.1	11.0	10.7	10.2	9.9	9.4	8.8	8.2	7.7	7.3
860	10.9	11.1	11.4	11.3	11.3	11.2	10.7	10.4	9.9	9.6	9.2	8.5	7.9
880	10.4	10.8	11.0	11.3	11.2	11.2	11.2	10.9	10.5	10.3	9.8	9.3	8.7
900	9.7	10.1	10.6	11.0	11.2	11.2	11.2	11.0	10.9	10.7	10.2	10.0	9.4
920	8.7	9.3	9.9	10.3	10.8	11.0	11.1	11.2	11.2	11.0	10.7	10.4	10.1
940	7.7	8.2	8.8	9.5	10.1	10.4	10.9	11.0	11.2	11.2	11.0	10.7	10.5
960	6.5	7.3	8.1	8.6	9.3	9.8	10.2	10.6	10.8	11.1	11.2	10.9	10.8
980	5.5	6.2	7.0	7.7	8.3	8.9	9.5	10.0	10.4	10.6	10.8	11.0	10.9
1000	4.5	5.2	5.9	6.6	7.3	8.0	8.5	9.0	9.5	10.0	10.4	10.7	10.9
	440	450	460	470	480	490	500	510	520	530	540	550	560

*Perturbations produced by Mars.*

Arguments II and IV.

IV.

II.	560	570	580	590	600	610	620	630	640	650	660	670	680
	"	"	"	"	"	"	"	"	"	"	"	"	"
0	10.9	10.8	10.6	10.4	10.3	10.0	9.7	9.2	8.9	8.5	8.1	7.9	7.7
20	11.4	10.6	10.7	10.6	10.4	10.2	9.9	9.7	9.3	9.0	8.8	8.5	8.1
40	9.8	10.1	10.4	10.4	10.5	10.3	10.2	9.9	9.6	9.4	9.1	8.9	8.5
60	9.1	9.4	9.8	10.2	10.2	10.3	10.2	10.1	9.9	9.6	9.3	9.0	8.8
80	8.2	8.7	9.0	9.3	9.6	9.8	10.0	9.9	9.8	9.7	9.5	9.3	9.1
100	7.4	7.9	8.4	8.7	9.0	9.4	9.6	9.7	9.8	9.7	9.7	9.5	9.2
120	6.6	6.9	7.6	8.1	8.3	8.6	9.0	9.2	9.4	9.5	9.5	9.4	9.3
140	5.9	6.3	6.8	7.2	7.7	8.0	8.3	8.7	8.9	9.1	9.2	9.3	9.3
160	5.3	5.8	6.0	6.5	6.9	7.4	7.7	8.0	8.4	8.5	8.8	8.9	9.0
180	5.0	5.2	5.6	6.0	6.3	6.7	7.1	7.2	7.7	8.1	8.1	8.4	8.6
200	4.8	5.0	5.3	5.4	5.8	6.1	6.5	6.7	7.1	7.3	7.7	7.8	8.0
220	5.0	5.0	5.1	5.3	5.5	5.7	6.0	6.3	6.6	6.8	7.0	7.3	7.5
240	5.2	5.2	5.3	5.3	5.4	5.5	5.7	5.9	6.1	6.4	6.6	6.8	7.1
260	5.4	5.5	5.5	5.5	5.5	5.5	5.5	5.7	5.8	6.0	6.3	6.4	6.5
280	5.8	5.8	5.8	5.9	5.8	5.8	5.8	5.9	5.9	5.9	6.0	6.1	6.2
300	6.2	6.1	6.2	6.1	6.1	6.1	6.2	6.1	6.0	5.9	5.9	6.0	6.1
320	6.6	6.5	6.6	6.6	6.5	6.5	6.6	6.5	6.5	6.3	6.1	6.0	6.0
340	6.9	6.9	6.9	7.0	7.0	6.9	6.8	6.9	6.9	6.8	6.6	6.5	6.3
360	6.9	7.0	7.2	7.3	7.3	7.3	7.4	7.3	7.3	7.1	7.1	7.0	6.7
380	6.9	7.0	7.2	7.4	7.5	7.6	7.7	7.7	7.7	7.6	7.5	7.4	7.2
400	6.8	7.0	7.1	7.3	7.6	7.9	8.0	8.0	8.1	8.1	8.1	7.9	7.8
420	6.7	6.9	7.0	7.2	7.6	7.8	8.0	8.2	8.3	8.4	8.4	8.5	8.4
440	6.4	6.6	6.9	7.0	7.3	7.5	7.9	8.2	8.4	8.6	8.8	8.8	8.9
460	6.1	6.2	6.5	6.9	7.1	7.2	7.6	8.0	8.4	8.7	9.0	9.1	9.2
480	5.8	5.9	6.0	6.2	6.7	7.1	7.2	7.6	7.9	8.5	8.9	9.2	9.3
500	5.4	5.5	5.6	5.9	6.1	6.4	6.9	7.2	7.7	7.9	8.4	9.0	9.4
520	5.1	5.2	5.2	5.3	5.6	5.9	6.3	6.7	7.0	7.6	8.0	8.4	9.0
540	4.8	4.8	4.8	5.0	5.1	5.4	5.6	6.0	6.4	6.7	7.5	8.1	8.5
560	4.6	4.5	4.5	4.5	4.7	4.8	5.0	5.3	5.8	6.2	6.6	7.1	7.8
580	4.4	4.3	4.3	4.3	4.3	4.3	4.5	4.7	5.2	5.5	5.9	6.4	6.9
600	4.3	4.3	4.2	4.1	4.0	4.0	4.1	4.2	4.5	4.8	5.1	5.7	6.2
620	4.1	4.0	4.0	3.9	3.9	3.8	3.8	3.8	3.8	4.0	4.4	4.9	5.4
640	4.0	3.9	4.0	3.8	3.8	3.8	3.7	3.5	3.5	3.6	3.8	4.0	4.5
660	4.0	4.0	3.9	3.8	3.7	3.5	3.5	3.4	3.3	3.3	3.4	3.5	3.7
680	3.9	4.0	3.9	3.8	3.6	3.5	3.4	3.3	3.2	3.1	3.1	3.1	3.1
700	4.1	3.9	3.9	3.9	3.7	3.5	3.4	3.3	3.2	3.0	3.0	3.0	2.9
720	4.3	4.1	4.0	3.9	3.8	3.8	3.5	3.4	3.1	2.9	2.9	2.7	2.7
740	4.5	4.2	4.2	4.2	4.0	3.7	3.6	3.4	3.3	3.0	2.8	2.6	2.5
760	4.9	4.7	4.5	4.3	4.2	4.1	3.8	3.6	3.3	3.1	2.9	2.8	2.5
780	5.5	5.1	4.9	4.5	4.4	4.3	4.1	3.9	3.8	3.4	3.2	3.0	2.7
800	5.8	5.6	5.2	5.0	4.6	4.5	4.4	4.3	4.1	3.8	3.5	3.1	2.8
820	6.6	6.1	5.8	5.5	5.3	5.0	4.8	4.6	4.4	4.2	4.0	3.6	3.3
840	7.3	6.8	6.5	6.1	5.7	5.5	5.2	5.0	4.7	4.6	4.3	4.1	3.8
860	7.9	7.5	7.0	6.7	6.4	5.9	5.8	5.4	5.1	5.0	4.8	4.6	4.4
880	8.7	8.2	7.8	7.3	6.9	6.6	6.3	6.0	5.7	5.4	5.2	5.0	4.7
900	9.4	9.0	8.5	8.0	7.6	7.2	6.8	6.6	6.3	5.9	5.6	5.4	5.2
920	10.1	9.8	9.2	8.7	8.3	7.8	7.4	7.0	6.7	6.4	6.0	5.8	5.7
940	10.5	10.2	9.8	9.4	8.8	8.5	8.0	7.6	7.3	6.9	6.6	6.2	6.1
960	10.8	10.5	10.2	10.0	9.5	9.1	8.6	8.2	7.8	7.5	7.1	6.8	6.6
980	10.9	10.7	10.3	10.2	9.9	9.6	9.2	9.0	8.5	8.0	7.7	7.4	7.2
1000	10.9	10.8	10.6	10.4	10.3	10.0	9.7	9.2	8.9	8.5	8.1	7.9	7.7
	560	570	580	590	600	610	620	630	640	650	660	670	680

*Perturbations produced by Mars.*

Arguments II. and IV.

IV.

II.	680	690	700	710	720	730	740	750	760	770	780	790	800
	"	"	"	"	"	"	"	"	"	"	"	"	"
0	7.7	7.4	6.9	6.8	6.7	6.4	6.1	5.8	5.5	5.2	4.8	4.4	3.7
20	8.1	7.8	7.4	7.0	7.1	6.9	6.7	6.4	6.1	5.8	5.5	5.1	4.7
40	8.5	8.3	7.8	7.5	7.2	7.1	7.0	6.9	6.6	6.4	6.1	5.8	5.3
60	8.8	8.6	8.3	8.1	7.8	7.6	7.5	7.4	7.1	6.9	6.7	6.3	6.0
80	9.1	8.9	8.7	8.4	8.1	8.0	7.8	7.6	7.4	7.3	7.1	6.9	6.5
100	9.2	8.9	8.8	8.7	8.6	8.3	8.0	7.7	7.6	7.6	7.6	7.3	7.0
120	9.3	9.2	9.0	8.7	8.6	8.4	8.2	8.1	7.9	7.8	7.7	7.6	7.5
140	9.3	9.2	9.0	9.0	8.7	8.5	8.4	8.3	8.0	7.8	7.7	7.7	7.7
160	9.0	9.0	8.9	8.8	8.7	8.6	8.5	8.4	8.2	8.0	7.9	7.8	7.8
180	8.6	8.6	8.7	8.7	8.7	8.6	8.5	8.3	8.3	8.0	8.2	7.8	7.9
200	8.0	8.2	8.3	8.3	8.5	8.4	8.4	8.4	8.2	8.1	8.1	8.1	7.9
220	7.5	7.7	7.9	8.1	8.2	8.2	8.1	8.2	8.2	8.0	8.1	8.0	8.0
240	7.1	7.2	7.4	7.5	7.6	7.7	7.8	7.8	7.9	8.0	8.0	7.8	7.8
260	6.5	6.7	6.9	7.1	7.2	7.3	7.4	7.5	7.6	7.6	7.7	7.7	7.8
280	6.2	6.3	6.5	6.7	6.7	6.9	7.1	7.2	7.3	7.3	7.3	7.3	7.4
300	6.1	6.0	6.2	6.4	6.4	6.5	6.6	6.7	6.9	6.9	6.9	7.1	7.1
320	6.0	6.0	6.0	6.0	6.2	6.1	6.2	6.3	6.5	6.5	6.6	6.6	6.8
340	6.3	6.2	6.0	6.0	6.0	6.0	6.1	6.1	6.2	6.2	6.3	6.3	6.4
360	6.7	6.6	6.4	6.1	6.0	5.9	6.0	5.9	5.9	5.9	6.0	6.1	6.2
380	7.2	7.1	6.8	6.6	6.4	6.2	6.1	5.9	5.8	5.7	5.6	5.8	5.9
400	7.8	7.7	7.4	7.1	6.8	6.6	6.4	6.1	6.0	5.8	5.6	5.5	5.6
420	8.4	8.2	8.0	7.8	7.5	7.2	6.8	6.5	6.2	6.0	5.7	5.5	5.4
440	8.9	8.8	8.7	8.4	8.2	7.8	7.5	7.1	6.6	6.2	6.0	5.7	5.6
460	9.2	9.2	9.2	9.0	8.8	8.5	8.2	7.9	7.5	6.9	6.5	6.3	6.0
480	9.3	9.5	9.6	9.6	9.4	9.2	9.1	8.6	8.3	7.8	7.2	6.9	6.5
500	9.4	9.6	9.8	10.0	9.9	9.8	9.6	9.4	9.1	8.7	8.2	7.6	7.2
520	9.0	9.5	9.8	10.1	10.2	10.3	10.3	10.0	9.8	9.5	9.1	8.5	8.0
540	8.5	9.1	9.5	10.0	10.3	10.5	10.6	10.6	10.4	10.1	9.8	9.5	9.0
560	7.8	8.5	9.0	9.5	9.9	10.4	10.8	10.8	10.9	10.8	10.6	10.2	9.9
580	6.9	7.6	8.3	9.0	9.7	10.0	10.4	10.7	11.1	11.2	11.0	11.0	10.6
600	6.2	6.8	7.4	8.0	8.9	9.6	10.1	10.4	10.9	11.3	11.4	11.3	11.2
620	5.4	5.9	6.5	7.1	7.8	8.6	9.4	10.3	10.6	11.0	11.5	11.7	11.7
640	4.5	5.0	5.5	6.2	6.8	7.6	8.4	9.2	10.0	10.7	11.1	11.6	11.8
660	3.7	4.1	4.7	5.2	5.9	6.5	7.3	8.3	9.1	9.8	10.5	11.2	11.5
680	3.1	3.4	3.8	4.3	4.8	5.5	6.2	7.0	7.8	8.7	9.6	10.2	11.0
700	2.9	2.8	3.0	3.4	3.9	4.5	5.2	6.0	6.7	7.5	8.5	9.4	10.1
720	2.7	2.6	2.5	2.7	3.1	3.5	4.0	4.8	5.6	6.4	7.3	8.2	9.1
740	2.5	2.4	2.4	2.4	2.5	2.7	3.1	3.6	4.5	5.2	6.1	6.9	7.8
760	2.5	2.3	2.2	2.1	2.1	2.3	2.4	2.8	3.2	4.1	4.7	5.7	6.6
780	2.7	2.5	2.3	2.1	2.0	1.9	2.1	2.2	2.5	2.9	3.6	4.4	5.2
800	2.8	2.7	2.4	2.2	2.0	1.8	1.8	1.8	2.0	2.3	2.5	3.2	4.0
820	3.3	3.0	2.7	2.3	2.1	1.9	1.8	1.5	1.7	1.7	2.0	2.2	2.9
840	3.8	3.5	3.0	2.6	2.3	2.1	1.9	1.6	1.5	1.5	1.6	1.7	2.2
860	4.4	4.0	3.5	3.2	2.8	2.3	1.9	1.7	1.4	1.3	1.2	1.4	1.6
880	4.7	4.4	4.1	3.7	3.3	3.0	2.5	2.1	1.7	1.4	1.3	1.2	1.2
900	5.2	5.0	4.6	4.3	4.0	3.6	3.2	2.7	2.2	1.6	1.3	1.2	1.1
920	5.7	5.3	5.1	5.0	4.6	4.2	3.8	3.4	2.9	2.3	1.9	1.3	1.1
940	6.1	5.9	5.6	5.4	5.2	4.8	4.5	3.9	3.5	3.1	2.6	2.1	1.5
960	6.6	6.4	6.2	5.9	5.6	5.4	5.1	4.7	4.3	3.7	3.2	2.8	2.3
980	7.2	6.9	6.6	6.4	6.2	5.9	5.6	5.3	5.0	4.6	4.0	3.5	3.0
1000	7.7	7.4	6.9	6.8	6.7	6.4	6.1	5.8	5.5	5.2	4.8	4.4	3.7
	680	690	700	710	720	730	740	750	760	770	780	790	800

*Perturbations produced by Mars.*

Arguments II. and IV.

· IV.

II.	800	810	820	830	840	850	860	870	880	890	900	910	920
	"	"	"	"	"	"	"	"	"	"	"	"	"
0	3.7	3.2	2.6	2.1	1.7	1.3	0.9	0.7	0.7	1.0	1.2	1.6	2.2
20	4.7	4.2	3.6	3.1	2.4	1.9	1.5	1.2	0.8	0.6	0.9	1.2	1.5
40	5.3	4.9	4.5	3.8	3.3	2.7	2.0	1.7	1.4	1.0	0.8	0.9	1.0
60	6.0	5.7	5.2	4.7	4.1	3.6	3.1	2.6	2.0	1.5	1.2	0.9	1.0
80	6.5	6.3	6.0	5.5	5.0	4.6	4.0	3.4	2.7	2.2	1.8	1.5	1.3
100	7.0	6.7	6.5	6.3	5.9	5.3	4.9	4.4	3.7	3.1	2.5	2.1	1.7
120	7.5	7.3	7.0	6.8	6.5	6.2	5.7	5.1	4.7	4.1	3.5	2.9	2.4
140	7.7	7.7	7.5	7.3	7.0	6.7	6.4	6.0	5.6	5.1	4.5	3.8	3.3
160	7.8	7.9	7.7	7.6	7.4	7.2	7.0	6.8	6.3	5.8	5.4	4.8	4.2
180	7.9	7.8	7.9	7.9	7.7	7.6	7.5	7.1	7.0	6.6	6.1	5.7	5.2
200	7.9	7.9	7.8	7.9	7.8	7.7	7.6	7.5	7.5	7.1	6.8	6.3	6.1
220	8.0	7.9	7.8	7.8	7.8	7.8	7.8	7.8	7.6	7.5	7.4	7.1	6.7
240	7.8	7.7	7.7	7.7	7.7	7.7	7.8	7.8	7.7	7.6	7.6	7.5	7.2
260	7.8	7.7	7.7	7.6	7.7	7.7	7.7	7.7	7.7	7.7	7.8	7.8	7.6
280	7.4	7.4	7.5	7.5	7.5	7.5	7.5	7.5	7.5	7.6	7.6	7.8	7.7
300	7.1	7.2	7.3	7.3	7.3	7.3	7.3	7.4	7.5	7.4	7.5	7.5	7.7
320	6.8	6.9	6.8	7.0	7.1	7.1	7.1	7.1	7.3	7.3	7.3	7.4	7.4
340	6.4	6.5	6.6	6.6	6.7	6.7	6.8	6.9	7.0	7.1	7.2	7.2	7.2
360	6.2	6.2	6.2	6.3	6.4	6.4	6.5	6.6	6.7	6.7	6.9	6.9	7.1
380	5.9	5.8	5.8	5.9	6.0	6.1	6.2	6.3	6.4	6.4	6.4	6.6	6.8
400	5.6	5.6	5.6	5.7	5.7	5.7	5.8	5.9	5.9	6.0	6.1	6.2	6.4
420	5.4	5.4	5.5	5.5	5.5	5.5	5.5	5.5	5.6	5.6	5.6	5.7	5.8
440	5.6	5.3	5.3	5.3	5.3	5.2	5.2	5.2	5.2	5.1	5.0	5.3	5.5
460	6.0	5.6	5.4	5.3	5.2	5.2	5.1	5.0	5.1	5.2	5.2	5.2	5.3
480	6.5	6.0	5.7	5.4	5.2	5.2	5.1	4.9	4.9	4.9	4.9	5.0	5.0
500	7.2	6.8	6.3	5.9	5.6	5.3	5.0	4.8	4.9	4.8	4.8	4.8	4.9
520	8.0	7.4	7.0	6.5	6.1	5.5	5.4	5.1	4.9	4.7	4.7	4.7	4.8
540	9.0	8.4	7.8	7.3	6.7	6.3	5.8	5.4	5.2	4.9	4.7	4.7	4.7
560	9.9	9.5	8.8	8.2	7.7	7.1	6.5	6.0	5.7	5.3	5.0	4.8	4.6
580	10.6	10.2	9.8	9.3	8.8	8.1	7.2	6.8	6.4	6.0	5.6	5.1	4.9
600	11.2	11.0	10.7	10.3	9.6	9.1	8.5	7.7	7.1	6.4	6.1	5.6	5.3
620	11.7	11.5	11.4	11.0	10.6	9.9	9.5	8.9	8.1	7.4	6.8	6.3	5.9
640	11.8	11.9	11.8	11.7	11.3	11.0	10.4	9.8	9.3	8.5	7.8	7.1	6.6
660	11.5	11.8	12.0	12.1	11.9	11.6	11.2	10.8	10.2	9.6	8.9	8.2	7.5
680	11.0	11.6	12.1	12.2	12.1	12.2	12.1	11.5	11.1	10.6	10.1	9.2	8.5
700	10.1	10.9	11.6	12.1	12.4	12.3	12.3	12.3	11.9	11.4	10.8	10.4	9.7
720	9.1	10.0	10.6	11.4	11.9	12.4	12.6	12.5	12.4	12.0	11.6	11.2	10.8
740	7.8	8.8	9.7	10.5	11.3	11.8	12.3	12.8	12.6	12.6	12.3	11.9	11.5
760	6.6	7.6	8.5	9.4	10.3	11.0	11.7	12.1	12.6	12.8	12.7	12.5	12.1
780	5.2	6.3	7.1	8.1	9.2	10.1	10.7	11.6	12.0	12.4	12.8	12.9	12.8
800	4.0	4.8	5.7	6.7	7.7	8.7	9.7	10.5	11.3	11.9	12.3	12.5	12.9
820	2.9	3.6	4.4	5.4	6.4	7.3	8.4	9.5	10.3	11.0	11.7	12.1	12.5
840	2.2	2.7	3.3	4.0	4.9	6.0	7.0	8.0	9.1	10.0	10.8	11.4	12.0
860	1.6	1.6	2.2	2.9	3.6	4.6	5.6	6.6	7.6	8.6	9.6	10.5	11.2
880	1.2	1.3	1.5	1.9	2.6	3.3	4.1	5.2	6.1	7.1	8.2	9.2	10.1
900	1.1	1.1	1.2	1.3	1.7	2.2	2.9	3.8	4.8	5.7	6.8	7.9	8.8
920	1.1	1.0	1.0	1.1	1.1	1.4	1.9	2.6	3.4	4.4	5.3	6.3	7.4
940	1.5	1.1	0.8	0.9	1.0	1.1	1.3	1.6	2.3	3.1	3.9	5.0	5.9
960	2.3	1.7	1.3	0.9	0.7	0.8	0.9	1.2	1.4	2.0	2.8	3.5	4.6
980	3.0	2.5	1.9	1.4	1.2	1.0	0.8	0.9	1.2	1.4	1.7	2.4	3.3
1000	3.7	3.2	2.6	2.1	1.7	1.3	0.9	0.7	0.7	1.0	1.2	1.6	2.2
	800	810	820	830	840	850	860	870	880	890	900	910	920



*Perturbations by Mars.**Pert's. by Jupiter*

Arguments II. and IV.

Arg's. II. and V.

IV.

V.

II.	920	930	940	950	960	970	980	990	1000	0	10	20	30
	"	"	"	"	"	"	"	"	"	"	"	"	"
0	2.2	3.0	3.8	4.8	5.8	6.9	7.8	8.4	9.5	15.3	15.1	15.0	15.0
20	1.5	2.1	2.6	3.4	4.4	5.5	6.5	7.6	8.7	14.9	14.9	14.7	14.8
40	1.0	1.4	1.8	2.5	3.2	4.0	5.2	6.0	7.1	14.7	14.6	14.6	14.5
60	1.0	1.1	1.3	1.8	2.3	3.0	3.7	4.8	5.8	14.4	14.4	14.4	14.4
80	1.3	1.1	1.2	1.4	1.6	2.2	2.7	3.6	4.5	13.4	13.9	14.0	14.2
100	1.7	1.3	1.2	1.2	1.3	1.6	2.0	2.6	3.3	13.2	13.4	13.6	13.7
120	2.4	2.0	1.5	1.4	1.4	1.4	1.7	1.9	2.4	12.3	12.7	13.0	13.3
140	3.3	2.8	2.3	2.0	1.7	1.5	1.5	1.8	2.1	11.3	11.8	12.1	12.5
160	4.2	3.6	3.1	2.6	2.1	2.0	1.7	1.7	1.9	10.2	10.7	11.2	11.7
180	5.2	4.6	4.0	3.5	3.1	2.5	2.0	2.0	1.9	9.1	9.6	10.1	10.7
200	6.1	5.5	5.0	4.4	3.9	3.5	2.8	2.7	2.9	7.8	8.3	8.9	9.5
220	6.7	6.3	5.8	5.4	4.9	4.4	3.9	3.2	3.0	6.8	7.2	7.7	8.3
240	7.2	6.9	6.6	6.1	5.6	5.3	4.8	4.2	3.7	5.7	6.2	6.6	7.2
260	7.6	7.5	7.1	6.8	6.5	6.0	5.6	5.2	4.8	4.8	5.2	5.6	6.1
280	7.7	7.7	7.5	7.3	7.1	6.7	6.3	5.9	5.5	3.9	4.1	4.7	5.2
300	7.7	7.7	7.7	7.7	7.4	7.2	7.0	6.6	6.1	3.4	3.5	3.9	4.3
320	7.4	7.4	7.6	7.7	7.6	7.6	7.3	7.1	6.9	3.2	3.1	3.4	3.6
340	7.2	7.2	7.3	7.5	7.7	7.6	7.6	7.6	7.7	3.2	3.0	3.0	3.1
360	7.1	7.1	7.1	7.2	7.2	7.6	7.6	7.6	7.5	3.5	3.2	2.9	2.9
380	6.8	6.9	7.0	7.0	7.0	7.1	7.3	7.5	7.5	4.5	4.0	3.4	3.1
400	6.4	6.6	6.6	6.7	6.7	6.9	7.0	7.1	7.3	5.0	4.3	3.8	3.5
420	5.8	5.9	6.2	6.3	6.6	6.5	6.7	6.7	6.9	6.1	5.2	4.6	4.1
440	5.5	5.6	5.7	5.8	6.0	6.1	6.3	6.5	6.5	7.5	6.6	5.8	4.9
460	5.3	5.3	5.4	5.7	5.7	5.7	5.9	6.1	6.2	9.0	7.9	7.0	6.3
480	5.0	5.0	5.0	5.1	5.3	5.4	5.5	5.6	5.8	10.5	9.5	8.5	7.6
500	4.9	4.9	5.0	5.0	5.0	5.1	5.2	5.3	5.3	12.3	11.3	10.0	9.1
520	4.8	4.8	4.8	4.8	4.8	4.7	4.9	5.0	5.1	14.0	12.7	11.7	10.7
540	4.7	4.7	4.6	4.6	4.6	4.5	4.6	4.6	4.7	15.6	14.5	13.3	12.3
560	4.6	4.5	4.5	4.4	4.5	4.5	4.5	4.5	4.4	17.1	16.1	15.1	14.0
580	4.9	4.7	4.6	4.5	4.4	4.4	4.4	4.4	4.2	18.6	17.4	16.5	15.7
600	5.3	4.9	4.8	4.7	4.5	4.4	4.4	4.3	4.1	19.8	19.0	17.9	17.0
620	5.9	5.5	5.1	4.8	4.6	4.5	4.4	4.3	4.2	20.8	20.1	19.2	18.4
640	6.6	6.1	5.6	5.4	5.0	4.7	4.6	4.5	4.3	21.6	20.9	20.2	19.5
660	7.5	6.8	6.3	5.9	5.5	5.3	4.9	4.8	4.6	22.1	21.6	21.0	20.4
680	8.5	7.8	7.3	6.5	6.1	5.6	5.4	5.1	4.8	22.3	22.0	21.6	21.2
700	9.7	8.9	8.1	7.6	7.0	6.3	5.9	5.6	5.3	22.2	22.0	21.7	21.5
720	10.8	10.0	9.3	8.5	7.9	7.2	6.6	6.1	5.8	22.0	21.9	21.7	21.6
740	11.5	11.0	10.2	9.7	8.9	8.2	7.6	6.9	6.5	21.6	21.6	21.5	21.5
760	12.1	11.8	11.3	10.5	10.0	9.3	8.5	7.9	7.3	21.2	21.1	21.1	21.2
780	12.8	12.3	11.9	11.4	10.9	10.2	9.6	9.0	8.2	20.4	20.5	20.6	20.7
800	12.9	12.9	12.5	12.1	11.7	11.2	10.5	9.8	9.2	19.6	19.8	19.9	20.1
820	12.5	12.7	12.8	12.7	12.2	11.9	11.2	10.7	10.1	18.8	19.0	19.2	19.4
840	12.0	12.4	12.6	12.8	12.6	12.4	12.2	11.5	10.9	18.1	18.2	18.4	18.6
860	11.2	11.8	12.3	12.5	12.7	12.5	12.5	12.3	11.7	17.4	17.5	17.6	17.9
880	10.1	11.0	11.5	12.1	12.3	12.6	12.6	12.4	12.3	16.9	16.9	16.9	17.1
900	8.8	9.8	10.6	11.3	11.8	12.2	12.4	12.5	12.4	16.3	16.4	16.4	16.5
920	7.4	8.4	9.3	10.2	11.0	11.5	12.1	12.2	12.3	16.0	15.9	15.9	16.0
940	5.9	7.1	8.1	8.9	9.9	10.7	11.2	11.7	12.1	15.8	15.7	15.7	15.6
960	4.6	5.6	6.7	7.7	8.7	9.4	10.2	10.9	11.4	15.5	15.4	15.3	15.4
980	3.3	4.2	5.2	6.2	7.3	8.2	8.9	9.9	10.6	15.3	15.2	15.2	15.1
1000	2.2	3.0	3.8	4.8	5.8	6.9	7.8	8.7	9.5	15.3	15.1	15.0	15.0
	920	930	940	950	960	970	980	990	1000	0	10	20	30



*Perturbations produced by Jupiter.*

Arguments II. and V.

V.

II.	30	40	50	60	70	80	90	100	110	120	130	140	150
	"	"	"	"	"	"	"	"	"	"	"	"	"
0	15.0	14.8	14.7	14.7	14.6	14.5	14.5	14.4	14.5	14.5	14.6	14.7	14.8
20	14.8	14.7	14.6	14.4	14.4	14.2	14.2	14.1	14.1	14.1	14.1	14.1	14.2
40	14.5	14.4	14.4	14.3	14.2	14.1	13.9	13.8	13.8	13.8	13.8	13.8	13.7
60	14.4	14.3	14.3	14.2	14.1	13.9	13.8	13.6	13.5	13.5	13.5	13.4	13.3
80	14.2	14.2	14.1	14.5	14.0	13.8	13.7	13.5	13.4	13.2	13.1	13.0	13.1
100	13.7	13.7	13.9	13.9	13.8	13.7	13.6	13.5	13.4	13.2	13.0	12.8	12.7
120	13.3	13.4	13.4	13.5	13.6	13.5	13.5	13.3	13.3	13.2	13.0	12.8	12.6
140	12.5	12.8	13.0	13.1	13.2	13.2	13.3	13.2	13.1	13.0	12.9	12.8	12.6
160	11.7	12.0	12.4	12.6	12.7	12.8	12.9	12.9	13.0	12.9	12.8	12.7	12.5
180	10.7	11.1	11.6	11.9	12.2	12.3	12.5	12.5	12.6	12.7	12.8	12.6	12.5
200	9.5	10.0	10.6	11.0	11.5	11.7	11.9	12.2	12.2	12.3	12.4	12.3	12.3
220	8.3	8.8	9.5	9.9	10.4	10.8	11.3	11.5	11.8	11.9	12.0	12.0	12.0
240	7.2	7.7	8.2	8.9	9.4	9.8	10.3	10.6	11.0	11.3	11.5	11.7	11.8
260	6.1	6.5	7.1	7.6	8.3	8.8	9.3	9.7	10.1	10.5	10.9	11.0	11.2
280	5.2	5.5	6.0	6.5	7.1	7.6	8.2	8.7	9.2	9.6	10.0	10.4	10.6
300	4.3	4.7	5.1	5.5	6.1	6.6	7.1	7.6	8.1	8.7	9.1	9.4	9.9
320	3.6	3.9	4.3	4.6	5.1	5.4	6.0	6.6	7.2	7.7	8.1	8.5	8.9
340	3.1	3.3	3.5	3.8	4.1	4.5	5.0	5.4	6.1	6.6	7.2	7.6	8.0
360	2.9	3.0	3.1	3.3	3.6	3.8	4.1	4.5	5.0	5.5	6.1	6.6	7.1
380	3.1	2.8	2.8	2.7	2.8	2.9	3.0	3.2	3.5	4.1	4.6	5.0	5.6
400	3.5	3.1	2.9	2.9	2.8	2.8	3.0	3.1	3.4	3.8	4.2	4.7	5.2
420	4.1	3.6	3.3	3.1	2.8	2.7	2.8	2.9	3.1	3.2	3.5	3.8	4.3
440	4.9	4.4	3.9	3.4	3.1	2.7	2.8	2.7	2.8	3.1	3.1	3.2	3.5
460	6.3	5.4	4.8	4.3	3.7	3.2	2.9	2.8	2.8	2.7	2.7	2.8	3.2
480	7.6	6.7	5.9	5.2	4.6	4.1	3.6	3.1	3.0	2.8	2.8	2.6	2.7
500	9.1	8.1	7.2	6.4	5.7	5.0	4.4	3.9	3.4	3.2	3.1	2.9	2.7
520	10.7	9.5	8.7	7.7	6.9	6.1	5.5	4.8	4.2	3.8	3.5	3.2	3.1
540	12.3	11.1	10.2	9.1	8.4	7.4	6.6	5.9	5.3	4.7	4.1	3.8	3.5
560	14.0	13.0	11.9	10.8	9.9	8.7	7.9	7.1	6.4	5.8	5.2	4.5	4.1
580	15.7	14.5	13.6	12.5	11.4	10.4	9.3	8.3	7.7	6.9	6.2	5.5	5.0
600	17.0	16.0	15.0	14.0	13.1	12.0	11.0	10.1	9.2	8.2	7.5	6.7	6.0
620	18.4	17.4	16.5	15.5	14.7	13.6	12.6	11.6	10.7	9.8	9.0	8.0	7.3
640	19.5	18.5	17.9	17.0	16.0	15.1	14.2	13.1	12.2	11.3	10.8	9.4	8.7
660	20.4	19.7	18.9	18.1	17.4	16.3	15.6	14.6	13.7	12.8	11.9	11.0	10.1
680	21.2	20.5	19.9	19.1	18.5	17.6	16.8	16.0	15.1	14.2	13.5	12.5	11.6
700	21.5	21.0	20.6	20.0	19.3	18.7	18.0	17.1	16.5	15.6	14.7	13.8	13.0
720	21.6	21.2	21.0	20.5	20.0	19.3	18.9	18.3	17.5	16.8	16.1	15.1	14.3
740	21.5	21.2	21.1	20.8	20.5	20.0	19.4	18.9	18.4	17.7	17.2	16.8	15.7
760	21.2	21.0	21.0	20.8	20.7	20.3	20.0	19.4	19.0	18.6	17.9	17.4	16.7
780	20.7	20.7	20.7	20.6	20.6	20.3	20.2	19.8	19.4	19.1	18.7	18.1	17.6
800	20.1	20.2	20.3	20.3	20.4	20.3	20.1	19.9	19.7	19.3	19.1	18.7	18.2
820	19.4	19.5	19.7	19.8	19.9	19.9	19.9	19.8	19.8	19.6	19.2	18.9	18.7
840	18.6	18.8	18.9	19.0	19.2	19.3	19.4	19.4	19.4	19.4	19.4	19.0	18.9
860	17.9	18.0	18.3	18.4	18.6	18.7	18.8	18.9	19.0	19.1	19.1	19.0	18.8
880	17.1	17.2	17.5	17.6	17.9	18.0	18.2	18.3	18.5	18.6	18.6	18.6	18.7
900	16.5	16.6	16.8	16.9	17.1	17.1	17.4	17.5	17.7	17.9	18.1	18.2	18.2
920	16.0	16.0	16.1	16.2	16.4	16.5	16.7	16.8	17.0	17.2	17.4	17.5	17.7
940	15.6	15.5	15.6	15.6	15.7	15.8	16.0	16.1	16.3	16.5	16.8	16.8	17.1
960	15.4	15.3	15.3	15.2	15.2	15.2	15.3	15.4	15.6	15.7	15.9	16.0	16.3
980	15.1	15.0	15.0	14.9	14.9	14.8	14.9	14.9	14.9	15.0	15.2	15.3	15.5
1000	15.0	14.8	14.7	14.7	14.6	14.5	14.5	14.4	14.5	14.5	14.6	14.7	14.8
	30	40	50	60	70	80	90	100	110	120	130	140	150

*Perturbations produced by Jupiter.*

Arguments II. and V.

V.

II.	150	160	170	180	190	200	210	220	230	240	250	260	270
	"	"	"	"	"	"	"	"	"	"	"	"	"
0	14.8	15.0	15.3	15.5	15.8	15.9	16.2	16.3	16.7	17.0	17.1	17.3	17.5
20	14.2	14.3	14.6	14.8	14.9	15.2	15.5	15.7	15.9	16.2	16.6	16.8	17.1
40	13.7	13.7	13.9	14.1	14.3	14.5	14.8	15.0	15.3	15.5	15.8	16.2	16.4
60	13.3	13.2	13.4	13.5	13.6	13.8	14.1	14.3	14.6	14.8	15.1	15.5	15.8
80	13.1	13.0	13.0	13.0	13.1	13.1	13.3	13.5	13.8	14.1	14.4	14.5	15.1
100	12.7	12.7	12.7	12.6	12.7	12.6	12.8	12.9	13.1	13.4	13.7	14.0	14.2
120	12.6	12.5	12.5	12.4	12.3	12.2	12.3	12.3	12.6	12.8	13.0	13.3	13.6
140	12.6	12.4	12.4	12.3	12.1	12.0	12.0	12.0	12.1	12.1	12.3	12.5	12.8
160	12.5	12.3	12.2	12.1	12.1	11.9	11.8	11.8	11.8	11.8	11.9	12.0	12.2
180	12.5	12.3	12.2	12.1	11.9	11.8	11.7	11.5	11.5	11.5	11.6	11.7	11.8
200	12.3	12.2	12.2	12.0	11.9	11.7	11.7	11.5	11.4	11.3	11.2	11.3	11.5
220	12.0	12.0	12.1	12.0	11.8	11.6	11.6	11.5	11.4	11.3	11.2	11.1	11.1
240	11.8	11.8	11.9	11.9	11.8	11.6	11.5	11.4	11.3	11.2	11.1	11.1	11.0
260	11.2	11.5	11.6	11.6	11.6	11.5	11.3	11.3	11.3	11.2	11.1	11.0	10.9
280	10.6	10.8	11.1	11.2	11.2	11.2	11.3	11.3	11.2	11.2	11.1	11.0	10.9
300	9.9	10.1	10.5	10.8	10.9	11.0	11.1	11.0	11.0	11.0	11.0	11.1	10.9
320	8.9	9.4	9.7	10.1	10.4	10.5	10.7	10.8	10.8	10.8	10.8	10.8	10.9
340	8.0	8.5	9.1	9.3	9.6	9.9	10.2	10.3	10.5	10.6	10.6	10.7	10.7
360	7.1	7.5	8.0	8.4	8.9	9.2	9.5	9.8	10.1	10.3	10.4	10.5	10.5
380	5.6	6.2	6.8	7.3	7.8	8.3	8.9	9.3	9.7	10.0	10.0	10.1	10.2
400	5.2	5.6	6.2	6.6	7.0	7.5	7.9	8.4	8.8	9.1	9.4	9.7	9.9
420	4.3	4.8	5.3	5.8	6.2	6.6	7.1	7.4	7.9	8.4	8.7	9.1	9.4
440	3.5	3.9	4.4	4.9	5.4	5.7	6.2	6.7	7.1	7.6	7.9	8.4	8.7
460	3.2	3.3	3.8	4.1	4.5	4.9	5.4	5.7	6.3	6.7	7.2	7.7	8.0
480	2.7	2.9	3.2	3.6	3.9	4.3	4.7	5.0	5.4	5.9	6.3	6.8	7.3
500	2.7	2.7	2.9	3.1	3.4	3.6	4.0	4.4	4.8	5.2	5.7	5.9	6.4
520	3.1	2.8	2.9	3.0	3.1	3.2	3.5	3.8	4.2	4.7	4.9	5.4	5.7
540	3.5	3.2	3.1	3.0	3.0	3.0	3.3	3.5	3.7	4.1	4.3	4.7	5.1
560	4.1	3.8	3.6	3.3	3.2	3.2	3.2	3.3	3.5	3.7	4.0	4.3	4.5
580	5.0	4.6	4.2	4.0	3.6	3.5	3.3	3.2	3.4	3.5	3.7	4.0	4.2
600	6.0	5.4	5.1	4.6	4.3	3.9	3.7	3.5	3.5	3.6	3.7	3.8	4.0
620	7.3	6.6	6.0	5.6	5.1	4.6	4.2	4.0	3.9	3.8	3.9	3.9	4.0
640	8.7	7.8	7.3	6.6	6.1	5.5	5.2	4.7	4.4	4.2	4.0	4.0	4.1
660	10.1	9.3	8.6	7.7	7.2	6.5	6.2	5.9	5.3	4.9	4.6	4.5	4.4
680	11.6	10.8	10.0	9.3	8.5	7.5	7.3	6.7	6.3	5.8	5.5	5.2	4.9
700	13.0	12.1	11.5	10.7	9.9	9.0	8.5	7.8	7.4	6.9	6.3	6.0	5.8
720	14.3	13.5	12.8	12.1	11.3	10.6	9.8	9.1	8.7	8.0	7.6	7.0	6.6
740	15.7	14.9	14.2	13.4	12.7	12.0	11.2	10.5	9.7	9.3	8.9	8.2	7.7
760	16.7	15.9	15.5	14.7	13.9	13.3	12.6	11.8	11.2	10.5	10.0	9.5	9.0
780	17.6	17.0	16.4	15.7	15.1	14.6	13.8	13.2	12.6	11.9	11.2	10.8	10.2
800	18.2	17.8	17.3	16.8	16.2	16.0	15.0	14.3	13.7	13.1	12.6	12.0	11.5
820	18.7	18.3	18.0	17.6	17.0	16.6	16.0	15.3	14.9	14.3	13.7	13.1	12.6
840	18.9	18.7	18.4	18.2	17.7	17.2	16.8	16.3	15.8	15.3	14.9	14.4	13.8
860	18.8	18.7	18.6	18.4	18.3	17.9	17.4	17.1	16.7	16.3	15.9	15.4	15.0
880	18.7	18.5	18.6	18.5	18.3	18.2	18.0	17.7	17.4	17.1	16.6	16.3	15.9
900	18.2	18.2	18.3	18.3	18.3	18.1	18.1	18.0	17.8	17.6	17.3	17.0	16.7
920	17.7	17.9	18.0	18.0	18.1	18.1	18.0	18.0	18.0	17.8	17.7	17.6	17.3
940	17.1	17.1	17.4	17.6	17.6	17.7	17.8	17.8	17.9	18.0	17.8	17.8	17.7
960	16.3	16.5	16.8	16.9	17.1	17.2	17.4	17.5	17.6	17.8	17.9	18.0	17.9
980	15.5	15.7	16.1	16.3	16.5	16.7	16.8	17.0	17.2	17.3	17.6	17.7	17.9
1000	14.8	15.0	15.3	15.5	15.8	15.9	16.2	16.3	16.7	17.0	17.1	17.3	17.5
	150	160	170	180	190	200	210	220	230	240	250	260	270

*Perturbations produced by Jupiter.*

Arguments II. and V

V

II.	270	280	290	300	310	320	330	340	350	360	370	380	390
0	17.5	17.5	17.7	17.8	17.9	17.9	18.0	18.0	17.9	17.7	17.6	17.5	17.5
20	17.1	17.3	17.5	17.6	17.8	17.8	18.0	18.1	18.1	18.1	18.0	18.0	18.0
40	16.4	16.8	16.9	17.2	17.6	17.7	17.9	18.1	18.3	18.3	18.4	18.4	18.6
60	15.8	16.0	16.4	16.7	16.9	17.3	17.6	17.9	18.2	18.3	18.5	18.5	18.7
80	15.1	15.4	15.7	16.1	16.4	16.7	17.0	17.5	17.8	18.0	18.3	18.5	18.8
100	14.2	14.6	15.1	15.0	15.8	16.1	16.5	17.0	17.2	17.5	17.9	18.3	18.7
120	13.6	13.7	14.2	14.5	15.0	15.4	15.8	16.2	16.7	17.1	17.3	17.9	18.3
140	12.8	13.1	13.3	13.7	14.2	14.4	15.1	15.5	15.9	16.3	16.8	17.3	17.7
160	12.2	12.4	12.6	12.9	13.4	13.8	14.1	14.6	15.2	15.5	16.0	16.5	17.1
180	11.8	11.9	12.1	12.3	12.5	12.8	13.3	13.7	14.4	14.7	15.2	15.7	16.3
200	11.5	11.5	11.6	11.7	12.0	12.1	12.5	13.0	13.4	13.8	14.3	14.7	15.5
220	11.1	11.1	11.2	11.3	11.6	11.7	11.9	12.3	12.7	13.0	13.5	14.0	14.5
240	11.0	10.9	10.9	11.0	11.2	11.3	11.5	11.8	12.1	12.3	12.8	13.2	13.8
260	10.9	10.8	10.8	10.8	10.9	10.9	11.1	11.3	11.4	11.6	12.0	12.3	13.0
280	10.9	10.8	10.7	10.6	10.7	10.6	10.8	11.0	11.2	11.3	11.5	11.8	12.2
300	10.9	10.8	10.7	10.6	10.6	10.5	10.6	10.7	10.8	10.9	11.1	11.4	11.8
320	10.9	10.7	10.7	10.6	10.6	10.5	10.5	10.6	10.7	10.6	10.7	11.0	11.2
340	10.7	10.7	10.6	10.5	10.5	10.4	10.5	10.5	10.6	10.5	10.6	10.7	10.8
360	10.5	10.5	10.5	10.5	10.5	10.4	10.4	10.4	10.4	10.3	10.5	10.6	10.8
380	10.2	10.3	10.3	10.3	10.4	10.3	10.4	10.4	10.4	10.3	10.3	10.4	10.6
400	9.9	10.0	10.0	10.2	10.3	10.2	10.2	10.3	10.4	10.3	10.3	10.3	10.5
420	9.4	9.6	9.8	9.9	10.1	10.2	10.1	10.2	10.2	10.2	10.3	10.3	10.4
440	8.7	9.0	9.2	9.4	9.7	9.8	10.0	10.1	10.2	10.1	10.1	10.2	10.4
460	8.0	8.4	8.6	8.8	9.1	9.3	9.6	9.9	10.1	10.0	10.0	10.2	10.3
480	7.3	7.6	7.9	8.4	8.7	8.9	9.1	9.4	9.6	9.7	9.8	10.0	10.1
500	6.4	6.9	7.2	7.6	8.0	8.3	8.6	8.9	9.2	9.4	9.5	9.7	9.9
520	5.7	6.1	6.6	6.9	7.3	7.6	7.9	8.3	8.6	8.9	9.1	9.4	9.7
540	5.1	5.4	5.8	6.2	6.7	7.0	7.4	7.7	8.0	8.3	8.6	8.9	9.2
560	4.5	4.9	5.1	5.5	6.0	6.3	6.7	7.2	7.5	7.7	8.0	8.3	8.7
580	4.2	4.4	4.8	5.0	5.3	5.7	6.1	6.6	6.9	7.1	7.4	7.7	8.1
600	4.0	4.2	4.3	4.7	4.9	5.2	5.6	6.0	6.3	6.5	6.8	7.2	7.6
620	4.0	4.0	4.1	4.3	4.7	4.8	5.1	5.5	5.8	6.1	6.4	6.7	7.0
640	4.1	4.1	4.2	4.2	4.4	4.6	4.8	5.1	5.4	5.6	5.9	6.3	6.6
660	4.4	4.3	4.3	4.3	4.5	4.5	4.7	4.9	5.1	5.3	5.5	5.8	6.2
680	4.9	4.9	4.7	4.6	4.7	4.5	4.6	4.8	5.0	5.1	5.3	5.5	5.8
700	5.8	5.4	5.2	5.1	5.0	4.9	4.9	4.9	5.1	5.2	5.3	5.4	5.6
720	6.6	6.2	5.9	5.7	5.6	5.5	5.4	5.3	5.3	5.3	5.3	5.4	5.5
740	7.7	7.2	6.8	6.5	6.4	6.1	6.0	5.9	5.8	5.7	5.6	5.5	5.7
760	9.0	8.2	7.9	7.5	7.2	6.9	6.7	6.5	6.3	6.1	5.9	5.9	6.0
780	10.2	9.7	9.1	8.4	8.2	7.7	7.6	7.4	7.2	6.9	6.6	6.5	6.5
800	11.5	11.0	10.4	9.8	9.4	8.7	8.5	8.3	8.0	7.7	7.6	7.3	7.1
820	12.6	12.1	11.7	11.2	10.6	10.1	9.7	9.2	9.1	8.6	8.3	8.1	7.9
840	13.8	13.2	12.8	12.3	11.9	11.3	10.9	10.5	10.2	9.6	9.4	9.1	8.9
860	15.0	14.4	13.8	13.5	13.1	12.6	12.1	11.7	11.2	10.7	10.4	10.1	10.0
880	15.9	15.4	15.0	14.4	14.2	13.7	13.4	12.9	12.5	12.0	11.5	11.3	11.1
900	16.7	16.4	15.9	15.5	15.2	14.8	14.4	14.1	13.7	13.2	12.8	12.4	12.2
920	17.3	17.1	16.8	16.5	16.2	15.7	15.5	15.2	14.8	14.3	14.0	13.6	13.3
940	17.7	17.5	17.3	17.1	16.9	16.6	16.3	16.1	16.0	15.5	15.0	14.7	14.5
960	17.9	17.8	17.6	17.5	17.4	17.2	17.0	16.9	16.8	16.4	16.2	15.8	15.6
980	17.9	17.8	17.8	17.8	17.8	17.8	17.6	17.5	17.3	17.2	17.0	16.8	16.6
1000	17.5	17.7	17.7	17.8	17.9	17.9	18.0	18.0	17.9	17.7	17.6	17.5	17.5
	270	280	290	300	310	320	330	340	350	360	370	380	390

*Perturbations produced by Jupiter.*

Arguments II. and V.

V.

II.	390	400	410	420	430	440	450	460	470	480	490	500	510
	"	"	"	"	"	"	"	"	"	"	"	"	"
0	17.5	17.1	17.0	16.7	16.5	16.3	16.1	15.8	15.6	15.1	14.6	14.3	13.9
20	18.0	18.1	17.7	17.5	17.5	17.2	17.1	16.8	16.7	16.3	16.0	15.6	15.3
40	18.6	18.6	18.5	18.4	18.3	18.1	18.0	17.8	17.6	17.3	17.2	16.8	16.5
60	18.7	18.9	18.9	18.9	18.9	18.7	18.8	18.6	18.7	18.4	18.1	17.9	17.7
80	18.8	18.9	19.2	19.3	19.4	19.3	19.3	19.3	19.3	19.2	19.2	18.9	18.8
100	18.7	18.9	19.1	19.4	19.7	19.8	19.8	19.8	19.8	19.8	19.9	19.7	19.7
120	18.3	18.6	18.9	19.2	19.5	19.8	20.0	20.1	20.3	20.3	20.4	20.4	20.4
140	17.7	18.2	18.6	18.9	19.2	19.6	20.0	20.3	20.5	20.6	20.7	20.8	21.0
160	17.1	17.6	17.9	18.5	19.0	19.3	19.8	20.2	20.5	20.6	20.9	21.1	21.2
180	16.3	16.8	17.3	17.9	18.3	18.8	19.3	19.8	20.3	20.6	20.9	21.1	21.4
200	15.5	16.0	16.5	17.1	17.7	18.2	18.6	19.1	19.8	20.2	20.7	21.0	21.4
220	14.5	15.0	15.6	16.1	16.9	17.4	18.0	18.6	19.0	19.7	20.3	20.7	21.1
240	13.8	14.2	14.7	15.2	15.9	16.5	17.1	17.7	18.4	18.9	19.5	20.1	20.7
260	13.0	13.4	13.9	14.4	15.0	15.5	16.3	16.9	17.5	18.0	18.6	19.3	20.0
280	12.2	12.7	13.0	13.5	14.2	14.7	15.3	15.9	16.7	17.2	17.8	18.4	19.1
300	11.8	11.9	12.4	12.8	13.3	13.8	14.4	14.9	15.7	16.3	17.0	17.6	18.2
320	11.2	11.5	11.8	12.2	12.7	13.0	13.6	14.1	14.7	15.3	16.0	16.6	17.4
340	10.8	11.2	11.4	11.6	12.1	12.4	12.9	13.4	13.9	14.4	15.1	15.7	16.4
360	10.8	10.8	11.0	11.2	11.6	11.9	12.3	12.6	13.2	13.6	14.2	14.8	15.5
380	10.6	10.6	10.7	10.9	11.2	11.4	11.9	12.2	12.6	12.9	13.5	13.9	14.5
400	10.5	10.5	10.6	10.6	10.9	11.1	11.4	11.8	12.2	12.5	12.9	13.3	13.8
420	10.4	10.4	10.5	10.6	10.7	10.9	11.2	11.3	11.7	11.9	12.4	12.8	13.3
440	10.4	10.4	10.4	10.5	10.7	10.8	10.9	11.1	11.3	11.6	11.9	12.2	12.7
460	10.3	10.4	10.4	10.4	10.6	10.6	10.7	10.9	11.2	11.3	11.7	11.9	12.2
480	10.1	10.2	10.3	10.4	10.6	10.6	10.7	10.8	11.0	11.2	11.4	11.7	12.0
500	9.9	10.0	10.1	10.2	10.4	10.5	10.7	10.8	10.9	11.0	11.2	11.3	11.7
520	9.7	9.8	9.8	10.0	10.2	10.3	10.5	10.6	10.9	10.8	11.1	11.3	11.5
540	9.2	9.4	9.6	9.8	10.0	10.2	10.3	10.4	10.6	10.7	10.9	11.1	11.4
560	8.7	8.9	9.1	9.3	9.7	9.8	10.1	10.3	10.5	10.6	10.7	10.8	11.2
580	8.1	8.5	8.7	8.7	9.2	9.4	9.7	9.9	10.2	10.4	10.6	10.7	10.9
600	7.6	7.9	8.2	8.5	8.8	9.0	9.3	9.5	9.8	10.0	10.3	10.5	10.7
620	7.0	7.3	7.6	7.9	8.2	8.5	8.8	9.0	9.4	9.6	10.0	10.1	10.4
640	6.6	6.8	7.1	7.4	7.7	7.9	8.2	8.6	8.9	9.1	9.4	9.7	10.1
660	6.2	6.4	6.6	6.9	7.3	7.6	7.9	8.1	8.3	8.6	8.9	9.2	9.5
680	5.8	6.1	6.2	6.5	6.8	7.0	7.4	7.6	7.9	8.1	8.4	8.7	9.0
700	5.6	5.8	6.0	6.2	6.4	6.6	6.9	7.1	7.4	7.6	7.9	8.2	8.5
720	5.5	5.6	5.7	5.9	6.2	6.3	6.5	6.8	7.1	7.2	7.5	7.7	8.0
740	5.7	5.7	5.7	5.8	6.0	6.1	6.2	6.4	6.7	6.9	7.1	7.2	7.5
760	6.0	6.0	6.0	6.0	6.0	6.1	6.2	6.3	6.4	6.5	6.7	6.8	7.1
780	6.5	6.3	6.2	6.2	6.3	6.3	6.3	6.3	6.4	6.4	6.5	6.7	6.8
800	7.1	7.0	6.7	6.6	6.7	6.5	6.5	6.4	6.5	6.5	6.5	6.6	6.7
820	7.9	7.6	7.5	7.3	7.2	7.0	7.0	6.8	6.8	6.7	6.6	6.6	6.7
840	8.9	8.6	8.3	8.1	7.8	7.7	7.6	7.4	7.3	7.1	7.0	6.8	6.8
860	10.0	9.7	9.3	9.0	8.7	8.4	8.2	8.1	7.9	7.7	7.6	7.3	7.2
880	11.1	10.5	10.4	10.0	9.7	9.5	9.2	8.9	8.7	8.4	8.2	7.9	7.7
900	12.2	11.8	11.5	11.0	10.8	10.5	10.3	9.9	9.7	9.4	9.0	8.8	8.5
920	13.3	13.0	12.6	12.3	12.1	11.5	11.3	11.0	10.6	10.2	10.1	9.7	9.4
940	14.5	14.1	13.8	13.5	13.2	12.8	12.5	11.9	11.8	11.3	11.0	10.7	10.4
960	15.6	15.3	14.9	14.6	14.4	14.0	13.7	13.3	13.0	12.5	12.1	11.8	11.5
980	16.6	16.3	16.0	15.7	15.6	15.2	14.9	14.6	14.2	13.8	13.6	12.9	12.7
1000	17.5	17.1	17.0	16.7	16.5	16.3	16.1	15.8	15.6	15.1	14.6	14.3	13.9
	390	400	410	420	430	440	450	460	470	480	490	500	510



*Perturbations produced by Jupiter.*

Arguments II. and V.

V.

II.	510	520	530	540	550	560	570	580	590	600	610	620	630
	"	"	"	"	"	"	"	"	"	"	"	"	"
0	13.9	13.4	13.1	12.7	12.1	11.8	11.3	10.8	10.2	9.9	9.4	8.9	8.4
20	15.3	14.9	14.4	13.9	13.5	13.1	12.5	12.1	11.5	11.0	10.4	10.0	9.4
40	16.5	16.3	15.7	15.4	15.0	14.3	13.8	13.4	12.8	12.3	11.7	11.1	10.5
60	17.7	17.3	17.0	16.6	16.1	15.8	15.3	14.7	14.3	13.7	13.0	12.4	11.8
80	18.8	18.5	18.1	17.9	17.4	17.1	16.6	16.2	15.7	15.1	14.5	13.9	13.2
100	19.7	19.5	19.2	19.0	18.8	18.4	17.9	17.6	17.0	16.5	16.0	15.2	14.7
120	20.4	20.3	20.2	20.0	19.7	19.5	19.1	18.8	18.4	18.0	17.3	16.8	16.2
140	21.0	21.1	21.0	20.8	20.7	20.4	20.2	19.9	19.6	19.3	18.8	18.3	17.7
160	21.2	21.5	21.5	21.6	21.5	21.3	21.2	21.0	20.6	20.4	20.1	19.6	19.1
180	21.4	21.6	21.8	22.0	22.0	22.1	21.9	21.8	21.6	21.4	21.1	20.7	20.3
200	21.4	21.7	21.9	22.1	22.3	22.5	22.5	22.5	22.4	22.3	22.1	21.8	21.5
220	21.1	21.5	21.8	22.2	22.5	22.8	23.1	23.1	22.9	22.8	22.9	22.6	22.5
240	20.7	21.1	21.5	21.9	22.3	22.7	23.0	23.3	23.4	23.5	23.4	23.3	23.2
260	20.0	20.6	21.0	21.6	22.0	22.4	22.8	23.2	23.5	23.8	23.8	23.8	23.9
280	19.1	19.9	20.4	20.9	21.5	22.0	22.4	23.0	23.3	23.7	24.0	24.1	24.1
300	18.2	19.0	19.6	20.3	20.7	21.3	21.8	22.3	23.0	23.4	23.8	24.1	24.3
320	17.4	18.9	18.7	19.4	20.0	20.6	21.1	21.8	22.3	22.9	23.3	23.7	24.2
340	16.4	17.0	17.6	18.5	19.2	19.9	20.4	21.1	21.6	22.2	22.8	23.3	23.7
360	15.5	16.2	16.7	17.4	18.2	18.9	19.5	20.1	20.8	21.5	22.0	22.6	23.2
380	14.5	15.2	15.9	16.6	17.1	17.9	18.6	19.3	19.8	20.5	21.1	21.8	22.5
400	13.8	14.4	14.9	15.6	16.2	16.8	17.6	18.4	19.1	19.7	20.3	20.9	21.5
420	13.3	13.7	14.2	14.8	15.3	16.0	16.5	17.4	18.0	18.7	19.4	20.0	20.6
440	12.7	13.1	13.6	14.1	14.6	15.2	15.7	16.4	17.1	17.8	18.4	18.9	19.6
460	12.2	12.7	13.0	13.5	13.9	14.4	15.0	15.6	16.1	16.9	17.5	18.2	18.7
480	12.0	12.2	12.5	13.0	13.4	13.9	14.3	14.8	15.3	15.9	16.6	17.3	17.9
500	11.7	12.0	12.2	12.6	12.9	13.3	13.8	14.3	14.7	15.2	15.7	16.4	16.9
520	11.5	11.9	12.0	12.3	12.6	13.0	13.2	13.8	14.2	14.7	15.1	15.5	16.2
540	11.4	11.6	11.9	12.2	12.4	12.7	12.9	13.3	13.7	14.2	14.6	15.0	15.4
560	11.2	11.4	11.5	11.9	12.1	12.4	12.7	13.1	13.4	13.8	14.1	14.5	14.9
580	10.9	11.2	11.4	11.6	11.9	12.2	12.4	12.8	13.1	13.5	13.8	14.2	14.5
600	10.7	10.8	11.1	11.5	11.7	12.0	12.2	12.5	12.8	13.1	13.4	13.8	14.2
620	10.4	10.7	10.7	11.1	11.4	11.6	12.0	12.3	12.5	12.9	13.1	13.4	13.8
640	10.1	10.4	10.6	10.7	11.0	11.3	11.6	12.0	12.3	12.6	12.9	13.2	13.5
660	9.5	9.9	10.2	10.5	10.6	11.0	11.3	11.6	11.9	12.3	12.6	12.9	13.2
680	9.0	9.3	9.6	10.0	10.3	10.5	10.8	11.3	11.5	11.9	12.2	12.4	12.8
700	8.5	8.9	9.1	9.5	9.8	10.1	10.3	10.7	11.1	11.4	11.8	12.1	12.4
720	8.0	8.3	8.5	9.0	9.2	9.6	9.9	10.2	10.5	10.9	11.3	11.7	12.0
740	7.5	7.8	8.0	8.3	8.6	9.0	9.3	9.7	9.9	10.4	10.8	11.1	11.5
760	7.1	7.3	7.5	7.9	8.1	8.4	8.6	9.1	9.4	9.7	10.1	10.5	10.9
780	6.8	7.0	7.1	7.3	7.6	7.9	8.1	8.5	8.8	9.2	9.4	9.8	10.2
800	6.7	6.8	6.8	7.0	7.1	7.3	7.5	7.8	8.2	8.5	8.8	9.1	9.5
820	6.7	6.8	6.6	6.8	6.9	7.0	7.1	7.4	7.6	7.9	8.1	8.4	8.7
840	6.8	6.8	6.8	6.8	6.8	6.9	6.9	7.1	7.2	7.4	7.6	7.9	8.1
860	7.2	7.1	7.1	7.0	6.9	6.9	6.8	6.8	6.9	7.1	7.2	7.3	7.6
880	7.7	7.5	7.4	7.3	7.1	7.0	6.8	6.8	6.7	6.8	6.8	7.0	7.2
900	8.5	8.2	7.9	7.7	7.5	7.3	7.2	7.1	6.9	6.9	6.8	6.8	6.8
920	9.4	9.2	8.7	8.4	8.1	7.9	7.6	7.4	7.1	7.0	6.9	6.8	6.7
940	10.4	10.0	9.7	9.4	8.9	8.6	8.3	8.1	7.7	7.4	7.1	6.9	6.7
960	11.5	11.2	10.7	10.4	9.8	9.5	9.1	8.8	8.5	8.1	7.7	7.4	7.1
980	12.7	12.3	11.8	11.5	11.1	10.6	10.0	9.7	9.2	8.9	8.5	8.1	7.7
1000	13.9	13.4	13.1	12.7	12.1	11.8	11.3	10.8	10.2	9.9	9.4	8.9	8.4
	510	520	530	540	550	560	570	580	590	600	610	620	630



*Perturbations produced by Jupiter.*

Arguments II. and V.

V.

II.	630	640	650	660	670	680	690	700	710	720	730	740	750
0	8.4	8.0	7.7	7.3	6.9	6.7	6.5	6.5	6.3	6.2	6.2	6.4	6.5
20	9.4	9.0	8.4	8.0	7.5	7.1	6.9	6.7	6.4	6.3	6.0	6.1	6.1
40	10.5	10.1	9.4	8.9	8.3	7.8	7.4	7.0	6.6	6.4	6.2	5.9	5.8
60	11.8	11.3	10.6	10.1	9.3	8.7	8.2	7.7	7.2	6.8	6.4	6.2	5.8
80	13.2	12.7	12.0	11.3	10.5	9.9	9.2	8.7	8.1	7.6	7.1	6.6	6.2
100	14.7	14.1	13.4	12.8	12.0	11.3	10.6	9.9	9.1	8.5	7.9	7.3	6.8
120	16.2	15.4	14.9	14.2	13.4	12.7	12.0	11.3	10.4	9.8	8.9	8.2	7.6
140	17.7	17.2	16.4	15.6	14.9	14.2	13.4	12.7	11.9	11.1	10.2	9.6	8.8
160	19.1	18.6	17.9	17.3	16.6	15.7	15.0	14.2	13.3	12.6	11.7	10.9	10.0
180	20.3	19.9	19.4	18.8	18.0	17.3	16.7	15.8	15.0	14.1	13.2	12.4	11.5
200	21.5	21.2	20.8	20.2	19.3	18.9	18.1	17.5	16.6	15.7	14.9	14.0	13.1
220	22.5	22.3	21.9	21.5	21.0	20.3	19.7	19.0	18.2	17.5	16.6	15.5	14.7
240	23.2	23.0	22.9	22.5	22.0	21.6	21.1	20.5	19.8	19.1	18.2	17.3	16.4
260	23.9	23.8	23.7	23.5	23.1	22.7	22.3	21.8	21.2	20.6	19.8	19.1	18.1
280	24.1	24.3	24.2	24.2	24.0	23.7	23.5	23.1	22.4	21.8	21.2	20.5	19.8
300	24.3	24.5	24.6	24.6	24.5	24.4	24.2	23.9	23.6	23.1	22.5	21.9	21.2
320	24.2	24.5	24.7	24.9	24.8	24.8	24.8	24.7	24.4	24.1	23.7	23.1	22.5
340	23.7	24.2	24.5	24.7	25.0	25.2	25.1	25.0	25.0	24.9	24.6	24.1	23.7
360	23.2	23.7	24.2	24.5	24.7	25.0	25.1	25.3	25.4	25.3	25.1	24.9	24.5
380	22.5	23.1	23.6	24.1	24.4	24.7	25.1	25.2	25.4	25.5	25.4	25.3	25.2
400	21.5	22.3	22.8	23.4	23.9	24.3	24.7	25.1	25.2	25.4	25.6	25.6	25.5
420	20.6	21.3	22.0	22.6	23.1	23.6	24.1	24.5	25.0	25.2	25.4	25.6	25.7
440	19.6	20.3	21.0	21.8	22.3	22.9	23.4	23.9	24.3	24.8	25.0	25.2	25.6
460	18.7	19.4	20.1	20.7	21.3	21.9	22.6	23.3	23.6	24.1	24.6	24.8	25.1
480	17.9	18.5	19.1	19.7	20.3	21.0	21.6	22.2	22.8	23.3	23.8	24.3	24.6
500	16.9	17.6	18.2	18.8	19.3	19.9	20.7	21.4	21.9	22.5	22.9	23.4	23.9
520	16.2	16.8	17.3	17.9	18.4	19.0	19.7	20.4	21.0	21.6	21.1	22.6	23.0
540	15.4	16.1	16.6	17.2	17.5	18.1	18.7	19.3	19.9	20.5	21.2	22.7	22.2
560	14.9	15.4	16.0	16.5	16.9	17.3	17.9	18.4	18.9	19.6	20.1	20.7	21.3
580	14.5	15.0	15.3	15.9	16.3	16.7	17.1	17.6	18.1	18.7	19.3	19.8	20.3
600	14.2	14.6	14.9	15.3	15.8	16.3	16.6	17.0	17.4	17.9	18.3	18.9	19.4
620	13.8	14.2	14.6	14.9	15.1	15.7	16.2	16.6	16.9	17.3	17.6	18.0	18.5
640	13.5	14.0	14.2	14.6	14.8	15.1	15.6	16.1	16.5	16.8	17.1	17.5	17.9
660	13.2	13.5	13.9	14.3	14.6	14.9	15.2	15.6	15.9	16.4	16.6	17.0	17.3
680	12.8	13.2	13.5	13.9	14.2	14.5	14.9	15.2	15.6	16.0	16.2	16.5	16.8
700	12.4	12.9	13.3	13.5	13.8	14.2	14.5	14.9	15.1	15.6	15.9	16.2	16.4
720	12.0	12.4	12.8	13.2	13.5	13.8	14.2	14.5	14.8	15.1	15.5	15.8	16.1
740	11.5	11.9	12.2	12.6	12.9	13.3	13.8	14.2	14.5	14.8	15.1	15.4	15.7
760	10.9	11.4	11.8	12.2	12.4	12.8	13.2	13.7	14.1	14.5	14.7	15.0	15.4
780	10.2	10.6	11.2	11.6	11.9	12.4	12.8	13.2	13.5	13.9	14.3	14.6	14.9
800	9.5	10.0	10.3	10.9	11.3	11.6	12.1	12.6	12.9	13.4	13.8	14.2	14.5
820	8.7	9.3	9.7	10.0	10.5	10.9	11.4	11.9	12.3	12.8	13.2	13.6	14.0
840	8.1	8.4	8.8	9.3	9.6	10.1	10.6	11.1	11.6	12.1	12.5	13.0	13.4
860	7.6	7.9	8.1	8.5	8.8	9.2	9.7	10.2	10.7	11.2	11.7	12.1	12.6
880	7.2	7.4	7.6	7.8	8.1	8.5	8.8	9.4	9.8	10.2	10.7	11.2	11.8
900	6.8	7.0	7.1	7.3	7.4	7.8	8.2	8.5	8.9	9.4	9.8	10.3	10.8
920	6.7	6.8	6.8	6.9	7.0	7.0	7.4	7.8	8.1	8.6	8.9	9.4	9.9
940	6.7	6.7	6.7	6.8	6.7	6.8	6.8	7.1	7.4	7.7	8.1	8.4	8.9
960	7.1	7.0	6.8	6.7	6.5	6.5	6.6	6.7	6.8	7.1	7.3	7.7	8.0
980	7.7	7.4	7.1	6.9	6.6	6.5	6.4	6.4	6.3	6.5	6.8	6.9	7.3
1000	8.4	8.0	7.7	7.3	6.9	6.7	6.5	6.5	6.3	6.2	6.2	6.4	6.5
	630	640	650	660	670	680	690	700	710	720	730	740	750

*Perturbations produced by Jupiter.*

Arguments II. and V.

V.

II.	750	760	770	780	790	800	810	820	830	840	850	860	870
0	"	"	"	"	"	"	"	"	"	"	"	"	"
20	6.5	6.8	7.2	7.5	8.0	8.4	8.8	9.5	10.1	10.5	11.0	11.6	12.4
40	6.1	6.2	6.5	6.7	7.0	7.4	7.9	8.4	9.0	9.5	10.0	10.6	11.1
60	5.8	5.9	5.9	6.2	6.4	6.6	6.9	7.4	7.8	8.2	8.8	9.5	10.0
80	5.8	5.7	5.7	5.7	5.9	6.1	6.2	6.5	6.9	7.2	7.7	8.3	8.8
100	6.2	5.8	5.7	5.6	5.4	5.6	5.7	5.9	6.1	6.3	6.7	7.3	7.8
120	6.8	6.3	5.9	5.6	5.5	5.3	5.3	5.4	5.4	5.6	5.9	6.3	6.8
140	7.6	7.4	6.5	6.0	5.7	5.5	5.1	5.2	5.1	5.1	5.2	5.5	5.8
160	8.8	8.1	7.4	6.8	6.2	5.8	5.4	5.2	5.0	4.9	4.8	5.0	5.1
180	10.0	9.3	8.5	7.8	7.2	6.5	5.9	5.5	5.1	5.9	4.7	4.7	4.7
200	11.5	10.6	9.7	9.0	8.2	7.5	6.9	6.3	5.8	5.2	4.8	4.7	4.5
220	13.1	12.2	11.2	10.4	9.5	8.8	7.9	7.1	6.5	5.9	5.3	5.0	4.7
240	14.7	13.8	12.9	12.0	11.1	10.2	9.3	8.4	7.5	6.7	6.1	5.5	5.2
260	16.4	15.3	14.5	13.6	12.6	11.7	10.7	9.8	8.8	7.9	7.0	6.5	5.9
280	18.1	17.2	16.3	15.3	14.3	13.3	12.2	11.4	10.4	9.4	8.3	7.7	6.9
300	19.8	18.9	17.9	17.0	16.1	15.0	14.0	13.0	11.9	10.9	9.9	8.9	8.0
320	21.2	20.4	19.6	18.7	17.7	16.8	15.8	14.7	13.7	12.6	11.5	10.5	9.4
340	22.5	21.9	21.2	20.4	19.4	18.5	17.4	16.5	15.5	14.2	13.2	12.3	11.2
360	23.7	23.0	22.4	21.8	21.1	20.2	19.2	18.3	17.1	16.1	15.0	13.9	12.9
380	24.5	24.0	23.6	23.0	22.4	21.6	20.8	19.9	18.9	17.9	16.8	15.9	14.7
400	25.2	24.9	24.5	24.0	23.5	22.8	22.1	21.4	20.5	19.5	18.5	17.6	16.5
420	25.5	25.4	25.1	24.8	24.5	23.9	23.4	22.7	21.9	21.0	20.1	19.2	18.2
440	25.7	25.6	25.5	25.3	25.0	24.5	24.2	23.7	23.2	22.3	21.5	20.7	19.8
460	25.6	25.6	25.7	25.7	25.5	25.3	24.9	24.6	24.1	23.4	22.7	22.0	21.2
480	25.1	25.3	25.5	25.6	25.8	25.7	25.4	25.2	24.8	24.3	23.7	23.1	22.5
500	24.6	24.9	25.2	25.4	25.6	25.6	25.5	25.4	25.2	24.9	24.5	24.1	23.5
520	23.9	24.2	24.7	25.0	25.3	25.4	25.5	25.5	25.4	25.2	24.9	24.7	24.3
540	23.0	23.6	23.9	24.3	24.7	24.9	25.2	25.4	25.4	25.3	25.2	25.1	24.8
560	22.2	22.6	23.2	23.6	24.0	24.4	24.6	24.9	25.1	25.0	25.1	25.1	25.0
580	21.3	21.7	22.2	22.8	23.2	23.7	24.0	24.3	24.6	24.7	24.8	24.9	24.9
600	20.3	20.8	21.3	21.8	22.3	22.7	23.2	23.7	23.9	24.1	24.4	24.6	24.7
620	19.4	19.9	20.4	20.8	21.4	21.9	22.2	22.7	23.1	23.4	23.7	24.1	24.3
640	18.5	19.0	19.5	20.1	20.5	20.9	21.4	21.8	22.2	22.6	22.9	23.3	23.6
660	17.9	18.3	18.7	19.2	19.7	20.1	20.5	22.0	21.3	21.7	22.1	22.5	22.8
680	17.3	17.6	18.1	18.5	18.9	19.4	19.6	20.1	20.5	20.7	21.2	21.7	22.0
700	16.8	17.1	17.4	17.8	18.2	18.6	18.9	19.4	19.7	20.1	20.4	20.7	21.2
720	16.4	16.7	16.9	17.3	17.7	18.0	18.3	18.7	18.9	19.2	19.6	20.0	20.3
740	16.1	16.3	16.5	16.9	17.2	17.6	17.8	18.0	18.3	18.5	18.7	19.2	19.5
760	15.7	16.0	16.2	16.5	16.7	17.0	17.3	17.6	17.8	17.9	18.1	18.5	18.8
780	15.4	15.7	16.0	16.1	16.4	16.6	16.7	17.2	17.4	17.4	17.8	18.0	18.2
800	14.9	15.3	15.6	15.9	16.1	16.3	16.5	16.7	16.9	17.1	17.3	17.6	17.7
820	14.5	14.7	15.2	15.5	15.8	15.9	16.2	16.5	16.6	16.8	16.9	17.1	17.3
840	14.0	14.4	14.7	15.1	15.4	15.7	15.8	16.1	16.3	16.4	16.6	16.9	17.0
860	13.4	13.7	14.1	14.5	15.1	15.4	15.4	15.8	15.9	16.1	16.2	16.6	16.7
880	12.6	13.1	13.5	13.9	14.3	14.8	15.2	15.5	15.6	15.8	16.0	16.3	16.4
900	11.8	12.3	12.8	13.3	13.7	14.1	14.5	15.0	15.3	15.4	15.6	15.9	16.1
920	10.8	11.3	11.9	12.4	13.0	13.4	13.7	14.2	14.7	15.0	15.2	15.5	15.7
940	9.9	10.3	10.8	11.4	12.0	12.5	12.9	13.4	14.0	14.3	14.7	15.0	15.3
960	8.9	9.4	9.9	10.4	11.0	11.6	12.1	12.5	13.0	13.6	13.9	14.4	14.7
980	8.0	8.3	8.8	9.4	10.0	10.6	11.1	11.7	12.2	12.5	13.1	13.7	14.1
1000	7.3	7.6	7.9	8.4	8.9	9.5	9.9	10.5	11.1	11.6	12.1	12.8	13.3
	6.5	6.8	7.2	7.5	8.0	8.4	8.8	9.5	10.0	10.5	11.0	11.6	12.4
	750	760	770	780	790	800	810	820	830	840	850	860	870

*Perturbations produced by Jupiter.*

Arguments II. and V.

V.

II.	870	880	890	900	910	920	930	940	950	960	970	980	990	1000
0	12.4	12.9	13.2	13.6	13.9	14.2	14.4	14.8	15.0	15.1	15.1	15.2	15.2	15.3
20	11.1	11.7	12.2	12.7	13.2	13.6	13.8	14.1	14.4	14.7	14.8	15.0	14.9	14.9
40	10.0	10.5	11.1	11.7	12.3	12.6	13.0	13.4	13.7	14.1	14.3	14.6	14.7	14.7
60	8.8	9.4	9.9	10.6	11.2	11.8	12.1	12.6	12.9	13.3	13.6	13.9	14.2	14.4
80	7.8	8.3	8.7	9.3	10.0	10.5	11.1	11.6	12.1	12.5	12.8	13.2	13.5	13.8
100	6.8	7.2	7.6	8.1	8.6	9.4	9.9	10.5	10.9	11.4	12.0	12.4	12.8	13.2
120	5.8	6.1	6.6	7.1	7.6	8.1	8.7	9.4	9.9	10.4	10.8	11.4	11.8	12.3
140	5.1	5.3	5.6	6.0	6.5	7.0	7.5	8.2	8.7	9.3	9.7	10.3	10.8	11.3
160	4.7	4.8	4.8	5.2	5.6	5.9	6.3	6.8	7.4	8.0	8.6	9.2	9.7	10.2
180	4.5	4.5	4.4	4.5	4.8	5.1	5.4	5.8	6.2	6.9	7.4	8.0	8.4	9.1
200	4.7	4.5	4.2	4.2	4.2	4.4	4.6	5.0	5.3	5.7	6.3	6.9	7.4	7.8
220	5.2	4.7	4.3	4.2	4.1	4.1	4.0	4.3	4.5	4.8	5.1	5.7	6.2	6.8
240	5.9	5.3	4.7	4.3	4.1	4.0	3.8	3.9	4.0	4.2	4.3	4.7	5.2	5.7
260	6.9	6.1	5.4	4.9	4.4	4.1	3.8	3.7	3.6	3.7	3.8	4.1	4.3	4.9
280	8.0	7.2	6.3	5.7	5.2	4.6	4.1	3.8	3.5	3.5	3.5	3.6	3.7	3.9
300	9.4	8.5	7.5	6.8	6.1	5.4	4.7	4.3	3.9	3.6	3.3	3.3	3.3	3.4
320	11.2	10.1	9.1	8.1	7.3	6.5	5.7	5.0	4.4	4.0	3.6	3.4	3.2	3.2
340	12.9	11.8	10.7	9.6	8.7	7.7	6.8	6.0	5.2	4.6	4.1	3.7	3.4	3.2
360	14.7	13.4	12.3	11.1	10.1	9.2	8.3	7.4	6.4	5.7	4.9	4.3	3.8	3.5
380	16.5	15.4	14.2	13.0	11.8	10.8	9.7	8.7	7.8	6.9	6.1	5.4	4.6	4.1
400	18.2	17.2	16.0	14.9	13.8	12.4	11.4	10.4	9.3	8.3	7.3	6.4	5.6	5.0
420	19.8	18.8	17.7	16.7	15.5	14.4	13.1	11.9	10.9	9.8	8.8	8.0	6.9	6.1
440	21.2	20.3	19.3	18.3	17.3	16.2	14.9	13.8	12.7	11.5	10.5	9.5	8.4	7.5
460	22.5	21.6	20.6	19.7	18.9	17.9	16.7	15.6	14.3	13.3	12.2	10.9	10.0	9.0
480	23.5	22.7	22.0	21.1	20.2	19.3	18.2	17.3	16.2	15.0	13.8	12.8	11.6	10.5
500	24.3	23.8	23.0	22.3	21.6	20.7	19.7	18.8	17.8	16.7	15.4	14.5	13.4	12.3
520	24.8	24.3	23.7	23.2	22.7	21.9	21.1	20.2	19.2	18.3	17.2	16.1	15.0	14.0
540	25.0	24.8	24.3	23.9	23.4	22.8	22.1	21.3	20.6	19.7	18.7	17.6	16.6	15.6
560	24.9	24.8	24.7	24.4	24.0	23.6	22.9	22.4	21.6	20.8	20.0	19.1	18.2	17.1
580	24.7	24.7	24.6	24.5	24.3	23.9	23.5	23.1	22.5	21.9	21.1	20.3	19.5	18.6
600	24.3	24.3	24.3	24.3	24.3	24.1	23.8	23.5	23.0	22.5	22.0	21.4	20.6	19.8
620	23.6	23.7	23.9	24.0	24.1	24.1	23.9	23.7	23.4	23.1	22.6	22.1	21.4	20.8
640	22.8	23.1	23.2	23.4	23.6	23.7	23.8	23.7	23.5	23.2	22.9	22.6	22.1	21.6
660	22.0	22.3	22.5	22.8	23.0	23.2	23.2	23.3	23.2	23.1	23.0	22.8	22.5	22.1
680	21.2	21.5	21.7	22.0	22.3	22.5	22.6	22.8	22.9	22.9	22.8	22.7	22.7	22.3
700	20.3	20.7	20.9	21.2	21.5	21.7	21.9	22.2	22.3	22.5	22.5	22.5	22.4	22.2
720	19.5	19.8	20.1	20.4	20.8	21.1	21.2	21.4	21.6	21.8	21.9	22.0	22.0	22.0
740	18.8	19.0	19.2	19.6	19.9	20.2	20.5	20.7	20.9	21.1	21.2	21.5	21.5	21.6
760	18.2	18.5	18.4	18.8	19.1	19.4	19.6	19.9	20.1	20.3	20.5	20.8	21.0	21.2
780	17.7	17.8	18.0	18.1	18.4	18.7	18.8	19.1	19.3	19.5	19.7	20.0	20.2	20.4
800	17.3	17.4	17.4	17.7	17.9	18.0	18.1	18.4	18.6	18.9	18.9	19.1	19.4	19.6
820	17.0	17.2	17.2	17.2	17.4	17.4	17.6	17.8	17.8	18.1	18.3	18.5	18.6	18.8
840	16.7	16.8	16.8	16.9	17.2	17.2	17.1	17.1	17.3	17.4	17.5	17.8	17.9	18.1
860	16.4	16.5	16.5	16.6	16.6	16.7	16.8	16.9	16.9	17.0	17.0	17.1	17.2	17.4
880	16.1	16.3	16.3	16.5	16.5	16.5	16.6	16.6	16.6	16.6	16.6	16.7	16.7	16.9
900	15.7	15.9	16.1	16.2	16.3	16.4	16.3	16.3	16.2	16.2	16.2	16.3	16.3	16.3
920	15.3	15.5	15.6	15.9	16.0	16.1	16.1	16.1	16.0	16.1	16.1	16.1	16.0	16.0
940	14.7	15.9	15.2	15.4	15.7	15.8	15.8	16.0	15.9	15.9	15.9	15.8	15.7	15.8
960	14.1	14.3	14.5	14.8	15.2	15.5	15.5	15.7	15.7	15.7	15.6	15.6	15.5	15.5
980	13.3	12.7	13.9	14.2	14.5	14.8	15.1	15.3	15.4	15.5	15.4	15.4	15.4	15.3
1000	12.4	12.9	13.2	13.6	13.9	14.2	14.4	14.8	15.0	15.1	15.1	15.2	15.2	15.3
	870	880	890	900	910	920	930	940	950	960	970	980	990	1000

*Perturbations produced by Saturn.*

Arguments II and VII.

VII.

II	0	100	200	300	400	500	600	700	800	900	1000
	"	"	"	"	"	"	"	"	"	"	"
0	1.2	1.5	1.4	1.0	0.7	0.6	0.5	0.5	0.4	0.8	1.2
100	0.9	1.2	1.3	1.1	0.9	0.8	0.7	0.7	0.6	0.7	0.9
200	0.7	0.9	1.0	1.1	1.0	0.9	0.8	0.8	0.9	0.8	0.7
300	0.9	0.8	0.7	0.8	0.9	1.0	1.0	1.0	1.0	1.0	0.9
400	1.0	0.9	0.6	0.4	0.6	0.9	1.0	1.1	1.1	1.1	1.0
500	1.1	1.0	0.8	0.4	0.2	0.5	1.0	1.3	1.3	1.2	1.1
600	1.2	1.1	0.9	0.6	0.2	0.2	0.5	1.1	1.5	1.5	1.2
700	1.4	1.1	1.0	0.8	0.4	0.1	0.3	0.8	1.4	1.7	1.4
800	1.6	1.3	1.0	0.8	0.6	0.4	0.1	0.3	1.0	1.6	1.6
900	1.5	1.4	1.1	0.9	0.7	0.6	0.3	0.2	0.6	1.2	1.5
1000	1.2	1.5	1.4	1.0	0.7	0.6	0.5	0.5	0.4	0.8	1.2

Constant, 1." 0

TABLE XXXIV.

*Variable Part of Sun's Aberration.*

Argument, Sun's Mean Anomaly.

	Os	Is	II <sup>s</sup>	III <sup>s</sup>	IV <sup>s</sup>	V <sup>s</sup>	
°	"	"	"	"	"	"	°
0	0.0	0.0	0.1	0.3	0.5	0.6	30
3	0.0	0.0	0.2	0.3	0.5	0.6	27
6	0.0	0.0	0.2	0.3	0.5	0.6	24
9	0.0	0.0	0.2	0.3	0.5	0.6	21
12	0.0	0.1	0.2	0.4	0.5	0.6	18
15	0.0	0.1	0.2	0.4	0.5	0.6	15
18	0.0	0.1	0.2	0.4	0.5	0.6	12
21	0.0	0.1	0.3	0.4	0.6	0.6	9
24	0.0	0.1	0.3	0.4	0.6	0.6	6
27	0.0	0.1	0.3	0.4	0.6	0.6	3
30	0.0	0.1	0.3	0.5	0.6	0.6	0
	XI <sup>s</sup>	X <sup>s</sup>	IX <sup>s</sup>	VIII <sup>s</sup>	VII <sup>s</sup>	VI <sup>s</sup>	

Constant, 0." 3



*Moon's Epochs.*

Years.	1	2	3	4	5	6	7	8	9	10	11	12	13
1830	00174	4541	4461	4638	9885	0635	5979	9921	7623	219	226	458	468
1831	00103	1749	4127	9381	2357	6432	7040	2378	6487	825	587	177	940
1832 B	00032	8957	3793	4125	4829	2229	8100	4835	5351	432	948	897	413
1833	00235	6816	4499	9156	7636	8399	9219	7683	4239	108	340	687	920
1834	00164	4024	4164	3900	0107	4196	0279	0140	3103	715	701	406	393
1835	00093	1232	3830	8644	2579	9993	1340	2598	1967	321	061	125	866
1836 B	00022	8441	3496	3388	5051	5791	2400	5055	0831	928	422	845	339
1837	00224	6299	4202	8419	7858	1960	3518	7903	9719	605	814	635	846
1838	00153	3508	3868	3163	0329	7757	4579	0360	8583	211	175	354	319
1839	00082	0716	3534	7907	2801	3555	5639	2818	7447	818	536	074	792
1840 B	00011	7925	3199	2651	5273	9352	6700	5275	6310	424	896	793	265
1841	00213	5783	3906	7682	8080	5522	7818	8123	5199	101	288	583	772
1842	00142	2991	3571	2425	0551	1319	8879	0580	4062	707	649	302	245
1843	00071	0200	3237	7169	3023	7116	9939	3038	2926	314	010	022	718
1844 B	00000	7408	2903	1913	5495	2914	1000	5495	1790	920	371	741	191
1845	00203	5266	3609	6944	8302	9083	2118	8343	0678	597	763	531	698
1846	00132	2475	3275	1688	0773	4880	3179	0800	9542	203	123	250	171
1847	00061	9683	2941	6432	3245	0678	4239	3257	8406	810	484	970	644
1848 B	99990	6892	2606	1176	5717	6475	5300	5715	7270	416	845	689	117
1849	00192	4750	3312	6207	8524	2644	6418	8563	6158	093	237	479	624
1850	00121	1958	2978	0951	0995	8442	7479	1020	5022	700	597	199	097
1851	00050	9167	2644	5695	3467	4239	8539	3477	3885	306	958	918	570
1852 B	99979	6375	2310	0439	5939	0036	9600	5935	2749	913	319	637	043
1853	00181	4233	3016	5469	8746	6206	0718	8782	1637	589	711	427	550
1854	00110	1442	2681	0213	1217	2003	1778	1240	0501	196	072	147	023
1855	00039	8650	2347	4957	3689	7801	2839	3697	9365	802	432	866	496
1856 B	99968	5859	2013	9701	6160	3598	3899	6155	8229	409	793	586	969
1857	00171	3717	2719	4732	8968	9767	5018	9002	7117	086	185	375	476
1858	00100	0925	2385	9476	1439	5565	6078	1460	5981	692	546	095	949
1859	00029	8134	2051	4220	3911	1362	7139	3917	4845	299	907	814	422
1860 B	99958	5342	1716	8964	6383	7159	8199	6374	3709	905	267	534	895
1861	00160	3200	2423	3995	9190	3329	9317	9222	2597	581	659	323	402
1862	00089	0409	2088	8739	1661	9126	0378	1679	1461	188	020	043	875
1863	00018	7617	1754	3483	4133	4923	1438	4137	0324	795	381	762	348
1864 B	99947	4826	1420	8227	6605	0721	2499	6594	9188	401	742	482	821
1865	00149	2684	2126	3257	9412	6890	3617	9442	8076	078	134	272	328
1866	00078	9893	1792	8001	1883	2687	4678	1899	6940	685	494	991	801
1867	00007	7101	1457	2745	4355	8485	5738	4357	5804	291	855	711	274
1868 B	99936	4309	1123	7489	6827	4282	6799	6814	4668	898	216	431	747
1869	00138	2168	1829	2520	9634	0452	7917	9662	3556	574	608	220	254
1870	00067	9376	1495	7264	2105	6249	8978	2119	2420	181	968	940	727



TABLE XXXV.

*Moon's Epochs.*

Years.	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31
1830	921	392	230	588	462	523	536	52	60	44	94	51	47	98	99	99	89	52
1831	115	532	589	940	937	296	703	30	70	41	65	53	94	48	24	24	51	44
1832 B	309	673	949	293	412	070	870	07	81	38	36	55	42	97	48	49	14	35
1833	602	844	345	688	913	845	037	85	92	45	07	61	92	53	77	77	77	27
1834	796	984	704	040	388	619	203	62	03	42	77	63	40	03	01	01	39	18
1835	989	124	063	393	863	392	370	39	13	38	48	65	87	51	26	26	02	10
1836 B	183	265	423	745	338	166	537	17	24	35	19	67	34	01	50	51	64	01
1837	476	436	819	140	840	942	704	94	35	42	90	73	85	58	79	79	27	93
1838	670	576	178	492	315	715	870	72	46	38	60	75	32	07	04	04	89	84
1839	864	716	537	845	790	489	037	49	56	35	31	77	80	56	28	28	52	76
1840 B	058	857	897	197	265	262	204	26	67	32	02	79	27	06	53	53	14	67
1841	351	028	293	592	766	038	371	04	78	39	73	85	77	62	81	81	77	59
1842	544	168	652	944	241	811	537	81	89	35	43	87	25	12	06	06	40	51
1843	738	308	012	297	716	585	704	58	99	32	14	89	72	61	30	31	02	42
1844 B	932	449	371	649	191	358	871	36	10	29	85	91	19	10	55	55	65	34
1845	225	620	767	044	692	134	038	13	21	36	56	97	70	67	84	83	27	26
1846	419	760	126	396	167	907	204	91	32	32	26	99	17	16	08	08	90	17
1847	613	901	486	749	643	681	371	68	42	29	97	01	65	65	33	33	52	09
1848 B	806	041	845	101	118	454	538	45	53	26	68	03	12	15	57	58	15	00
1849	099	212	241	496	619	230	705	23	64	33	39	09	63	71	86	86	77	92
1850	293	352	600	848	094	003	871	00	75	29	09	10	10	20	10	10	40	83
1851	487	493	960	201	569	777	038	78	85	26	80	12	57	70	35	35	02	75
1852 B	681	633	319	553	044	550	205	55	96	23	51	14	04	19	59	60	65	66
1853	974	804	715	948	545	326	372	33	07	30	22	20	55	76	88	88	28	58
1854	168	944	074	300	020	099	539	10	18	26	93	22	03	25	12	12	90	50
1855	361	085	434	653	495	873	705	87	28	23	63	24	50	74	37	37	53	41
1856 B	555	225	793	005	970	646	872	65	39	20	34	26	97	23	61	62	15	33
1857	848	390	189	400	471	422	039	42	50	27	05	32	48	80	90	90	78	24
1858	042	537	548	752	947	195	206	20	61	24	76	34	95	29	15	15	40	16
1859	236	677	908	105	422	969	372	97	71	20	46	36	42	79	39	40	03	07
1860 B	430	817	267	457	897	742	539	74	82	17	17	38	89	28	64	64	65	99
1861	723	988	663	852	398	518	706	52	93	24	88	44	41	84	92	92	28	91
1862	916	129	022	204	873	291	873	29	04	20	60	46	88	34	17	17	91	82
1863	110	269	382	557	348	065	039	06	14	17	29	48	35	82	41	42	53	74
1864 B	304	409	741	909	823	838	206	84	25	14	00	50	82	32	66	66	16	65
1865	597	580	137	304	324	614	373	61	36	21	71	56	33	89	95	94	78	57
1866	791	721	496	657	799	387	540	39	47	17	42	58	80	38	19	19	41	49
1867	985	861	856	009	274	161	707	16	57	14	12	60	28	87	44	44	03	40
1868 B	178	001	215	362	749	934	873	93	68	11	83	62	75	37	68	69	66	32
1869	471	172	611	756	251	710	040	71	79	18	54	68	26	93	97	97	28	23
1870	665	313	970	109	726	483	207	48	90	15	26	69	73	43	21	21	91	15

*Moon's Epochs.*

Years.	Evection.				Anomaly.				Variation.				Longitude.			
	s	o	'	"	s	o	'	"	s	o	'	"	s	o	'	"
1830	5	17	4	12	11	24	31	4.5	2	13	2	39	11	22	55	37.7
1831	11	7	35	41	2	23	14	24.6	6	22	40	4	4	2	18	42.8
1832 B	4	28	7	11	5	21	57	44.4	11	2	17	28	8	11	41	48.0
1833	10	29	57	40	9	3	44	58.5	3	24	6	21	1	4	15	28.4
1834	4	20	29	11	0	2	28	18.5	8	3	43	45	5	13	38	33.6
1835	10	11	0	40	3	1	11	38.6	0	13	21	10	9	23	1	38.8
1836 B	4	1	32	9	5	29	54	58.7	4	22	58	34	2	2	24	44.0
1837	10	3	22	39	9	11	42	12.8	9	14	47	27	6	24	58	24.5
1838	3	23	54	9	0	10	25	32.9	1	24	24	51	11	4	21	29.8
1839	9	14	25	38	3	9	8	53.1	6	4	2	16	3	13	44	35.0
1840 B	3	4	57	8	6	7	52	13.2	10	13	39	42	7	23	7	40.4
1841	9	6	47	37	9	19	39	27.5	3	5	28	33	0	15	41	20.9
1842	2	27	19	7	0	18	22	47.6	7	15	5	58	4	25	4	26.2
1843	8	17	50	37	3	17	6	7.9	11	24	43	23	9	4	27	31.6
1844 B	2	8	22	7	6	15	49	28.1	4	4	20	48	1	13	50	37.0
1845	8	10	12	36	9	27	36	42.5	8	26	9	40	6	6	24	17.5
1846	2	0	44	6	0	26	20	2.8	1	5	47	5	10	15	47	23.0
1847	7	21	15	35	3	25	3	23.2	5	15	24	30	2	25	10	28.3
1848 B	1	11	47	5	6	23	46	43.5	9	25	1	55	7	4	33	33.7
1849	7	13	37	35	10	5	33	57.9	2	16	50	47	11	27	7	14.5
1850	1	4	9	4	1	4	17	18.3	6	26	28	12	4	6	30	19.9
1851	6	24	40	35	4	3	0	38.6	11	6	5	37	8	15	53	25.4
1852 B	0	15	12	5	7	1	43	59.2	3	15	43	3	0	25	16	31.0
1853	6	17	2	34	10	13	31	13.7	8	7	31	54	5	17	50	11.6
1854	0	7	34	4	1	12	14	34.1	0	17	9	20	9	27	13	17.2
1855	5	28	5	33	4	10	57	54.7	4	26	46	44	2	6	36	22.7
1856 B	11	18	37	3	7	9	41	15.2	9	6	24	10	6	15	59	28.2
1857	5	20	27	33	10	21	28	29.8	1	28	13	2	11	8	33	9.1
1858	11	10	59	2	1	20	11	50.3	6	7	50	27	3	17	56	14.6
1859	5	1	30	33	4	18	55	10.9	10	17	27	53	7	27	19	20.1
1860 B	10	22	2	3	7	17	38	31.4	2	27	5	18	0	6	42	25.8
1861	4	23	52	32	10	29	25	46.1	7	18	54	10	4	29	16	6.6
1862	10	14	24	2	1	28	9	6.6	11	28	31	35	9	8	39	12.2
1863	4	4	55	32	4	26	52	27.3	4	8	9	1	1	18	2	17.9
1864 B	9	25	27	2	7	25	35	48.0	8	17	46	25	5	27	25	23.5
1865	3	27	17	31	11	7	23	2.7	1	9	35	18	10	19	59	4.3
1866	9	17	49	2	2	6	6	23.3	5	19	12	43	2	29	22	10.1
1867	3	8	20	31	5	4	49	44.0	9	28	50	9	7	8	45	15.7
1868 B	8	28	52	2	8	3	33	4.7	2	8	27	34	11	18	8	21.4
1869	3	0	42	33	11	15	20	19.6	7	0	16	26	4	10	42	2.3
1870	8	21	14	2	2	14	3	40.3	11	9	53	51	8	20	5	8.0

*Moon's Epochs.*

Years.	Supp. of Node.	II	V	VI	VII	VIII	IX	X	XI	XII
	<i>s</i> <i>o</i> ' "	<i>s</i> <i>o</i> "								
1830	6 7 7 11.0	10 24 46	498	502	900	904	427	062	025	433
1831	6 26 26 53.3	2 15 18	912	914	208	210	506	001	211	710
1832 B	7 15 46 35.5	6 5 50	326	327	516	516	586	940	397	986
1833	8 5 9 28.4	10 7 31	774	779	852	856	702	885	624	297
1834	8 24 29 10.7	1 28 3	187	191	159	163	782	825	810	573
1835	9 13 48 53.0	5 18 35	601	603	467	469	861	764	996	850
1836 B	10 3 8 35.2	9 9 8	015	016	775	775	941	703	182	127
1837	10 22 31 28.1	1 10 49	463	468	111	116	057	648	409	437
1838	11 11 51 10.4	5 1 21	876	880	419	423	137	588	595	714
1839	0 1 10 52.6	8 21 53	290	292	726	729	217	527	781	991
1840 B	0 20 30 34.9	0 12 25	704	705	034	035	296	466	967	268
1841	1 9 53 27.7	4 14 6	152	157	370	375	412	411	194	578
1842	1 29 13 10.0	8 4 38	566	569	678	682	492	350	380	855
1843	2 18 32 52.2	11 25 10	980	980	986	988	572	290	566	131
1844 B	3 7 52 34.5	3 15 42	393	394	293	294	651	229	752	408
1845	3 27 15 27.4	7 17 23	840	846	629	634	767	174	979	718
1846	4 16 35 9.6	11 7 55	254	258	937	941	847	113	165	995
1847	5 5 54 51.8	2 28 27	668	670	245	247	927	053	351	272
1848 B	5 25 14 34.1	6 18 59	082	083	553	553	006	992	537	549
1849	6 14 37 27.0	10 20 40	531	535	889	893	122	937	764	859
1850	7 3 57 9.2	2 11 12	944	947	196	200	202	876	950	136
1851	7 23 16 51.5	6 1 44	358	359	504	506	282	816	136	413
1852 B	8 12 36 33.6	9 22 17	772	772	812	812	362	755	322	689
1853	9 1 59 26.5	1 23 58	220	223	148	152	477	700	549	000
1854	9 21 19 8.8	5 14 30	634	636	456	459	557	639	735	276
1855	10 10 38 51.1	9 5 2	047	048	763	765	637	579	921	553
1856 B	10 29 58 33.3	0 25 34	461	461	71	071	717	518	107	830
1857	11 19 21 26.2	4 27 15	909	912	407	411	832	463	334	140
1858	0 8 41 8.4	8 17 47	323	325	715	718	912	402	520	417
1859	0 28 0 50.7	0 8 19	736	737	023	024	992	342	706	694
1860 B	1 17 20 32.9	3 28 51	150	150	330	330	072	281	892	971
1861	2 6 43 25.8	8 0 32	598	601	666	670	187	226	119	281
1862	2 26 3 8.0	11 21 4	012	014	974	977	267	165	305	558
1863	3 15 22 50.1	3 11 36	426	426	282	283	347	105	491	834
1864 B	4 4 42 32.3	7 2 8	839	839	590	589	427	044	677	111
1865	4 24 5 25.2	11 3 49	287	291	926	929	542	989	904	422
1866	5 13 25 7.3	2 24 21	701	703	233	236	622	928	090	698
1867	6 2 44 49.5	6 14 53	115	115	541	542	702	868	276	975
1868 B	6 22 4 31.7	10 5 26	529	528	849	848	782	807	462	252
1869	7 11 27 24.6	2 7 7	977	980	185	188	897	752	689	562
1870	8 0 47 6.7	5 27 39	390	392	493	495	977	691	875	839

*Moon's Motions for Months.*

Months.	1	2	3	4	5	6	7	8	9	10	11	12	13
January	00000	0000	0000	0000	0000	0000	0000	0000	0000	000	000	000	000
February	08487	0146	2246	8896	0402	1533	1789	2099	0753	175	965	184	059
March	Com.	16153	8343	1371	6931	9797	1951	3404	3027	1433	139	836	157 016
	Bis.	16427	8993	2411	7218	0132	2323	3462	3418	1457	209	868	228 050
April	Com.	24640	8490	3616	5827	0199	3484	5193	5126	2186	314	801	342 076
	Bis.	24914	9140	4657	6114	0534	3856	5251	5517	2210	384	832	412 110
May	Com.	32853	7986	4822	4436	0265	4646	6924	6835	2914	419	735	456 101
	Bis.	33127	8636	5862	4723	0600	5018	6982	7226	2938	489	766	526 135
June	Com.	41340	8133	7067	3332	0666	6179	8713	8934	3667	593	700	640 160
	Bis.	41614	8783	8107	3619	1002	6551	8771	9325	3691	663	731	710 194
July	Com.	49554	7629	8273	1942	0732	7341	0444	0643	4396	698	634	754 185
	Bis.	49828	8279	9313	2228	1068	7713	0502	1034	4420	768	665	824 219
Aug.	Com.	58041	7776	0518	0838	1134	8874	2233	2742	5148	873	599	938 245
	Bis.	58315	8426	1558	1125	1470	9246	2290	3133	5173	943	630	009 279
Sept.	Com.	66528	7922	2764	9734	1536	0408	4021	4842	5901	048	563	123 304
	Bis.	66802	8572	3804	0021	1871	0780	4079	5232	5925	118	595	193 338
Oct.	Com.	74741	7419	3969	8343	1602	1569	5752	6550	6630	152	497	237 329
	Bis.	75015	8069	5009	8630	1938	1941	5810	6941	6654	222	528	307 363
Nov.	Com.	83228	7565	6215	7239	2004	3102	7541	8649	7382	327	462	421 383
	Bis.	83502	8215	7255	7526	2339	3475	7599	9040	7407	397	493	492 423
Dec.	Com.	91442	7062	7420	5848	2070	4264	9272	0358	8111	432	396	535 414
	Bis.	91716	7712	8460	6135	2405	4636	9330	0749	8135	502	427	606 448

## TABLE XXXVI.

*Moon's Motions for Months.*

Months.	Evection.				Anomaly.				Variation.				Longitude.			
	s	o	'	"	s	o	'	"	s	o	'	"	s	o	'	"
January	0	0	0	0	0	0	0	0.0	0	0	0	0	0	0	0	0.0
February	11	20	48	42	1	15	0	53.1	0	17	54	48	1	18	28	5.8
March	Com.	10	7	40	26	1	20	50	4.2	11	29	15	1	27	24	26.6
	Bis.	10	18	59	26	2	3	53	58.2	0	11	26	42	2	10	35 1.6
April	Com.	9	28	29	8	3	5	50	57.3	0	17	10	3	3	15	52 32.5
	Bis.	10	9	48	8	3	18	54	51.2	0	29	21	29	3	29	3 7.5
May	Com.	9	7	58	51	4	7	47	56.4	0	22	53	24	4	21	10 3.3
	Bis.	9	19	17	50	4	20	51	50.3	1	5	4	50	5	4	20 38.3
June	Com.	8	28	47	33	5	22	48	49.4	1	10	48	11	6	9	38 9.1
	Bis.	9	10	6	33	6	5	52	43.4	1	22	59	38	6	22	48 44.1
July	Com.	8	8	17	16	6	24	45	48.5	1	16	31	32	7	14	55 39.9
	Bis.	8	19	36	15	7	7	49	42.5	1	28	42	59	7	28	6 15.0
Aug.	Com.	7	29	5	59	8	9	46	41.6	2	4	26	20	9	3	23 45.8
	Bis.	8	10	24	58	8	22	50	35.5	2	16	37	47	9	16	34 20.8
Sept.	Com.	7	19	54	41	9	24	47	34.6	2	22	21	7	10	21	51 51.6
	Bis.	8	1	13	40	10	7	51	28.6	3	4	32	34	11	5	2 26.7
Oct.	Com.	6	29	24	24	10	26	44	33.7	2	28	4	28	11	27	9 22.4
	Bis.	7	10	43	23	11	9	48	27.7	3	10	15	55	0	10	19 57.5
Nov.	Com.	6	20	13	6	0	11	45	26.8	3	15	59	16	1	15	37 28.3
	Bis.	7	1	32	5	0	24	49	20.7	3	28	10	43	1	28	48 3.3
Dec.	Com.	5	29	42	49	1	13	42	25.9	3	21	42	37	2	20	54 59.1
	Bis.	6	11	1	48	1	26	46	19.8	4	3	54	4	3	4	5 34.1



*Moon's Motions for Months.*

Months.	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31
January	000	000	000	000	000	000	000	00	00	00	00	00	00	00	00	00	00	00
February	074	946	135	304	805	066	014	24	26	14	82	28	14	17	29	96	05	07
March { Com.	851	801	159	482	532	125	027	45	50	98	57	43	18	12	46	82	10	15
March { Bis.	950	831	196	524	558	127	027	46	51	08	59	47	21	19	51	85	10	15
April { Com.	925	747	294	786	336	191	041	68	77	12	39	70	32	29	76	77	15	23
April { Bis.	024	778	331	828	362	193	042	69	77	22	42	74	36	36	80	80	16	23
May { Com.	899	663	392	047	115	254	055	91	02	15	19	94	43	38	01	70	21	30
May { Bis.	999	693	429	089	141	256	055	92	03	26	22	98	47	45	05	73	21	30
June { Com.	973	609	527	351	920	320	069	15	28	29	01	21	57	55	31	65	26	38
June { Bis.	073	639	563	393	946	322	069	15	29	40	04	25	61	62	35	68	26	38
July { Com.	948	525	625	613	699	384	083	37	54	33	81	45	68	64	56	58	31	45
July { Bis.	047	555	661	655	725	386	083	38	55	43	84	49	72	71	60	61	31	46
Aug. { Com.	022	471	759	917	503	449	097	61	80	47	64	72	82	81	85	53	36	53
Aug. { Bis.	121	501	796	959	529	451	097	62	81	57	66	77	86	88	90	56	36	53
Sept. { Com.	096	417	894	221	308	515	111	85	07	61	46	00	97	97	15	49	42	61
Sept. { Bis.	195	447	931	263	334	517	111	85	08	71	49	04	01	04	19	52	42	61
Oct. { Com.	071	333	992	483	087	578	125	07	32	65	26	23	08	07	40	41	47	68
Oct. { Bis.	170	363	029	525	113	581	126	08	33	75	28	28	11	14	44	44	47	69
Nov. { Com.	145	279	127	787	892	644	139	31	59	79	08	51	22	23	70	37	52	76
Nov. { Bis.	244	309	163	829	918	646	140	32	60	89	11	55	26	30	74	40	52	76
Dec. { Com.	120	194	225	049	670	708	153	54	85	83	88	74	33	33	95	29	57	84
Dec. { Bis.	219	225	261	091	696	710	153	54	86	93	90	79	37	40	99	32	57	84

TABLE XXXVI.

*Moon's Motions for Months.*

Months.	Supp. of Node.				II		V	VI	VII	VIII	IX	X	XI	XII
	<i>s</i>	<i>o</i>	<i>'</i>	<i>''</i>	<i>s</i>	<i>o</i>	<i>'</i>							
January	0	0	0	0.0	0	0	0	000	000	000	000	000	000	000
February	0	1	38	29.7	11	15	43	054	224	875	045	111	165	290
March { Com.	0	3	7	27.5	9	27	59	007	330	666	989	114	313	455
March { Bis.	0	3	10	38.2	10	9	8	041	369	694	023	150	319	496
April { Com.	0	4	45	57.3	9	13	42	061	554	542	034	225	478	745
April { Bis.	0	4	49	7.9	9	24	51	095	593	570	068	261	484	787
May { Com.	0	6	21	16.4	8	18	15	081	738	389	046	300	638	993
May { Bis.	0	6	24	27.0	8	29	25	115	778	417	080	336	643	034
June { Com.	0	7	59	46.1	8	3	58	136	962	264	091	411	802	282
June { Bis.	0	8	2	56.7	8	15	8	170	002	293	124	447	808	324
July { Com.	0	9	35	5.2	7	8	32	156	147	112	103	486	962	531
July { Bis.	0	9	38	15.9	7	19	41	190	186	140	136	522	967	572
Aug. { Com.	0	11	13	35.0	6	24	15	210	371	987	147	597	126	820
Aug. { Bis.	0	11	16	45.6	7	5	24	244	411	015	182	633	132	862
Sept. { Com.	0	12	52	4.7	6	9	58	265	595	862	193	708	291	110
Sept. { Bis.	0	12	55	15.4	6	21	7	299	635	891	227	744	296	152
Oct. { Com.	0	14	27	23.8	5	14	32	285	780	710	204	783	451	358
Oct. { Bis.	0	14	30	34.4	5	25	41	319	819	738	238	819	456	400
Nov. { Com.	0	16	5	53.5	5	0	15	339	004	585	250	894	615	648
Nov. { Bis.	0	16	9	4.2	5	11	24	373	043	613	283	930	621	690
Dec. { Com.	0	17	41	12.6	4	4	49	359	188	432	261	969	775	896
Dec. { Bis.	0	17	44	23.3	4	15	58	393	228	461	295	005	780	938



*Moon's Motions for Days.*

D.	1	2	3	4	5	6	7	8	9	10	11	12	13
1	00000	0000	0000	0000	0000	0000	0000	0000	0000	000	000	000	000
2	00274	0650	1040	0287	0336	0372	0058	0390	0024	070	031	070	034
3	00548	1300	2080	0574	0671	0744	0115	0781	0049	140	062	141	068
4	00821	1950	3121	0861	1007	1116	0173	1171	0073	210	093	211	103
5	01095	2600	4161	1148	1342	1488	0231	1561	0097	281	125	282	137
6	01369	3249	5201	1435	1678	1860	0289	1952	0121	351	156	352	171
7	01643	3899	6241	1722	2013	2232	0346	2342	0146	421	187	423	205
8	01916	4549	7281	2009	2349	2604	0404	2732	0170	491	218	493	239
9	02190	5199	8321	2296	2684	2976	0462	3122	0194	561	249	564	273
10	02464	5849	9362	2583	3020	3348	0519	3513	0219	631	280	634	308
11	02738	6499	0402	2870	3355	3720	0577	3903	0243	702	311	705	342
12	03012	7149	1442	3157	3691	4093	0635	4293	0267	772	342	775	376
13	03285	7799	2482	3444	4026	4465	0692	4684	0291	842	374	845	410
14	03559	8449	3522	3731	4362	4837	0750	5074	0316	912	405	916	444
15	03833	9098	4563	4018	4698	5209	0808	5464	0340	982	436	986	478
16	04107	9748	5603	4305	5033	5581	0866	5854	0364	052	467	057	513
17	04380	0398	6643	4592	5369	5953	0923	6245	0389	122	498	127	547
18	04654	1048	7683	4878	5704	6325	0981	6635	0413	193	529	198	581
19	04928	1698	8723	5165	6040	6697	1039	7025	0437	263	560	268	615
20	05202	2348	9763	5452	6375	7069	1096	7416	0461	333	591	339	649
21	05476	2998	0804	5739	6711	7441	1154	7806	0486	403	623	409	683
22	05749	3648	1844	6026	7046	7813	1212	8196	0510	473	654	480	718
23	06023	4298	2884	6313	7382	8185	1269	8586	0534	543	685	550	752
24	06297	4947	3924	6600	7717	8557	1327	8977	0559	614	716	621	786
25	06571	5597	4964	6887	8053	8929	1385	9367	0583	684	747	691	820
26	06844	6247	6005	7174	8389	9301	1443	9757	0607	754	778	762	854
27	07118	6897	7045	7461	8724	9673	1500	0148	0631	824	809	832	888
28	07392	7547	8085	7748	9060	0045	1558	0538	0656	894	840	903	923
29	07666	8197	9125	8035	9395	0417	1616	0928	0680	964	872	973	957
30	07940	8847	0165	8322	9731	0789	1673	1319	0704	034	903	043	991
31	08213	9497	1205	8609	0066	1161	1731	1709	0729	105	934	114	025

*Moon's Motion for Days.*

D	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31
1	000	000	000	000	000	000	000	00	00	00	00	00	00	00	00	00	00	00
2	099	031	037	042	026	002	000	01	01	10	03	04	04	07	04	03	00	00
3	198	061	073	084	052	004	001	02	02	20	05	08	07	14	08	06	00	00
4	297	092	110	126	078	006	001	02	03	30	08	12	11	21	13	09	01	01
5	397	122	146	168	104	008	002	03	03	41	11	16	15	28	17	12	01	01
6	496	153	183	210	130	011	002	04	04	51	13	21	18	35	21	15	01	01
7	595	183	220	252	156	013	003	05	05	61	16	25	22	42	25	18	01	01
8	694	214	256	294	182	015	003	05	06	71	19	29	26	49	29	22	01	02
9	793	244	293	336	208	017	004	06	07	81	21	33	30	56	33	25	01	02
10	892	275	329	379	234	019	004	07	08	91	24	37	33	63	38	28	02	02
11	992	305	366	421	260	021	005	08	09	01	27	41	37	70	42	31	02	02
12	091	336	403	463	286	023	005	08	09	11	29	45	41	77	46	34	02	03
13	190	366	439	505	312	025	005	09	10	22	32	49	44	84	50	37	02	03
14	289	397	476	547	337	028	006	10	11	32	34	53	48	91	54	40	02	03
15	388	427	512	589	363	030	006	11	12	42	37	58	52	98	58	43	02	03
16	487	458	549	631	389	032	007	11	13	52	40	62	55	05	63	46	03	04
17	587	488	586	673	415	034	007	12	14	62	42	66	59	12	67	49	03	04
18	686	519	622	715	441	036	008	13	14	72	45	70	63	19	71	52	03	04
19	785	549	659	757	467	038	008	14	15	82	48	74	66	26	75	55	03	04
20	884	580	695	799	493	040	009	14	16	92	50	78	70	33	79	59	03	05
21	983	611	732	841	519	042	009	15	17	03	53	82	74	40	84	62	03	05
22	082	641	769	883	545	044	010	16	18	13	56	86	77	47	88	65	04	05
23	182	672	805	925	571	047	010	17	19	23	58	90	81	54	92	68	04	05
24	281	702	842	967	597	049	011	17	20	33	61	95	85	61	96	71	04	06
25	380	733	878	009	623	051	011	18	20	43	64	99	89	68	00	74	04	06
26	479	763	915	052	649	053	011	19	21	53	66	03	92	75	04	77	04	06
27	578	794	952	094	675	055	012	20	22	63	69	07	96	82	09	80	04	06
28	677	824	988	136	701	057	012	20	23	73	72	11	00	89	13	83	05	06
29	777	855	025	178	727	059	013	21	24	84	74	15	03	96	17	86	05	07
30	876	885	061	220	753	061	013	22	25	94	77	19	07	03	21	89	05	07
31	975	916	098	262	779	064	014	23	26	04	80	23	11	10	25	92	05	07

*Moon's Motions for Days.*

D.	Evection.	Anomaly.	Variation.	M. Longitude.
	<i>s</i> <i>o</i> <i>'</i>	<i>s</i> <i>o</i> <i>'</i> <i>''</i>	<i>s</i> <i>o</i> <i>'</i> <i>''</i>	<i>s</i> <i>o</i> <i>'</i> <i>''</i>
1	0 0 0 0	0 0 0 0.0	0 0 0 0	0 0 0 0.0
2	0 11 18 59	0 13 3 54.0	0 12 11 27	0 13 10 35.0
3	0 22 37 59	0 26 7 47.9	0 24 22 53	0 26 21 10.1
4	1 3 56 58	1 9 11 41.9	1 6 34 20	1 9 31 45.1
5	1 15 15 58	1 22 15 35.9	1 18 45 47	1 22 42 20.1
6	1 26 34 57	2 5 19 29.8	2 0 57 13	2 5 52 55.1
7	2 7 53 57	2 18 23 23.8	2 13 8 40	2 19 3 30.2
8	2 19 12 56	3 1 27 17.8	2 25 20 7	3 2 14 5.2
9	3 0 31 55	3 14 31 11.7	3 7 31 34	3 15 24 40.2
10	3 11 50 55	3 27 35 5.7	3 19 43 0	3 28 35 15.2
11	3 23 9 54	4 10 38 59.7	4 1 54 27	4 11 45 50.3
12	4 4 28 54	4 23 42 53.7	4 14 5 54	4 24 56 25.3
13	4 15 47 53	5 6 46 47.6	4 26 17 20	5 8 7 0.3
14	4 27 6 53	5 19 50 41.6	5 8 28 47	5 21 17 35.4
15	5 8 25 52	6 2 54 35.6	5 20 40 14	6 4 28 10.4
16	5 19 44 51	6 15 58 29.5	6 2 51 40	6 17 38 45.4
17	6 1 3 51	6 29 2 23.5	6 15 3 7	7 0 49 20.4
18	6 12 22 50	7 12 6 17.5	6 27 14 34	7 13 59 55.5
19	6 23 41 50	7 25 10 11.4	7 9 26 1	7 27 10 30.5
20	7 5 0 49	8 8 14 5.4	7 21 37 27	8 10 21 5.5
21	7 16 19 49	8 21 17 59.4	8 3 48 54	8 23 31 40.5
22	7 27 38 48	9 4 21 53.4	8 16 0 21	9 6 42 15.6
23	8 8 57 47	9 17 25 47.3	8 28 11 47	9 19 52 50.6
24	8 20 16 47	10 0 29 41.3	9 10 23 14	10 3 3 25.6
25	9 1 35 46	10 13 33 35.3	9 22 34 41	10 16 14 0.7
26	9 12 54 46	10 26 37 29.2	10 4 46 7	10 29 24 35.7
27	9 24 13 45	11 9 41 23.2	10 16 57 34	11 12 35 10.7
28	10 5 32 45	11 22 45 17.2	10 29 9 1	11 25 45 45.7
29	10 16 51 44	0 5 49 11.1	11 11 20 28	0 8 56 20.8
30	10 28 10 43	0 18 53 5.1	11 23 31 54	0 22 6 55.8
31	11 9 29 43	1 1 56 59.1	0 5 43 21	1 5 17 30.8

*Moon's Motions for Days.*

D.	Supp. of Node.			II	V	VI	VII	VIII	IX	X	XI	XII
	<i>s</i>	<i>o</i>	<i>'</i>	<i>s</i>	<i>o</i>	<i>'</i>						
1	0	0	0	0	0	0	000	000	000	000	000	000
2	0	0	3	0	11	9	034	039	028	034	005	042
3	0	0	6	0	22	18	068	079	056	067	011	083
4	0	0	9	1	3	27	102	118	085	101	016	125
5	0	0	12	1	14	37	136	158	113	135	143	166
6	0	0	15	1	25	46	170	197	141	169	179	208
7	0	0	19	2	6	55	204	237	169	202	215	250
8	0	0	22	2	18	4	238	276	198	236	251	291
9	0	0	25	2	29	13	272	316	226	270	287	333
10	0	0	28	3	10	22	306	355	254	303	323	374
11	0	0	31	3	21	31	340	395	282	337	358	416
12	0	0	34	4	2	40	374	434	311	371	394	458
13	0	0	38	4	13	50	408	474	339	405	430	499
14	0	0	41	4	24	59	442	513	367	438	466	541
15	0	0	44	5	6	8	476	553	395	472	502	583
16	0	0	47	5	17	17	510	592	424	506	538	624
17	0	0	50	5	28	26	544	632	452	539	573	666
18	0	0	54	6	9	35	578	671	480	573	609	707
19	0	0	57	6	20	44	612	711	508	607	645	749
20	0	1	0	7	1	53	646	750	537	641	681	791
21	0	1	3	7	13	3	680	790	565	674	717	832
22	0	1	6	7	24	12	714	829	593	708	753	874
23	0	1	9	8	5	21	748	869	621	742	788	915
24	0	1	13	8	16	30	782	908	650	775	824	957
25	0	1	16	8	27	39	816	948	678	809	860	999
26	0	1	19	9	8	48	850	987	706	843	896	1040
27	0	1	22	9	19	57	884	1027	734	877	932	1082
28	0	1	25	10	1	6	918	1066	762	910	968	1123
29	0	1	28	10	12	16	952	1106	791	944	1003	1165
30	0	1	32	10	23	25	986	1145	819	978	1039	1207
31	0	1	35	11	4	34	1020	1185	847	1011	1075	1248

*Moon's Motions for Hours.*

H.	1	2	3	4	5	6	7	8	9	10	11	12	13
1	11	27	43	12	14	16	2	16	1	3	1	3	1
2	23	54	87	24	28	31	5	33	2	6	3	6	3
3	34	81	130	36	42	47	7	49	3	9	4	9	4
4	46	108	173	48	56	62	10	65	4	12	5	12	6
5	57	135	217	60	70	78	12	81	5	15	6	15	7
6	68	162	260	72	84	93	14	98	6	18	8	18	9
7	80	190	303	84	98	109	17	114	7	20	9	20	10
8	91	217	347	96	112	124	19	130	8	23	10	23	11
9	103	244	390	108	126	140	22	146	9	26	12	26	13
10	114	271	433	120	140	155	24	163	10	29	13	29	14
11	125	298	477	131	154	171	26	179	11	32	14	32	16
12	137	325	520	143	168	186	29	195	12	35	16	35	17
13	148	352	563	155	182	202	31	211	13	38	17	38	18
14	160	379	607	167	196	217	34	228	14	41	18	41	20
15	171	406	650	179	210	233	36	244	15	44	19	44	21
16	182	433	693	191	224	248	38	260	16	47	21	47	23
17	194	460	737	203	238	264	41	276	17	50	22	50	24
18	205	487	780	215	252	279	43	293	18	53	23	53	25
19	217	515	823	227	266	295	46	309	19	56	25	56	27
20	228	542	867	239	280	310	48	325	20	58	26	58	28
21	239	569	910	251	294	326	50	341	21	61	27	61	30
22	251	596	953	263	308	341	53	358	22	64	28	64	31
23	262	623	997	275	322	357	55	374	23	67	30	67	33
24	274	650	1040	287	336	372	58	390	24	70	31	70	34

Hours.	Evection.	Anomaly.	Variation.	Longitude.
	° ' "	° ' "	° ' "	° ' "
1	0 28 17	0 32 39.7	0 30 29	0 32 56.5
2	0 56 35	1 5 19.5	1 0 57	1 5 52.9
3	1 24 52	1 37 59.2	1 31 26	1 38 49.4
4	1 53 10	2 10 39.0	2 1 54	2 11 45.8
5	2 21 27	2 43 18.7	2 32 23	2 44 42.3
6	2 49 45	3 15 58.5	3 2 52	3 17 38.8
7	3 18 2	3 48 38.2	3 33 20	3 50 35.2
8	3 46 20	4 21 18.0	4 3 49	4 23 31.7
9	4 14 37	4 53 57.7	4 34 17	4 56 28.1
10	4 42 55	5 26 37.5	5 4 46	5 29 24.6
11	5 11 12	5 59 17.2	5 35 15	6 2 21.0
12	5 39 30	6 31 57.0	6 5 43	6 35 17.5
13	6 7 47	7 4 36.7	6 36 12	7 8 14.0
14	6 36 5	7 37 16.5	7 6 40	7 41 10.4
15	7 4 22	8 9 56.2	7 37 9	8 14 6.9
16	7 32 40	8 42 36.0	8 7 38	8 47 3.4
17	8 0 57	9 15 15.7	8 38 6	9 19 59.8
18	8 29 15	9 47 55.5	9 8 35	9 52 56.3
19	8 57 32	10 20 35.2	9 39 3	10 25 52.7
20	9 25 50	10 53 15.0	10 9 32	10 58 49.2
21	9 54 7	11 25 54.7	10 40 1	11 31 45.6
22	10 22 24	11 58 34.5	11 10 29	12 4 42.1
23	10 50 42	12 31 14.2	11 40 58	12 37 38.6
24	11 18 59	13 3 54.0	12 11 27	13 10 35.0



*Moon's Motions for Hours.*

H.	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29
1	4	1	2	2	1	0	0	0	0	0	0	0	0	0	0	0
2	8	3	3	4	2	0	0	0	0	1	0	0	0	1	0	0
3	12	4	5	5	3	0	0	0	0	1	0	1	0	1	1	0
4	16	5	6	7	4	0	0	0	0	2	0	1	1	1	1	1
5	21	6	8	9	5	0	0	0	0	2	1	1	1	1	1	1
6	25	8	9	11	6	0	0	0	0	3	1	1	1	2	1	1
7	29	9	11	12	8	1	0	0	0	3	1	1	1	2	1	1
8	33	10	12	14	9	1	0	0	0	3	1	1	1	2	1	1
9	37	11	14	16	10	1	0	0	0	4	1	2	1	3	1	1
10	41	13	15	18	11	1	0	0	0	4	1	2	2	3	2	1
11	45	14	17	19	12	1	0	0	0	5	1	2	2	3	2	1
12	49	15	18	21	13	1	0	0	0	5	1	2	2	3	2	2
13	54	16	20	23	14	1	0	0	0	5	1	2	2	4	2	2
14	58	18	21	25	15	1	0	0	0	6	2	2	2	4	2	2
15	62	19	23	26	16	1	0	0	0	6	2	3	2	4	3	2
16	66	20	25	28	17	1	0	1	1	7	2	3	2	5	3	2
17	70	21	26	30	18	1	0	1	1	7	2	3	3	5	3	2
18	74	23	28	32	19	2	0	1	1	8	2	3	3	5	3	2
19	78	24	29	33	21	2	0	1	1	8	2	3	3	6	3	3
20	83	25	31	35	22	2	0	1	1	8	2	3	3	6	3	3
21	87	26	32	37	23	2	0	1	1	9	2	4	3	6	4	3
22	91	28	34	39	24	2	0	1	1	9	2	4	3	6	4	3
23	95	29	35	40	25	2	0	1	1	10	3	4	4	7	4	3
24	99	31	37	42	26	2	0	1	1	10	3	4	4	7	4	3

H.	Sup. of Nod.	II	V	VI	VII	VIII	IX	X	XI	XII
1	0 7.9	0 28	1	2	1	1	1	0	2	1
2	0 15.9	0 56	3	3	2	3	3	0	3	3
3	0 23.8	1 24	4	5	4	4	4	1	5	4
4	0 31.8	1 52	6	7	5	6	6	1	7	6
5	0 39.7	2 19	7	8	6	7	7	1	9	7
6	0 47.7	2 47	9	10	7	9	9	1	10	9
7	0 55.6	3 15	10	12	8	10	10	2	12	10
8	1 3.6	3 43	11	13	9	11	12	2	14	11
9	1 11.5	4 11	13	15	11	13	13	2	15	13
10	1 19.4	4 39	14	16	12	14	15	2	17	14
11	1 27.4	5 7	16	18	13	15	16	2	19	15
12	1 35.3	5 35	17	20	14	17	18	3	21	17
13	1 43.3	6 2	18	21	15	18	19	3	23	18
14	1 51.2	6 30	20	23	16	19	21	3	24	19
15	1 59.2	6 58	21	25	18	21	22	3	26	21
16	2 7.1	7 26	23	26	19	22	24	4	28	22
17	2 15.0	7 54	24	28	20	24	25	4	29	24
18	2 23.0	8 22	26	29	21	25	27	4	31	25
19	2 30.9	8 50	27	31	22	27	28	4	33	27
20	2 38.9	9 18	28	32	24	28	30	4	35	28
21	2 46.8	9 45	30	34	25	29	31	5	37	29
22	2 54.8	10 13	31	36	26	31	33	5	38	31
23	3 2.7	10 41	33	38	27	32	34	5	40	32
24	3 10.6	11 9	34	39	28	34	36	5	42	34

*Moon's Motions for Minutes.*

Min.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
2	0	1	1	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0
3	1	1	2	1	1	1	0	1	0	0	0	0	0	0	0	0	0	0
4	1	2	3	1	1	1	0	1	0	0	0	0	0	0	0	0	0	0
5	1	2	4	1	1	1	0	1	0	0	0	0	0	0	0	0	0	0
6	1	3	4	1	1	2	0	2	0	0	0	0	0	0	0	0	0	0
7	1	3	5	1	2	2	0	2	0	0	0	0	0	0	0	0	0	0
8	2	4	6	2	2	2	0	2	0	0	0	0	0	1	0	0	0	0
9	2	4	6	2	2	2	0	2	0	0	0	0	0	1	0	0	0	0
10	2	5	7	2	2	3	0	3	0	0	0	0	0	1	0	0	0	0
11	2	5	8	2	3	3	0	3	0	1	0	1	0	1	0	0	0	0
12	2	5	9	2	3	3	0	3	0	1	0	1	0	1	0	0	0	0
13	2	6	9	3	3	3	1	4	0	1	0	1	0	1	0	0	0	0
14	3	6	10	3	3	4	1	4	0	1	0	1	0	1	0	0	0	0
15	3	7	11	3	3	4	1	4	0	1	0	1	0	1	0	0	0	0
16	3	7	12	3	4	4	1	4	0	1	0	1	0	1	0	0	0	0
17	3	8	12	3	4	4	1	5	0	1	0	1	0	1	0	0	0	0
18	3	8	13	4	4	5	1	5	0	1	0	1	0	1	0	0	1	0
19	4	9	14	4	4	5	1	5	0	1	0	1	0	1	0	0	1	0
20	4	9	14	4	5	5	1	5	0	1	0	1	0	1	0	1	1	0
21	4	10	15	4	5	5	1	6	0	1	0	1	0	1	0	1	1	0
22	4	10	16	4	5	6	1	6	0	1	0	1	1	2	0	1	1	0
23	4	10	17	5	5	6	1	6	0	1	0	1	1	2	0	1	1	0
24	5	11	17	5	6	6	1	7	0	1	1	1	1	2	1	1	1	0
25	5	11	18	5	6	6	1	7	0	1	1	1	1	2	1	1	1	0
26	5	12	19	5	6	7	1	7	0	1	1	1	1	2	1	1	1	0
27	5	12	19	5	6	7	1	7	0	1	1	1	1	2	1	1	1	0
28	5	13	20	6	7	7	1	8	0	1	1	1	1	2	1	1	1	0
29	6	13	21	6	7	7	1	8	0	1	1	1	1	2	1	1	1	0
30	6	14	22	6	7	8	1	8	0	1	1	1	1	2	1	1	1	0

*Moon's Motions for Minutes.*

Min.	Evec.	Anom.	Varia.	Long.	Sup. Nod.	II	V	VI	VII	VIII	IX	XI	XII
1	0 28	0 32.7	0 30	0 32.9	0.1	0	0	0	0	0	0	0	0
2	0 57	1 5.3	1 1	1 5.9	0.3	1	0	0	0	0	0	0	0
3	1 25	1 38.0	1 31	1 38.8	0.4	1	0	0	0	0	0	0	0
4	1 53	2 10.6	2 2	2 11.8	0.5	2	0	0	0	0	0	0	0
5	2 21	2 43.3	2 32	2 44.7	0.7	2	0	0	0	0	0	0	0
6	2 50	3 16.0	3 3	3 17.6	0.8	3	0	0	0	0	0	0	0
7	3 18	3 48.6	3 33	3 50.6	0.9	3	0	0	0	0	0	0	0
8	3 46	4 21.3	4 4	4 23.5	1.1	4	0	0	0	0	0	0	0
9	4 15	4 54.0	4 34	4 56.5	1.2	4	0	0	0	0	0	0	0
10	4 43	5 26.6	5 5	5 29.4	1.3	5	0	0	0	0	0	0	0
11	5 11	5 59.3	5 35	6 2.4	1.5	5	0	0	0	0	0	0	0
12	5 40	6 31.9	6 6	6 35.3	1.6	6	0	0	0	0	0	0	0
13	6 8	7 4.6	6 36	7 8.2	1.7	6	0	0	0	0	0	0	0
14	6 36	7 37.3	7 7	7 41.2	1.9	7	0	0	0	0	0	0	0
15	7 4	8 9.9	7 37	8 14.1	2.0	7	0	0	0	0	0	0	0
16	7 33	8 42.6	8 8	8 47.1	2.1	7	0	0	0	0	0	0	0
17	8 1	9 15.3	8 38	9 20.0	2.3	8	0	0	0	0	0	0	0
18	8 29	9 47.9	9 9	9 52.9	2.4	8	0	0	0	0	0	1	0
19	8 58	10 20.6	9 39	10 25.9	2.5	9	0	0	0	0	0	1	0
20	9 26	10 53.2	10 10	10 58.8	2.6	9	0	1	0	0	0	1	0
21	9 54	11 25.9	10 40	11 31.8	2.8	10	0	1	0	0	0	1	0
22	10 22	11 58.6	11 11	12 4.7	2.9	10	1	1	0	0	1	1	0
23	10 51	12 31.2	11 41	12 37.6	3.0	11	1	1	0	0	1	1	0
24	11 19	13 3.9	12 12	13 10.6	3.2	11	1	1	0	1	1	1	1
25	11 47	13 36.6	12 42	13 43.5	3.3	12	1	1	0	1	1	1	1
26	12 16	14 9.2	13 13	14 16.5	3.4	12	1	1	1	1	1	1	1
27	12 44	14 41.9	13 43	14 49.4	3.6	13	1	1	1	1	1	1	1
28	13 12	15 14.6	14 13	15 22.3	3.7	13	1	1	1	1	1	1	1
29	13 40	15 47.2	14 44	15 55.3	3.8	13	1	1	1	1	1	1	1
30	14 9	16 19.9	15 14	16 28.2	4.0	14	1	1	1	1	1	1	1

*Moon's Motions for Minutes.*

Min.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
31	6	14	22	6	7	8	1	8	0	1	1	1	1	2	1	1	1	1
32	6	14	23	6	7	8	1	9	1	2	1	2	1	2	1	1	1	1
33	6	15	24	7	8	9	1	9	1	2	1	2	1	2	1	1	1	1
34	6	15	25	7	8	9	1	9	1	2	1	2	1	2	1	1	1	1
35	7	16	25	7	8	9	1	10	1	2	1	2	1	2	1	1	1	1
36	7	16	26	7	8	9	1	10	1	2	1	2	1	3	1	1	1	1
37	7	17	27	7	9	10	1	10	1	2	1	2	1	3	1	1	1	1
38	7	17	27	8	9	10	2	10	1	2	1	2	1	3	1	1	1	1
39	7	18	28	8	9	10	2	11	1	2	1	2	1	3	1	1	1	1
40	8	18	29	8	9	10	2	11	1	2	1	2	1	3	1	1	1	1
41	8	19	30	8	10	11	2	11	1	2	1	2	1	3	1	1	1	1
42	8	19	30	8	10	11	2	11	1	2	1	2	1	3	1	1	1	1
43	8	19	31	9	10	11	2	12	1	2	1	2	1	3	1	1	1	1
44	8	20	32	9	10	11	2	12	1	2	1	2	1	3	1	1	1	1
45	9	20	32	9	10	12	2	12	1	2	1	2	1	3	1	1	1	1
46	9	21	33	9	11	12	2	12	1	2	1	2	1	3	1	1	1	1
47	9	21	34	9	11	12	2	13	1	2	1	2	1	3	1	1	1	1
48	9	22	35	10	11	12	2	13	1	2	1	2	1	3	1	1	1	1
49	9	22	35	10	11	13	2	13	1	2	1	2	1	3	1	1	1	1
50	9	23	36	10	11	13	2	13	1	2	1	2	1	3	1	1	1	1
51	10	23	37	10	12	13	2	14	1	2	1	2	1	4	1	1	1	1
52	10	24	38	10	12	13	2	14	1	3	1	3	1	4	1	1	1	1
53	10	24	38	11	12	14	2	14	1	3	1	3	1	4	1	1	1	1
54	10	24	39	11	12	14	2	14	1	3	1	3	1	4	1	1	2	1
55	10	25	40	11	13	14	2	15	1	3	1	3	1	4	1	1	2	1
56	11	25	40	11	13	14	2	15	1	3	1	3	1	4	1	1	2	1
57	11	26	41	11	13	15	2	15	1	3	1	3	1	4	1	1	2	1
58	11	26	42	12	13	15	2	16	1	3	1	3	1	4	1	2	2	1
59	11	27	43	12	14	15	2	16	1	3	1	3	1	4	1	2	2	1
60	11	27	43	12	14	15	2	16	1	3	1	3	1	4	1	2	2	1

*Moon's Motions for Minutes.*

Min.	Evcc.	Anom.	Varia.	Long.	Sup. Nod.	II	V	VI	VII	VIII	IX	XI	XII
	' "	' "	' "	' "	"								
31	14 37	16 52.5	15 45	17 1.2	4.1	14	1	1	1	1	1	1	1
32	15 5	17 25.2	16 15	17 34.1	4.2	15	1	1	1	1	1	1	1
33	15 34	17 57.9	16 46	18 7.1	4.4	15	1	1	1	1	1	1	1
34	16 2	18 30.5	17 16	18 40.0	4.5	16	1	1	1	1	1	1	1
35	16 30	19 3.2	17 47	19 12.9	4.7	16	1	1	1	1	1	1	1
36	16 58	19 35.8	18 17	19 45.9	4.8	17	1	1	1	1	1	1	1
37	17 27	20 8.5	18 48	20 18.8	4.9	17	1	1	1	1	1	1	1
38	17 55	20 41.2	19 18	20 51.8	5.0	18	1	1	1	1	1	1	1
39	18 23	21 13.8	19 49	21 24.7	5.2	18	1	1	1	1	1	1	1
40	18 52	21 46.5	20 19	21 57.6	5.3	19	1	1	1	1	1	1	1
41	19 20	22 19.2	20 50	22 30.6	5.4	19	1	1	1	1	1	1	1
42	19 48	22 51.8	21 20	23 3.5	5.6	20	1	1	1	1	1	1	1
43	20 16	23 24.5	21 51	23 36.5	5.7	20	1	1	1	1	1	1	1
44	20 45	23 57.1	22 21	24 9.4	5.8	21	1	1	1	1	1	1	1
45	21 13	24 29.8	22 52	24 42.3	6.0	21	1	1	1	1	1	1	1
46	21 41	25 2.5	23 22	25 15.3	6.1	21	1	1	1	1	1	1	1
47	22 10	25 35.1	23 53	25 48.2	6.2	22	1	1	1	1	1	1	1
48	22 38	26 7.8	24 23	26 21.2	6.4	22	1	1	1	1	1	1	1
49	23 6	26 40.5	24 54	26 54.1	6.5	23	1	1	1	1	1	1	1
50	23 34	27 13.1	25 24	27 27.0	6.6	23	1	1	1	1	1	1	1
51	24 3	27 45.8	25 55	28 0.0	6.8	24	1	1	1	1	1	1	1
52	24 31	28 18.5	26 25	28 32.9	6.9	24	1	1	1	1	1	1	1
53	24 59	28 51.1	26 56	29 5.9	7.0	25	1	1	1	1	1	1	1
54	25 28	29 23.8	27 26	29 38.8	7.1	25	1	1	1	1	1	2	1
55	25 56	29 56.4	27 56	30 11.8	7.3	26	1	1	1	1	1	2	1
56	26 24	30 29.1	28 27	30 44.7	7.4	26	1	1	1	1	1	2	1
57	26 52	31 1.8	28 57	31 17.6	7.5	27	1	2	1	1	1	2	1
58	27 21	31 34.4	29 28	31 50.6	7.7	27	1	2	1	1	1	2	1
59	27 49	32 7.1	29 58	32 23.5	7.8	28	1	2	1	1	1	2	1
60	28 17	32 39.8	30 29	32 56.5	7.9	28	1	2	1	1	1	2	1



*Moon's Motions for Seconds.*

Sec.	Evec.	Anom.	Var.	Long.	Sec.	Evec.	Anom.	Var.	Long.
	"	"	"	"		"	"	"	"
1	0	0.5	1	0.5	31	15	16.9	16	17.0
2	1	1.1	1	1.1	32	15	17.4	16	17.6
3	1	1.6	2	1.6	33	16	18.0	17	18.1
4	2	2.2	2	2.2	34	16	18.5	17	18.7
5	2	2.7	3	2.7	35	17	19.1	18	19.2
6	3	3.3	3	3.3	36	17	19.6	18	19.8
7	3	3.8	4	3.8	37	18	20.1	19	20.3
8	4	4.3	4	4.4	38	18	20.7	19	20.9
9	4	4.9	5	4.9	39	18	21.2	20	21.4
10	5	5.4	5	5.5	40	19	21.8	20	22.0
11	5	6.0	6	6.0	41	19	22.3	21	22.5
12	6	6.5	6	6.6	42	20	22.9	21	23.1
13	6	7.1	7	7.1	43	20	23.4	22	23.6
14	7	7.6	7	7.7	44	21	24.0	22	24.2
15	7	8.2	8	8.2	45	21	24.5	23	24.7
16	8	8.7	8	8.8	46	22	25.0	23	25.3
17	8	9.2	9	9.3	47	22	25.6	24	25.8
18	9	9.8	9	9.9	48	23	26.1	24	26.4
19	9	10.3	10	10.4	49	23	26.7	25	26.9
20	9	10.9	10	11.0	50	24	27.2	25	27.4
21	10	11.4	11	11.5	51	24	27.8	26	28.0
22	10	12.0	11	12.1	52	25	28.3	26	28.5
23	11	12.5	12	12.6	53	25	28.9	27	29.1
24	11	13.1	12	13.2	54	26	29.4	27	29.6
25	12	13.6	13	13.7	55	26	29.9	28	30.2
26	12	14.1	13	14.3	56	26	30.5	28	30.7
27	13	14.7	14	14.8	57	27	31.0	29	31.3
28	13	15.2	14	15.4	58	27	31.6	29	31.8
29	14	15.8	15	15.9	59	28	32.1	30	32.4
30	14	16.3	15	16.5	60	28	32.7	30	32.9

*First Equation of Moon's Longitude.—Argument 1.*

Arg.	1	Diff. for 10	Arg.	1	Diff. for 10	Arg.	1	Diff. for 10	Arg.	1	Diff. for 10
0	12 40.0	"	2500	1 40.7	"	5000	12 40.0	"	7500	23 39.3	"
50	12 18.8	4.24	2550	1 41.5	0.16	5050	13 0.3	4.06	7550	23 39.4	0.02
100	11 57.7	4.22	2600	1 42.9	0.28	5100	13 20.5	4.04	7600	23 38.9	0.10
150	11 36.6	4.22	2650	1 45.0	0.42	5150	13 40.7	4.04	7650	23 37.7	0.24
200	11 15.6	4.20	2700	1 47.7	0.54	5200	14 0.9	4.04	7700	23 35.8	0.38
250	10 54.7	4.18	2750	1 51.0	0.66	5250	14 20.9	4.00	7750	23 33.3	0.50
		4.16			0.80			4.00			0.62
300	10 33.9		2800	1 55.0		5300	14 40.9		7800	23 30.2	
350	10 13.2	4.14	2850	1 59.6	0.92	5350	15 0.8	3.98	7850	23 26.4	0.76
400	9 52.6	4.12	2900	2 4.8	1.04	5400	15 20.5	3.94	7900	23 22.0	0.88
450	9 32.3	4.06	2950	2 10.7	1.18	5450	15 40.1	3.92	7950	23 16.9	1.02
500	9 12.1	4.04	3000	2 17.1	1.28	5500	15 59.6	3.90	8000	23 11.2	1.14
		4.00			1.42			3.84			1.26
550	8 52.1		3050	2 24.2		5550	16 18.8		8050	23 4.9	
600	8 32.4	3.94	3100	2 31.9	1.54	5600	16 37.8	3.80	8100	22 57.9	1.40
650	8 13.0	3.88	3150	2 40.1	1.64	5650	16 56.7	3.78	8150	22 50.3	1.52
700	7 53.8	3.84	3200	2 48.9	1.76	5700	17 15.3	3.72	8200	22 42.0	1.66
750	7 34.9	3.78	3250	2 58.3	1.88	5750	17 33.6	3.66	8250	22 33.2	1.76
		3.70			1.98			3.60			1.90
800	7 16.4		3300	3 8.2		5800	17 51.6		8300	22 23.7	
850	6 58.2	3.64	3350	3 18.7	2.10	5850	18 9.4	3.56	8350	22 13.7	2.00
900	6 40.3	3.58	3400	3 29.7	2.20	5900	18 26.9	3.50	8400	22 3.1	2.12
950	6 22.8	3.50	3450	3 41.3	2.32	5950	18 44.0	3.42	8450	21 51.9	2.24
1000	6 5.7	3.42	3500	3 53.4	2.42	6000	19 0.8	3.36	8500	21 40.1	2.36
		3.34			2.50			3.28			2.46
1050	5 49.0		3550	4 5.9		6050	19 17.2		8550	21 27.8	
1100	5 32.8	3.24	3600	4 19.0	2.62	6100	19 33.3	3.22	8600	21 15.0	2.56
1150	5 17.0	3.16	3650	4 32.5	2.70	6150	19 49.0	3.14	8650	21 1.6	2.68
1200	5 1.6	3.08	3700	4 46.5	2.80	6200	20 4.2	3.04	8700	20 47.7	2.78
1250	4 46.7	2.98	3750	5 0.9	2.88	6250	20 19.1	2.98	8750	20 33.3	2.88
		2.88			2.98			2.88			2.98
1300	4 32.3		3800	5 15.8		6300	20 33.5		8800	20 18.4	
1350	4 18.4	2.78	3850	5 31.0	3.04	6350	20 47.5	2.80	8850	20 3.0	3.08
1400	4 5.0	2.68	3900	5 46.7	3.14	6400	21 1.0	2.70	8900	19 47.2	3.16
1450	3 52.2	2.56	3950	6 2.8	3.22	6450	21 14.1	2.62	8950	19 31.0	3.24
1500	3 39.9	2.46	4000	6 19.2	3.28	6500	21 26.6	2.50	9000	19 14.3	3.34
		2.36			3.36			2.42			3.42
1550	3 28.1		4050	6 36.0		6550	21 38.7		9050	18 57.2	
1600	3 16.9	2.24	4100	6 53.1	3.42	6600	21 50.3	2.32	9100	18 39.7	3.50
1650	3 6.3	2.12	4150	7 10.6	3.50	6650	22 1.3	2.20	9150	18 21.8	3.58
1700	2 56.3	2.00	4200	7 28.4	3.56	6700	22 11.8	2.10	9200	18 3.6	3.64
1750	2 46.8	1.90	4250	7 46.4	3.60	6750	22 21.7	1.98	9250	17 45.1	3.70
		1.76			3.66			1.88			3.78
1800	2 38.0		4300	8 4.7		6800	22 31.1		9300	17 26.2	
1850	2 29.7	1.66	4350	8 23.3	3.72	6850	22 39.9	1.76	9350	17 7.0	3.84
1900	2 22.1	1.52	4400	8 42.2	3.78	6900	22 48.1	1.64	9400	16 47.6	3.88
1950	2 15.1	1.40	4450	9 1.2	3.80	6950	22 55.8	1.54	9450	16 27.9	3.94
2000	2 8.8	1.26	4500	9 20.4	3.84	7000	23 2.9	1.42	9500	16 7.9	4.00
		1.14			3.90			1.28			4.04
2050	2 3.1		4550	9 39.9		7050	23 9.3		9550	15 47.7	
2100	1 58.0	1.02	4600	9 59.5	3.92	7100	23 15.2	1.18	9600	15 27.4	4.06
2150	1 53.6	0.88	4650	10 19.2	3.94	7150	23 20.4	1.04	9650	15 6.8	4.12
2200	1 49.8	0.76	4700	10 39.1	3.98	7200	23 25.0	0.92	9700	14 46.1	4.14
2250	1 46.7	0.62	4750	10 59.1	4.00	7250	23 29.0	0.80	9750	14 25.3	4.16
		0.50			4.00			0.66			4.18
2300	1 44.2		4800	11 19.1		7300	23 32.3		9800	14 4.4	
2350	1 42.3	0.38	4850	11 39.3	4.04	7350	23 35.0	0.54	9850	13 43.4	4.20
2400	1 41.1	0.24	4900	11 59.5	4.04	7400	23 37.1	0.42	9900	13 22.3	4.22
2450	1 40.6	0.10	4950	12 19.7	4.04	7450	23 38.5	0.28	9950	13 1.2	4.22
2500	1 40.7	0.02	5000	12 40.0	4.06	7500	23 39.3	0.16	10000	12 40.0	4.24

*Equations 2 to 7 of Moon's Longitude. Arguments 2 to 7.*

Arg.	2	diff	3	diff	4	diff	5	diff	6	diff	7	diff	Arg.
2500	4 57.3	"	0 2.3	"	6 30.3	"	3 39.4	"	0 6.2	"	0 0.8	"	2500
2600	4 57.0	0.3	0 2.4	0.1	6 29.9	0.4	3 39.2	0.2	0 6.4	0.2	0 0.9	0.1	2400
2700	4 56.1	0.9	0 2.8	0.4	6 28.8	1.1	3 38.5	0.7	0 6.9	0.5	0 1.3	0.4	2300
2800	4 54.7	1.4	0 3.3	0.5	6 26.9	1.9	3 37.5	1.0	0 7.7	0.8	0 1.8	0.5	2200
2900	4 52.7	2.0	0 4.1	0.8	6 24.3	2.6	3 36.0	1.5	0 8.8	1.1	0 2.7	0.9	2100
3000	4 50.1	2.6	0 5.1	1.0	6 21.0	3.3	3 34.1	1.9	0 10.3	1.5	0 3.7	1.0	2000
3100	4 47.0	3.1	0 6.4	1.3	6 16.9	4.1	3 31.7	2.4	0 12.1	1.8	0 5.0	1.3	1900
3200	4 43.3	3.7	0 7.8	1.4	6 12.2	4.7	3 29.0	2.7	0 14.2	2.1	0 6.4	1.4	1800
3300	4 39.1	4.2	0 9.4	1.6	6 6.8	5.4	3 25.9	3.1	0 16.6	2.4	0 8.1	1.7	1700
3400	4 34.4	4.7	0 11.3	1.9	6 0.7	6.1	3 22.4	3.5	0 19.2	2.6	0 10.0	1.9	1600
3500	4 29.2	5.2	0 13.3	2.0	5 54.0	6.7	3 18.5	3.9	0 22.2	3.0	0 12.1	2.1	1500
3600	4 23.5	5.7	0 15.5	2.2	5 46.6	7.4	3 14.3	4.2	0 25.4	3.2	0 14.4	2.3	1400
3700	4 17.4	6.1	0 17.9	2.4	5 38.7	7.9	3 9.7	4.6	0 28.9	3.5	0 16.8	2.4	1300
3800	4 10.8	6.6	0 20.5	2.6	5 30.3	8.4	3 4.9	4.8	0 32.7	3.8	0 19.5	2.7	1200
3900	4 3.9	6.9	0 23.2	2.7	5 21.3	9.0	2 59.7	5.2	0 36.6	3.9	0 22.3	2.8	1100
4000	3 56.6	7.3	0 26.1	2.9	5 11.9	9.4	2 54.3	5.4	0 40.7	4.1	0 25.2	2.9	1000
4100	3 48.9	7.7	0 29.1	3.0	5 2.0	9.9	2 48.0	5.7	0 45.1	4.4	0 28.3	3.1	900
4200	3 41.0	7.9	0 32.2	3.1	4 51.7	10.3	2 42.7	5.9	0 49.6	4.5	0 31.5	3.2	800
4300	3 32.7	8.3	0 35.4	3.2	4 41.0	10.7	2 36.6	6.1	0 49.6	4.7	0 31.5	3.3	700
4400	3 24.2	8.5	0 38.8	3.4	4 30.1	10.9	2 30.3	6.3	0 54.3	4.9	0 34.8	3.4	600
4500	3 15.5	8.7	0 42.2	3.4	4 18.8	11.3	2 23.8	6.5	0 59.2	4.9	0 38.2	3.5	500
4600	3 6.6	8.9	0 45.7	3.5	4 7.3	11.5	2 17.2	6.6	1 4.1	5.1	0 41.7	3.6	400
4700	2 57.6	9.0	0 49.2	3.5	3 55.7	11.6	2 10.5	6.7	1 9.2	5.1	0 45.3	3.6	300
4800	2 48.5	9.1	0 52.8	3.6	3 43.9	11.8	2 3.7	6.8	1 14.3	5.2	0 48.9	3.7	200
4900	2 39.2	9.3	0 56.4	3.6	3 31.9	12.0	1 56.9	6.8	1 19.5	5.2	0 52.6	3.7	100
5000	2 30.0	9.2	1 0.0	3.6	3 20.0	11.9	1 50.0	6.9	1 24.7	5.3	0 56.3	3.7	0
5100	2 20.8	9.2	1 3.6	3.6	3 8.1	11.9	1 43.1	6.9	1 30.0	5.3	1 0.0	3.7	9900
5200	2 11.5	9.3	1 7.2	3.6	2 56.1	12.0	1 36.3	6.8	1 35.3	5.2	1 3.7	3.7	9800
5300	2 2.4	9.1	1 10.8	3.6	2 44.3	11.8	1 29.5	6.8	1 40.5	5.2	1 7.4	3.7	9700
5400	1 53.4	9.0	1 14.3	3.5	2 32.7	11.6	1 22.8	6.7	1 45.7	5.1	1 11.1	3.6	9600
5500	1 44.5	8.9	1 17.8	3.5	2 21.2	11.5	1 16.2	6.6	1 50.8	5.1	1 14.7	3.6	9500
5600	1 35.8	8.7	1 21.2	3.4	2 9.9	11.3	1 9.7	6.5	1 55.9	4.9	1 18.3	3.5	9400
5700	1 27.3	8.5	1 24.6	3.4	1 59.0	10.9	1 3.4	6.3	2 0.8	4.9	1 21.8	3.4	9300
5800	1 19.0	8.3	1 27.8	3.2	1 48.3	10.7	0 57.3	6.1	2 5.7	4.7	1 25.2	3.3	9200
5900	1 11.1	7.9	1 30.9	3.1	1 38.0	10.3	0 51.4	5.9	2 10.4	4.5	1 28.5	3.2	9100
6000	1 3.4	7.7	1 33.9	3.0	1 28.1	9.9	0 45.7	5.7	2 14.9	4.4	1 31.7	3.1	9000
6100	0 56.1	7.3	1 36.8	2.9	1 18.7	9.4	0 40.3	5.4	2 19.3	4.1	1 34.8	2.9	8900
6200	0 49.2	6.9	1 39.5	2.7	1 9.7	9.0	0 35.1	5.2	2 23.4	3.9	1 37.7	2.8	8800
6300	0 42.6	6.6	1 42.1	2.6	1 1.3	8.4	0 30.3	4.8	2 27.3	3.8	1 40.5	2.7	8700
6400	0 36.5	6.1	1 44.5	2.4	0 53.4	7.9	0 25.7	4.6	2 31.1	3.5	1 43.2	2.4	8600
6500	0 30.8	5.7	1 46.7	2.2	0 46.0	7.4	0 21.5	4.2	2 34.6	3.2	1 45.6	2.3	8500
6600	0 25.6	5.2	1 48.7	2.0	0 39.3	6.7	0 17.6	3.9	2 37.8	3.0	1 47.9	2.1	8400
6700	0 20.9	4.7	1 50.6	1.9	0 33.2	6.1	0 14.1	3.5	2 40.8	2.6	1 50.0	1.9	8300
6800	0 16.7	4.2	1 52.2	1.6	0 27.8	5.4	0 11.0	3.1	2 43.4	2.4	1 51.9	1.7	8200
6900	0 13.0	3.7	1 53.6	1.4	0 23.1	4.7	0 8.3	2.7	2 45.8	2.1	1 53.6	1.4	8100
7000	0 9.9	3.1	1 54.9	1.3	0 19.0	4.1	0 5.9	2.4	2 47.9	1.8	1 55.0	1.3	8000
7100	0 7.3	2.6	1 55.9	1.0	0 15.7	3.3	0 4.0	1.9	2 49.7	1.5	1 56.3	1.0	7900
7200	0 5.3	2.0	1 56.7	0.8	0 13.1	2.6	0 2.5	1.5	2 51.2	1.1	1 57.3	0.9	7800
7300	0 3.9	1.4	1 57.2	0.5	0 11.2	1.9	0 1.5	1.0	2 52.3	0.8	1 58.2	0.5	7700
7400	0 3.0	0.9	1 57.6	0.4	0 10.1	1.1	0 0.8	0.7	2 53.1	0.5	1 58.7	0.4	7600
7500	0 2.7	0.3	1 57.7	0.1	0 9.7	0.4	0 0.6	0.2	2 53.8	0.2	1 59.2	0.1	7500

TABLE XLIII.

Equations 8 and 9.

Arg.	8	9	Arg.	8	9
" "	" "	" "	" "	" "	" "
0	1 20.0	1 20 0	5000	1 20.0	1 20.0
100	1 15.5	1 28 7	5100	1 24.4	1 25.8
200	1 11.1	1 37.3	5200	1 28.8	1 31.4
300	1 6.7	1 45.7	5300	1 33.1	1 36.9
400	1 2.3	1 53.7	5400	1 37.4	1 42.0
500	0 58.0	2 1.3	5500	1 41.6	1 46.8
600	0 53.8	2 8.3	5600	1 45.8	1 51.0
700	0 49.7	2 14.7	5700	1 49.8	1 54.6
800	0 45.7	2 20.2	5800	1 53.8	1 57.6
900	0 41.9	2 25.0	5900	1 57.6	1 59.8
1000	0 38.2	2 28.9	6000	2 1.2	2 1.3
1100	0 34.7	2 31.9	6100	2 4.7	2 1.9
1200	0 31.4	2 33.9	6200	2 8.0	2 1.7
1300	0 28.2	2 34.9	6300	2 11.2	2 0.7
1400	0 25.3	2 35.0	6400	2 14.1	1 58.8
1500	0 22.6	2 34.1	6500	2 16.8	1 56.1
1600	0 20.1	2 32.2	6600	2 19.3	1 52.5
1700	0 17.9	2 29.5	6700	2 21.6	1 48.3
1800	0 15.9	2 25.9	6800	2 23.7	1 43.4
1900	0 14.2	2 21.5	6900	2 25.4	1 37.8
2000	0 12.7	2 16.4	7000	2 27.0	1 31.7
2100	0 11.5	2 10.7	7100	2 28.2	1 25.1
2200	0 10.5	2 4.4	7200	2 29.2	1 18.2
2300	0 9.9	1 57.7	7300	2 30.0	1 11.1
2400	0 9.5	1 50.7	7400	2 30.4	1 3.8
2500	0 9.4	1 43.5	7500	2 30.6	0 56.5
2600	0 9.6	1 36.2	7600	2 30.5	0 49.3
2700	0 10.1	1 28.9	7700	2 30.1	0 42.3
2800	0 10.8	1 21.8	7800	2 29.5	0 35.6
2900	0 11.8	1 14.9	7900	2 28.5	0 29.3
3000	0 13.0	1 8.3	8000	2 27.3	0 23.6
3100	0 14.6	1 2.2	8100	2 25.8	0 18.5
3200	0 16.3	0 56.6	8200	2 24.1	0 14.1
3300	0 18.4	0 51.7	8300	2 22.1	0 10.5
3400	0 20.7	0 47.5	8400	2 19.9	0 7.8
3500	0 23.2	0 43.9	8500	2 17.4	0 5.9
3600	0 25.9	0 41.2	8600	2 14.7	0 5.0
3700	0 28.8	0 39.3	8700	2 11.8	0 5.1
3800	0 32.0	0 38.3	8800	2 8.6	0 6.1
3900	0 35.3	0 38.1	8900	2 5.3	0 8.1
4000	0 38.8	0 38.7	9000	2 1.8	0 11.1
4100	0 42.4	0 40.2	9100	1 58.1	0 15.0
4200	0 46.2	0 42.4	9200	1 54.3	0 19.8
4300	0 50.2	0 45.4	9300	1 50.3	0 25.3
4400	0 54.2	0 49.0	9400	1 46.2	0 31.7
4500	0 58.4	0 53.2	9500	1 42.0	0 38.7
4600	1 2.6	0 58.0	9600	1 37.7	0 46.3
4700	1 6.9	1 3.1	9700	1 33.3	0 54.3
4800	1 11.2	1 8.6	9800	1 28.9	1 2.7
4900	1 15.6	1 14.2	9900	1 24.5	1 11.3
5000	1 20.0	1 20.0	10000	1 20.0	1 20.0

TABLE XLIV. 67

Equations 10 and 11.

Arg.	10	11	Arg.	10	11
" "	" "	" "	" "	" "	" "
0	10.0	10.0	500	10.0	10.0
10	9.3	11.1	510	9.6	10.8
20	8.6	12.1	520	9.2	11.5
30	8.0	13.1	530	8.9	12.3
40	7.4	14.1	540	8.5	12.9
50	6.8	15.0	550	8.2	13.6
60	6.2	15.8	560	7.9	14.2
70	5.7	16.6	570	7.7	14.6
80	5.3	17.3	580	7.5	15.0
90	4.9	17.9	590	7.4	15.4
100	4.6	18.3	600	7.3	15.6
110	4.3	18.6	610	7.2	15.7
120	4.1	18.9	620	7.3	15.7
130	4.0	19.0	630	7.4	15.6
140	4.0	18.9	640	7.5	15.4
150	4.0	18.8	650	7.8	15.1
160	4.2	18.6	660	8.1	14.7
170	4.4	18.2	670	8.4	14.2
180	4.6	17.7	680	8.7	13.5
190	4.9	17.1	690	9.2	12.8
200	5.3	16.5	700	9.7	12.1
210	5.7	15.7	710	10.2	11.3
220	6.2	14.9	720	10.7	10.4
230	6.7	14.1	730	11.2	9.5
240	7.2	13.2	740	11.7	8.6
250	7.7	12.3	750	12.3	7.7
260	8.3	11.4	760	12.8	6.8
270	8.8	10.5	770	13.3	5.9
280	9.3	9.6	780	13.8	5.1
290	9.8	8.7	790	14.3	4.3
300	10.3	7.9	800	14.7	3.5
310	10.8	7.2	810	15.1	2.9
320	11.3	6.5	820	15.4	2.3
330	11.6	5.8	830	15.6	1.8
340	11.9	5.3	840	15.8	1.4
350	12.2	4.9	850	16.0	1.2
360	12.5	4.6	860	16.0	1.1
370	12.6	4.4	870	16.0	1.0
380	12.7	4.3	880	15.9	1.1
390	12.8	4.3	890	15.7	1.4
400	12.7	4.4	900	15.4	1.7
410	12.6	4.6	910	15.1	2.1
420	12.5	5.0	920	14.7	2.7
430	12.3	5.4	930	14.3	3.4
440	12.1	5.8	940	13.8	4.2
450	11.8	6.4	950	13.2	5.0
460	11.5	7.1	960	12.6	5.9
470	11.1	7.7	970	12.0	6.9
480	10.8	8.5	980	11.4	7.9
490	10.4	9.2	990	10.7	8.9
500	10.0	10.0	1000	10.0	10.0



TABLE XLV.  
Equations 12 to 19.

Arg.	12	13	14	15	16	17	18	19	Arg.
	"	"	"	"	"	"	"	"	
250	2.3	1.6	7.8	0.0	33.7	3.4	16.7	0.4	250
260	2.3	1.6	7.8	0.0	33.7	3.4	16.7	0.4	240
270	2.4	1.7	7.9	0.1	33.6	3.5	16.6	0.4	230
280	2.6	1.9	8.0	0.2	33.5	3.5	16.6	0.5	220
290	2.9	2.2	8.2	0.3	33.2	3.6	16.5	0.5	210
300	3.2	2.5	8.4	0.5	33.0	3.7	16.4	0.6	200
310	3.5	2.9	8.7	0.7	32.7	3.9	16.2	0.7	190
320	4.0	3.4	9.0	1.0	32.4	4.0	16.1	0.8	180
330	4.5	3.9	9.3	1.2	32.0	4.2	15.9	1.0	170
340	5.1	4.4	9.7	1.6	31.6	4.4	15.7	1.1	160
350	5.7	5.1	10.1	1.9	31.1	4.7	15.4	1.3	150
360	6.4	5.8	10.6	2.3	30.6	4.9	15.2	1.5	140
370	7.1	6.6	11.1	2.7	30.1	5.2	14.9	1.7	130
380	7.9	7.4	11.7	3.2	29.4	5.5	14.6	1.9	120
390	8.7	8.3	12.2	3.6	28.7	5.8	14.3	2.1	110
400	9.6	9.2	12.8	4.1	28.0	6.1	13.9	2.3	100
410	10.5	10.1	13.5	4.6	27.3	6.5	13.6	2.5	90
420	11.5	11.1	14.1	5.2	26.6	6.8	13.2	2.8	80
430	12.5	12.2	14.8	5.7	25.8	7.2	12.9	3.1	70
440	13.5	13.2	15.5	6.3	25.0	7.6	12.5	3.3	60
450	14.5	14.3	16.2	6.9	24.2	8.0	12.1	3.6	50
460	15.6	15.4	17.0	7.5	23.4	8.4	11.7	3.9	40
470	16.7	16.5	17.7	8.1	22.6	8.8	11.3	4.1	30
480	17.8	17.7	18.5	8.7	21.7	9.2	10.8	4.4	20
490	18.9	18.8	19.2	9.4	20.9	9.6	10.4	4.7	10
500	20.0	20.0	20.0	10.0	20.0	10.0	10.0	5.0	0
510	21.1	21.2	20.8	10.6	19.1	10.4	9.6	5.3	990
520	22.2	22.3	21.5	11.3	18.3	10.8	9.2	5.6	980
530	23.3	23.5	22.3	11.9	17.4	11.2	8.7	5.9	970
540	24.4	24.6	23.0	12.5	16.6	11.6	8.3	6.1	960
550	25.5	25.7	23.8	13.1	15.8	12.0	7.9	6.4	950
560	26.5	26.8	24.5	13.7	15.0	12.4	7.5	6.7	940
570	27.5	27.8	25.2	14.3	14.2	12.8	7.1	6.9	930
580	28.5	28.9	25.9	14.8	13.4	13.2	6.8	7.2	920
590	29.5	29.9	26.5	15.4	12.7	13.5	6.4	7.5	910
600	30.4	30.8	27.2	15.9	12.0	13.9	6.1	7.7	900
610	31.3	31.7	27.8	16.4	11.3	14.2	5.7	7.9	890
620	32.1	32.6	28.3	16.8	10.6	14.5	5.4	8.1	880
630	32.9	33.4	28.9	17.3	9.9	14.8	5.1	8.3	870
640	33.6	34.2	29.4	17.7	9.4	15.1	4.8	8.5	860
650	34.3	34.9	29.9	18.1	8.9	15.3	4.6	8.7	850
660	34.9	35.6	30.3	18.4	8.4	15.6	4.3	8.9	840
670	35.5	36.1	30.7	18.8	8.0	15.8	4.1	9.0	830
680	36.0	36.6	31.0	19.0	7.6	16.0	3.9	9.2	820
690	36.5	37.1	31.3	19.3	7.3	16.1	3.8	9.3	810
700	36.8	37.5	31.6	19.5	7.0	16.3	3.6	9.4	800
710	37.1	37.8	31.8	19.7	6.8	16.4	3.5	9.5	790
720	37.4	38.1	32.0	19.8	6.5	16.5	3.4	9.5	780
730	37.6	38.3	32.1	19.9	6.4	16.5	3.4	9.6	770
740	37.7	38.4	32.2	20.0	6.3	16.6	3.3	9.6	760
750	37.7	38.4	32.2	20.0	6.3	16.6	3.3	9.6	750

TABLE XLVI.  
Equation 20.

Arg.	20	Arg.
	"	
0	10.0	500
10	10.9	510
20	11.8	520
30	12.7	530
40	13.5	540
50	14.3	550
60	15.0	560
70	15.7	570
80	16.2	580
90	16.7	590
100	17.0	600
110	17.2	610
120	17.4	620
130	17.4	630
140	17.2	640
150	17.0	650
160	16.7	660
170	16.2	670
180	15.7	680
190	15.0	690
200	14.3	700
210	13.5	710
220	12.7	720
230	11.8	730
240	10.9	740
250	10.0	750
260	9.1	760
270	8.2	770
280	7.3	780
290	6.5	790
300	5.7	800
310	5.0	810
320	4.3	820
330	3.8	830
340	3.3	840
350	3.0	850
360	2.8	860
370	2.6	870
380	2.6	880
390	2.8	890
400	3.0	900
410	3.3	910
420	3.8	920
430	4.3	930
440	5.0	940
450	5.7	950
460	6.5	960
470	7.3	970
480	8.2	980
490	9.1	990
500	10.0	1000



TABLE XLVII.

TABLE XLVIII. 69

Equations 21 to 29.

Equations 30 and 31.

Arg	21	22	23	24	25	26	27	28	29	Arg
25	7.8	3.2	7.1	6.1	5.9	4.1	5.8	4.3	5.7	25
27	7.8	3.2	7.1	6.1	5.9	4.1	5.8	4.3	5.7	23
29	7.7	3.3	7.0	6.1	5.9	4.1	5.8	4.3	5.7	21
31	7.6	3.3	7.0	6.0	5.8	4.2	5.7	4.3	5.7	19
33	7.5	3.4	6.8	6.0	5.8	4.2	5.7	4.4	5.6	17
35	7.3	3.5	6.7	5.9	5.7	4.3	5.6	4.4	5.6	15
37	7.0	3.7	6.5	5.8	5.7	4.3	5.6	4.5	5.5	13
39	6.8	3.9	6.3	5.7	5.6	4.4	5.5	4.6	5.4	11
41	6.5	4.0	6.1	5.6	5.5	4.5	5.4	4.6	5.4	09
43	6.2	4.2	5.9	5.5	5.4	4.6	5.3	4.7	5.3	07
45	5.9	4.4	5.6	5.3	5.3	4.7	5.2	4.8	5.2	05
47	5.5	4.7	5.4	5.2	5.2	4.8	5.1	4.9	5.1	03
49	5.2	4.9	5.1	5.1	5.1	4.9	5.0	5.0	5.0	01
51	4.8	5.1	4.9	4.9	4.9	5.1	5.0	5.0	5.0	99
53	4.5	5.3	4.6	4.8	4.8	5.2	4.9	5.1	4.9	97
55	4.1	5.6	4.4	4.7	4.7	5.3	4.8	5.2	4.8	95
57	3.8	5.8	4.1	4.5	4.6	5.4	4.7	5.3	4.7	93
59	3.5	6.0	3.9	4.4	4.5	5.5	4.6	5.4	4.6	91
61	3.2	6.1	3.7	4.3	4.4	5.6	4.5	5.4	4.6	89
63	3.0	6.3	3.5	4.2	4.3	5.7	4.4	5.5	4.5	87
65	2.7	6.5	3.3	4.1	4.3	5.7	4.4	5.6	4.4	85
67	2.5	6.6	3.2	4.0	4.2	5.8	4.3	5.6	4.4	83
69	2.4	6.7	3.0	4.0	4.2	5.8	4.3	5.7	4.3	81
71	2.3	6.7	3.0	3.9	4.1	5.9	4.2	5.7	4.3	79
73	2.2	6.8	2.9	3.9	4.1	5.9	4.2	5.7	4.3	77
75	2.2	6.8	2.9	3.9	4.1	5.9	4.2	5.7	4.3	75

Arg.	30	31
0	5.0	5.0
2	5.0	5.0
4	4.9	5.1
6	4.9	5.1
8	4.8	5.2
10	4.8	5.2
12	4.7	5.3
14	4.6	5.4
16	4.5	5.5
18	4.4	5.5
20	4.2	5.6
22	4.1	5.7
24	4.0	5.8
26	3.9	5.8
28	3.8	5.9
30	3.7	5.9
32	3.7	5.9
34	3.7	5.9
36	3.7	5.9
38	3.8	5.8
40	3.9	5.7
42	4.1	5.6
44	4.3	5.5
46	4.5	5.3
48	4.8	5.2
50	5.0	5.0
52	5.2	4.8
54	5.5	4.7
56	5.7	4.5
58	5.9	4.4
60	6.1	4.3
62	6.2	4.2
64	6.3	4.1
66	6.3	4.1
68	6.3	4.1
70	6.3	4.1
72	6.2	4.1
74	6.2	4.2
76	6.0	4.2
78	5.9	4.3
80	5.8	4.4
82	5.7	4.5
84	5.5	4.6
86	5.4	4.6
88	5.3	4.7
90	5.2	4.8
92	5.1	4.8
94	5.1	4.9
96	5.0	4.9
98	5.0	5.0
100	5.0	5.0

TABLE XLIX.

Equation 32. Argument, Supp. of Node.

	III <sup>s</sup>	IV <sup>s</sup>	V <sup>s</sup>	VI <sup>s</sup>	VII <sup>s</sup>	VIII <sup>s</sup>	
0	3.1	4.0	6.5	10.0	13.5	16.0	30
2	3.1	4.2	6.8	10.2	13.7	16.1	28
4	3.1	4.3	7.0	10.5	13.8	16.2	26
6	3.1	4.4	7.2	10.7	14.0	16.3	24
8	3.2	4.6	7.4	11.0	14.2	16.4	22
10	3.2	4.7	7.6	11.2	14.4	16.5	20
12	3.3	4.9	7.9	11.4	14.6	16.6	18
14	3.3	5.0	8.1	11.7	14.8	16.6	16
16	3.4	5.2	8.3	11.9	15.0	16.7	14
18	3.4	5.4	8.6	12.1	15.1	16.7	12
20	3.5	5.6	8.8	12.4	15.3	16.8	10
22	3.6	5.8	9.0	12.6	15.4	16.8	8
24	3.7	6.0	9.3	12.8	15.6	16.9	6
26	3.8	6.2	9.5	13.0	15.7	16.9	4
28	3.9	6.3	9.8	13.2	15.8	16.9	2
30	4.0	6.5	10.0	13.5	16.0	16.9	0
	II <sup>s</sup>	I <sup>s</sup>	O <sup>s</sup>	XI <sup>s</sup>	X <sup>s</sup>	IX <sup>s</sup>	

Constant 55"

TABLE L.

*Evection.*

Argument.    Evection, corrected.

	O <sup>s</sup>		I <sup>s</sup>		II <sup>s</sup>		III <sup>s</sup>		IV <sup>s</sup>		V <sup>s</sup>	
Diff.	i°	Diff.	2°	Diff.	2°	Diff.	2°	Diff.	2°	Diff.	2°	Diff.
0	30 00.0	85.5	10 43.5	73.2	40 9.7	40.9	50 25.5	2.0	39 8.3	43.4	9 42.0	72.7
1	31 25.5	85.4	11 56.7	72.3	40 50.6	39.5	50 23.5	3.4	38 24.9	44.5	8 29.3	73.0
2	32 50.9	85.4	13 9.0	71.6	41 30.1	38.2	50 20.1	4.9	37 40.4	45.8	7 16.0	74.0
3	34 16.3	85.3	14 20.6	70.7	42 8.3	36.8	50 15.2	6.4	36 54.6	47.0	6 2.0	74.6
4	35 41.6	85.1	15 31.3	69.8	42 45.1	35.5	50 8.8	7.8	36 7.6	48.1	4 47.4	75.2
5	37 6.7	85.1	16 41.1	69.0	43 20.6	34.1	50 1.0	9.3	35 19.5	49.3	3 32.2	75.9
6	38 31.8	84.9	17 50.1	68.1	43 54.7	32.7	49 51.7	10.7	34 30.2	50.5	2 16.3	76.4
7	39 56.7	84.7	18 58.2	67.1	44 27.4	31.4	49 41.0	12.2	33 39.7	51.6	0 59.9	76.9
8	41 21.4	84.4	20 5.3	66.2	44 58.8	29.9	49 28.8	13.7	32 48.1	52.7	59 43.0	77.4
9	42 45.8	84.3	21 11.5	65.2	45 28.7	28.6	49 15.1	14.9	31 55.4	53.8	58 25.6	78.0
10	44 10.1	83.9	22 16.7	64.3	45 57.3	27.2	49 0.2	16.7	31 1.6	54.9	57 7.6	78.4
11	45 34.0	83.7	23 21.0	63.2	46 24.5	25.7	48 43.5	17.9	30 6.7	56.0	55 49.2	78.9
12	46 57.7	83.4	24 24.2	62.2	46 50.2	24.3	48 25.6	19.3	29 10.7	57.0	54 30.3	79.3
13	48 21.1	83.0	25 26.4	61.2	47 14.5	22.9	48 6.3	20.8	28 13.7	58.0	53 11.0	79.7
14	49 44.1	82.6	26 27.6	60.0	47 37.4	21.4	47 45.5	22.2	27 15.7	59.1	51 51.3	80.1
15	51 6.7	82.2	27 27.6	59.0	47 58.8	20.0	47 23.3	23.5	26 16.6	60.0	50 31.2	80.5
16	52 28.9	81.8	28 26.6	58.0	48 18.8	18.6	46 59.8	25.0	25 16.6	61.0	49 10.7	80.8
17	53 50.7	81.3	29 24.6	56.8	48 37.4	17.1	46 34.8	26.3	24 15.6	62.0	47 49.9	81.1
18	55 12.0	80.9	30 21.4	55.6	48 54.5	15.6	46 8.5	27.8	23 13.6	62.9	46 28.8	81.3
19	56 32.9	80.3	31 17.0	54.5	49 10.1	14.3	45 40.7	29.1	22 10.7	63.9	45 7.5	81.7
20	57 53.2	79.8	32 11.5	53.3	49 24.4	12.7	45 11.6	30.4	21 6.8	64.7	43 45.8	81.9
21	59 13.0	79.3	33 4.8	52.2	49 37.1	11.2	44 41.2	31.7	20 2.1	65.7	42 23.9	82.1
22	0 32.3	78.7	33 57.0	50.9	49 48.3	9.8	44 9.5	33.1	18 56.4	66.5	41 1.8	82.3
23	1 51.0	78.1	34 47.9	49.8	49 58.1	8.3	43 36.4	34.5	17 49.9	67.3	39 39.5	82.5
24	3 9.1	77.4	35 37.7	48.5	50 6.4	6.9	43 1.9	35.7	16 42.6	68.2	38 17.0	82.6
25	4 26.5	76.8	36 26.2	47.2	50 13.3	5.4	42 26.2	37.0	15 34.4	68.9	36 54.4	82.7
26	5 43.3	76.1	37 13.4	46.0	50 18.7	3.9	41 49.2	38.4	14 25.5	69.8	35 31.7	82.9
27	6 59.4	75.5	37 59.4	44.8	50 22.6	2.4	41 10.8	39.6	13 15.7	70.5	34 8.8	82.9
28	8 14.9	74.7	38 44.2	43.4	50 25.0	1.0	40 31.2	40.8	12 5.2	71.2	32 45.9	82.9
29	9 29.6	73.9	39 27.6	42.1	50 26.0	0.5	39 50.4	42.1	10 54.0	72.0	31 23.0	83.0
30	10 43.5		40 9.7		50 25.5		39 8.3		9 42.0		30 0.0	
	2°		2°		2°		2°		2°		1°	



*Equation of Moon's Centre.*

Argument.	Anomaly corrected.
<p>1. <math>\frac{1}{2} \pi</math> is the angle between the direction of the magnetic field and the direction of the electric field.</p> <p>2. The angle between the direction of the magnetic field and the direction of the electric field is <math>\frac{1}{2} \pi</math>.</p>	<p>1. <math>\frac{1}{2} \pi</math> is the angle between the direction of the magnetic field and the direction of the electric field.</p> <p>2. The angle between the direction of the magnetic field and the direction of the electric field is <math>\frac{1}{2} \pi</math>.</p>

O <sub>s</sub>		I <sub>s</sub>		II <sub>s</sub>		III <sub>s</sub>		IV <sub>s</sub>		V <sub>s</sub>	
7°	Diff for10	10°	Diff for10	12°	Diff for10	13°	Diff for10	12°	Diff for10	9°	Diff for10
0 0	0 0.0	20 57.9	59.2	38 43.6	30.1	17 35.2	4.8	16 20.8	35.2	58 28.9	55.0
30	3 32.6	70 9	58.9	40 14.0	29.6	17 20.9	5.4	14 35.3	35.6	55 43.8	55.3
1 0	7 5.2	70 9	58.5	41 42.7	29.0	17 4.8	5.9	12 48.5	36.0	52 58.0	55.5
30	10 37.8	70.8	58.1	43 9.6	28.4	16 47.1	6.5	11 0.4	36.4	50 11.6	55.7
2 0	14 10.3	70.8	57.7	44 34.9	27.8	16 27.6	7.0	9 11.1	36.9	47 24.5	55.9
30	17 42.7	70.8	57.3	45 58.4	27.3	16 6.5	7.6	7 20.5	37.3	44 36.8	56.1
3 0	21 15.0	70.8	57.0	47 20.2	26.7	15 43.7	8.2	5 28.7	37.7	41 48.5	56.3
30	24 47.3	70.7	56.5	48 40.3	26.1	15 19.2	8.7	3 35.6	38.1	38 59.5	56.5
4 0	28 19.4	70.6	56.1	49 58.7	25.5	14 53.1	9.3	1 41.3	38.5	36 10.0	56.7
30	31 51.2	70.6	55.7	51 15.3	25.0	14 25.2	9.8	59 45.8	38.9	33 19.8	56.9
5 0	35 23.0	70.5	55.3	52 30.2	24.4	13 55.8	10.4	57 49.1	39.3	30 29.1	57.1
30	38 54.5	70.4	54.9	53 43.3	23.8	13 24.7	10.9	55 51.1	39.7	27 37.8	57.3
6 0	42 25.8	70.4	54.5	54 54.7	23.2	12 51.9	11.5	53 52.0	40.1	24 45.9	57.5
30	45 56.9	70.3	54.0	56 4.4	22.6	12 17.4	12.0	51 51.7	40.5	21 53.5	57.6
7 0	49 27.7	70.2	53.6	57 12.3	22.1	11 41.4	12.6	49 50.3	40.9	19 0.6	57.8
30	52 58.2	70.1	53.2	58 18.5	21.5	11 3.7	13.1	47 47.6	41.3	16 7.1	58.0
8 0	56 28.5	70.0	52.7	59 22.9	20.9	10 24.3	13.6	45 43.8	41.7	13 13.1	58.2
30	59 58.4	69.9	52.3	0 25.6	20.3	9 43.4	14.2	43 38.9	42.0	10 18.6	58.3
9 0	3 28.0	69.7	51.8	1 26.5	19.7	9 0.8	14.7	41 32.8	42.4	7 23.6	58.5
30	6 57.2	69.6	51.4	2 25.7	19.1	8 16.6	15.3	39 25.6	42.8	4 28.1	58.6
10 0	10 26.0	69.5	50.9	3 23.0	18.6	7 30.8	15.8	37 17.3	43.1	1 32.2	58.8
30	13 54.5	69.3	50.5	4 18.7	17.9	6 43.4	16.3	35 7.9	43.5	58 35.8	59.0
11 0	17 22.5	69.2	50.0	5 12.5	17.4	5 54.4	16.8	32 57.4	43.9	55 38.9	59.1
30	20 50.1	69.1	49.6	6 4.6	16.8	5 3.9	17.4	30 45.8	44.2	52 41.7	59.3
12 0	24 17.3	68.9	49.1	6 54.9	16.2	4 11.7	17.9	28 33.1	44.6	49 43.9	59.4
30	27 44.0	68.7	48.6	7 43.5	15.6	3 18.0	18.4	26 19.4	44.9	46 45.8	59.5
13 0	31 10.2	68.5	48.1	8 30.3	15.0	2 22.7	19.0	24 4.6	45.3	43 47.3	59.6
30	34 35.8	68.4	47.7	9 15.4	14.4	1 25.8	19.5	21 48.8	45.6	40 48.4	59.8
14 0	38 1.0	68.2	47.2	9 58.6	13.8	0 27.4	20.0	19 31.9	45.9	37 49.1	59.9
30	41 25.6	68.0	46.6	10 40.1	13.3	59 27.4	20.5	17 14.1	46.3	34 49.5	60.0
15 0	44 49.6	40 49.3		11 19.9		58 25.9		14 55.2		31 49 4	
8°		11°		13°		12°		11°		8°	

*Equation of Moon's Centre.*

Argument.      Anomaly corrected.

VI <sup>s</sup>		VII <sup>s</sup>		VIII <sup>s</sup>		IX <sup>s</sup>		X <sup>s</sup>		XI <sup>s</sup>		
	7°	Diff for 10	4°	Diff for 10	1°	Diff for 10	0°	Diff for 10	1°	Diff for 10	3°	Diff for 10
°	°	°	°	°	°	°	°	°	°	°	°	°
0 0	0 0.0	61.8	131.1	54.8	43 39.2	34.7	42 24.8	4.2	21 16.4	30.7	39 2.1	59.6
30	56 54.6	61.8	58 46.7	54.6	41 55.0	34.3	42 12.1	3.6	22 48.5	31.2	42 0.8	60.0
1 0	53 49.2	61.8	56 3.0	54.3	40 12.0	33.8	42 1.2	3.1	24 22.2	31.8	45 0.7	60.3
30	50 43.9	61.8	53 20.0	54.1	38 30.5	33.4	41 52.0	2.5	25 57.7	32.4	48 1.7	60.7
2 0	47 38.6	61.7	50 37.7	53.8	36 50.3	33.0	41 44.4	1.9	27 34.8	33.0	51 3.7	61.0
30	44 33.4	61.8	47 56.2	53.6	35 11.3	32.5	41 38.7	1.4	29 13.7	33.5	54 6.7	61.3
3 0	41 28.1	61.7	45 15.4	53.4	33 33.7	32.1	41 34.6	0.8	30 54.2	34.0	57 10.7	61.7
30	38 23.0	61.7	42 35.3	53.1	31 57.5	31.6	41 32.2	0.2	32 36.3	34.6	0 15.8	62.0
4 0	35 18.0	61.7	39 56.0	52.9	30 22.6	31.2	41 31.6	0.4	34 20.2	35.1	3 21.8	62.3
30	32 13.0	61.6	37 17.4	52.6	28 49.0	30.7	41 32.7	1.0	36 5.6	35.7	6 28.8	62.7
5 0	29 8.1	61.6	34 39.6	52.3	27 16.8	30.2	41 35.6	1.5	37 52.8	36.2	9 36.8	63.0
30	26 3.4	61.5	32 2.7	52.1	25 46.1	29.8	41 40.1	2.1	39 41.5	36.8	12 45.7	63.3
6 0	22 58.8	61.5	29 26.5	51.8	24 16.7	29.3	41 46.4	2.7	41 32.0	37.3	15 55.5	63.6
30	19 54.3	61.4	26 51.1	51.5	22 48.7	28.9	41 54.5	3.3	43 24.0	37.9	19 6.2	63.9
7 0	16 50.0	61.4	24 16.6	51.2	21 22.1	28.4	42 4.3	3.9	45 17.7	38.4	22 17.8	64.2
30	13 45.8	61.3	21 42.9	51.0	19 56.9	27.9	42 15.9	4.4	47 12.9	39.0	25 30.3	64.5
8 0	10 41.9	61.3	19 10.0	50.7	18 33.1	27.4	42 29.2	5.0	49 9.8	39.5	28 43.7	64.7
30	7 38.0	61.2	16 38.0	50.4	17 10.8	27.0	42 44.2	5.6	51 8.3	40.0	31 57.8	65.0
9 0	4 34.4	61.1	14 6.9	50.1	15 49.8	26.5	43 1.1	6.2	53 8.4	40.6	35 12.9	65.3
30	1 31.0	61.1	11 36.6	49.8	14 30.4	26.0	43 19.6	6.8	55 10.1	41.1	38 28.7	65.5
10 0	58 27.8	61.0	9 7.3	49.5	13 12.5	25.5	43 39.9	7.4	57 13.3	41.6	41 45.2	65.8
30	55 24.9	60.9	6 38.9	49.2	11 55.9	25.0	44 2.0	8.0	59 18.2	42.1	45 2.6	66.0
11 0	52 22.2	60.8	4 11.3	48.9	10 40.9	24.5	44 25.9	8.5	1 24.5	42.6	48 20.7	66.3
30	49 19.7	60.7	1 44.7	48.6	9 27.3	24.0	44 51.5	9.1	3 32.4	43.2	51 39.6	66.5
12 0	46 17.5	60.6	59 18.9	48.2	8 15.2	23.5	45 18.8	9.7	5 41.9	43.7	54 59.1	66.7
30	43 15.6	60.5	56 54.2	47.9	7 4.6	23.1	45 48.0	10.3	7 52.9	44.2	58 19.3	67.0
13 0	40 14.0	60.5	54 30.4	47.6	5 55.4	22.5	46 18.9	10.9	10 5.5	44.7	1 40.3	67.2
30	37 12.6	60.3	52 7.5	47.3	4 47.8	22.0	46 51.5	11.5	12 19.5	45.2	5 1.9	67.4
14 0	34 11.6	60.2	49 45.6	47.0	3 41.7	21.5	47 26.0	12.1	14 35.1	45.7	8 24.1	67.6
30	31 10.9	60.1	47 24.7	46.6	2 37.1	21.0	48 2.2	12.6	16 52.1	46.2	11 46.9	67.8
15 0	28 10.6		45 4.8		1 34.1		48 40.1		19 10.7		15 10.4	
	5°		2°		1°		0°		2°		5°	



*Equation of Moon's Centre.*

Argument. Anomaly corrected.

Os		Is		IIs		IIIs		IVs		Vs		
	8°	Diff for 10	11°	Diff for 10	13°	Diff for 10	12°	Diff for 10	11°	Diff for 10	8°	Diff for 10
15 0	44 49.6	67.8	40 49.3	46.2	11 19.9	12.6	58 25.9	21.0	14 55.2	46.6	31 49.4	60.1
30	48 13.1	67.6	43 7.9	45.7	11 57.8	12.1	57 22.9	21.5	12 35.3	47.0	28 49.1	60.2
16 0	51 35.9	67.4	45 24.9	45.2	12 34.0	11.5	56 18.3	22.0	10 14.4	47.3	25 48.4	60.3
30	54 58.1	67.2	47 40.5	44.7	13 8.5	10.9	55 12.2	22.5	7 52.5	47.7	22 47.4	60.5
17 0	58 19.7	67.0	49 54.5	44.2	13 41.1	10.3	54 4.6	23.1	5 29.6	47.9	19 46.0	60.5
30	1 40.7	66.7	52 7.1	43.7	14 12.0	9.7	52 55.4	23.5	3 5.8	48.3	16 44.4	60.6
18 0	5 0.9	66.5	54 18.1	43.2	14 41.2	9.1	51 44.8	24.0	0 41.1	48.6	13 42.5	60.7
30	8 20.4	66.3	56 27.6	42.6	15 8.5	8.5	50 32.7	24.5	58 15.3	48.9	10 40.3	60.8
19 0	11 39.3	66.0	58 35.5	42.1	15 34.1	8.0	49 19.1	25.0	55 48.7	49.2	7 37.8	60.9
30	14 57.4	65.8	0 41.8	41.6	15 58.0	7.4	48 4.1	25.5	53 21.1	49.5	4 35.1	61.0
20 0	18 14.8	65.5	2 46.7	41.1	16 20.1	6.8	46 47.5	26.0	50 52.7	49.8	1 32.2	61.1
30	21 31.3	65.3	4 49.9	40.6	16 40.4	6.2	45 29.6	26.5	48 23.4	50.1	58 29.0	61.1
21 0	24 47.1	65.0	6 51.6	40.0	16 58.9	5.6	44 10.2	27.0	45 53.1	50.4	55 25.6	61.2
30	28 2.2	64.7	8 51.7	39.5	17 15.8	5.0	42 49.2	27.4	43 22.0	50.7	52 22.0	61.3
22 0	31 16.3	64.5	10 50.2	39.0	17 30.8	4.4	41 26.9	27.9	40 50.0	51.0	49 18.1	61.3
30	34 29.7	64.2	12 47.1	38.4	17 44.1	3.9	40 3.1	28.4	38 17.1	51.2	46 14.2	61.4
23 0	37 42.2	63.9	14 42.3	37.9	17 55.7	3.3	38 37.9	28.9	35 43.4	51.5	43 10.0	61.4
30	40 53.8	63.6	16 36.0	37.3	18 5.5	2.7	37 11.3	29.3	33 8.9	51.8	40 5.7	61.5
24 0	44 4.5	63.2	18 28.0	36.8	18 13.6	2.1	35 43.3	29.8	30 33.5	52.1	37 1.2	61.5
30	47 14.3	63.0	20 18.5	36.2	18 19.9	1.5	34 13.9	30.2	27 57.3	52.3	33 56.6	61.6
25 0	50 23.2	62.7	22 7.2	35.7	18 24.4	1.0	32 43.2	30.7	25 20.4	52.6	30 51.9	61.6
30	53 31.2	62.3	23 54.4	35.1	18 27.3	0.4	31 11.0	31.2	22 42.6	52.9	27 47.0	61.7
26 0	56 38.2	62.0	25 39.8	34.6	18 28.4	0.2	29 37.4	31.6	20 4.0	53.1	24 42.0	61.7
30	59 44.2	61.7	27 23.7	34.1	18 27.8	0.8	28 2.5	32.1	17 24.7	3.3	21 37.0	61.7
27 0	2 49.3	61.3	29 5.8	33.5	18 25.4	1.4	26 26.3	32.5	14 44.7	53.6	18 31.8	61.7
30	5 53.3	61.0	30 46.3	33.0	18 21.3	1.9	24 48.7	33.0	12 3.8	53.8	15 26.6	61.7
28 0	8 56.3	60.7	32 25.2	32.4	18 15.6	2.5	23 9.7	33.4	9 22.3	54.1	12 21.4	61.8
30	11 58.3	60.3	34 2.3	31.8	18 8.0	3.1	21 29.5	33.8	6 40.0	54.3	9 16.1	61.8
29 0	14 59.3	60.0	35 37.8	31.2	17 58.8	3.6	19 48.0	34.3	3 57.0	54.6	6 10.8	61.8
30	17 59.2	59.6	37 11.5	30.7	17 47.9	4.2	18 5.0	34.7	1 13.3	54.8	3 5.4	61.8
30 0	20 57.9		38 43.6		17 35.2		16 20.8		58 28.9		0 0.0	
	10°		12°		13°		12°		9°		7°	

*Equation of Moon's Centre.*

Argument.      Anomaly corrected.

VI <sub>s</sub>		VII <sub>s</sub>		VIII <sub>s</sub>		IX <sub>s</sub>		X <sub>s</sub>		XI <sub>s</sub>	
5°	Diff for 10	2°	Diff for 10	1°	Diff for 10	0°	Diff for 10	2°	Diff for 10	5°	Diff for 10
° "	" "	" "	" "	" "	" "	" "	" "	" "	" "	" "	" "
15 0	28 10.6	45 4.8	46.3	1 34.1	20.5	48 40.1	13.3	19 10.7	46.6	15 10.4	68.0
30	25 10.5	42 45.9	45.9	0 32.6	20.0	49 19.9	13.8	21 30.6	47.2	8 34.4	68.2
16 0	22 10.9	40 28.1	45.6	59 32.6	19.5	50 1.4	14.4	23 52.1	47.7	21 59.0	68.4
30	19 11.6	38 11.2	45.3	58 34.2	19.0	50 44.6	15.0	26 15.1	48.1	25 24.2	68.5
17 0	16 12.7	35 55.4	44.9	57 37.3	18.4	51 29.7	15.6	28 39.5	48.6	28 49.8	68.7
30	13 14.2	33 40.6	44.6	56 42.0	17.9	52 16.5	16.2	31 5.3	49.1	32 16.0	68.9
18 0	10 16.1	31 26.9	44.2	55 48.3	17.4	53 5.1	16.8	33 32.5	49.6	35 42.7	69.1
30	7 18.3	29 14.2	43.9	54 56.1	16.8	53 55.4	17.4	36 1.2	50.0	39 9.9	69.2
19 0	4 21.1	27 2.6	43.5	54 5.6	16.3	54 47.5	17.9	38 31.2	50.5	42 37.5	69.3
30	1 24.2	24 52.1	43.1	53 16.6	15.8	55 41.3	18.6	41 2.7	50.9	46 5.5	69.5
20 0	58 27.8	22 42.7	42.8	52 29.2	15.3	56 37.0	19.1	43 35.5	51.4	49 34.0	69.6
30	55 31.9	20 34.4	42.4	51 43.4	14.7	57 34.3	19.7	46 9.7	51.8	53 2.8	69.7
21 0	52 36.4	18 27.2	42.0	50 59.2	14.2	58 33.5	20.3	48 45.2	52.3	56 31.9	69.9
30	49 41.4	16 21.1	41.6	50 16.6	13.6	59 34.4	20.9	51 22.1	52.7	0 1.6	70.0
22 0	46 46.9	14 16.2	41.3	49 35.7	13.1	0 37.1	21.5	54 0.3	53.2	3 31.5	70.1
30	43 52.9	12 12.4	40.9	48 56.3	12.6	1 41.5	22.1	56 39.9	53.6	7 1.8	70.2
23 0	40 59.4	10 9.7	40.5	48 18.6	12.0	2 47.7	22.6	59 20.7	54.0	10 32.3	70.3
30	38 6.5	8 8.3	40.1	47 42.6	11.5	3 55.6	23.2	2 2.8	54.5	14 3.1	70.4
24 0	35 14.1	6 8.0	39.7	47 8.1	10.9	5 5.3	23.8	4 46.2	54.9	17 34.2	70.4
30	32 22.2	4 8.9	39.3	46 35.3	10.4	6 16.7	24.4	7 30.9	55.3	21 5.5	70.5
25 0	29 30.9	2 10.9	38.9	46 4.2	9.8	7 29.8	25.0	10 16.8	55.7	24 37.0	70.6
30	26 40.2	0 14.2	38.5	45 34.8	9.3	8 44.7	25.5	13 4.0	56.1	28 8.8	70.6
26 0	23 50.0	58 18.7	38.1	45 6.9	8.7	10 1.3	26.1	15 52.4	56.5	31 40.7	70.7
30	21 0.5	55 24.4	37.7	44 40.8	8.2	11 19.7	26.7	18 42.0	57.0	35 12.8	70.7
27 0	18 11.5	54 31.3	37.3	44 16.3	7.6	12 39.8	27.3	21 32.9	57.3	38 45.1	70.8
30	15 23.2	52 39.5	36.9	43 53.5	7.0	14 1.6	27.8	24 24.8	57.7	42 17.3	70.8
28 0	12 35.5	50 48.9	36.4	43 32.4	6.5	15 25.1	28.4	27 18.0	58.1	45 49.7	70.8
30	9 48.4	48 59.6	36.0	43 12.9	5.9	16 50.4	29.0	30 12.3	58.5	49 22.2	70.9
29 0	7 2.0	47 11.5	35.6	42 55.2	5.4	18 17.3	29.6	33 7.8	58.9	52 54.8	70.9
30	4 16.2	45 24.7	35.2	42 39.1	4.8	19 46.0	30.1	36 4.4	59.2	56 27.4	70.9
30 0	1 31.1	43 39.2		42 24.8		21 16.4		39 2.1		0 0.0	
4°		1°		0°		1°		3°		7°	

*Variation.*

Argument. Variation, corrected.

	O <sup>s</sup>		I <sup>s</sup>		II <sup>s</sup>		III <sup>s</sup>		IV <sup>s</sup>		V <sup>s</sup>	
Diff.	0°	Diff.	1°	Diff.	1°	Diff.	0°	Diff.	0°	Diff.	0°	Diff.
	' "		' "		' "		' "		' "		' "	
0	38 0.0	"	8 1.5	"	6 57.9	"	35 54.4	"	5 29.5	"	6 1.6	"
1	39 13.3	73.3	8 35.5	34.0	6 18.0	39.9	34 40.4	74.0	4 54.2	35.3	6 41.6	40.0
2	40 26.5	73.3	9 7.2	31.7	5 35.9	42.1	33 26.6	73.8	4 21.3	32.9	7 23.9	42.3
3	41 39.5	73.0	9 36.5	29.3	4 51.7	44.2	32 13.0	73.6	3 50.6	30.7	8 8.4	44.5
4	42 52.2	72.7	10 3.4	26.9	4 5.5	46.2	30 59.6	73.4	3 22.3	28.3	8 55.0	46.6
5	44 4.5	72.3	10 27.9	24.5	3 17.3	48.2	29 46.7	72.9	2 56.5	25.8	9 43.7	48.7
		71.9		22.0		50.1		72.4		23.4		50.8
6	45 16.4	71.3	10 49.9	19.5	2 27.2	51.9	28 34.3	71.9	2 33.1	21.0	10 34.5	52.8
7	46 27.7	70.7	11 9.4	17.0	1 35.3	53.7	27 22.4	71.2	2 12.1	18.4	11 27.3	54.7
8	47 38.4	69.9	11 26.4	14.5	0 41.6	55.5	26 11.2	70.5	1 53.7	15.9	12 22.0	56.6
9	48 48.3	69.1	11 40.9	12.0	59 46.1	57.1	25 0.7	69.6	1 37.8	13.3	13 18.6	58.3
10	49 57.4	68.2	11 52.9	9.3	58 49.0	58.8	23 51.1	68.8	1 24.5	10.8	14 16.9	60.1
11	51 5.6	67.2	12 2.2	6.8	57 50.2	60.2	22 42.3	67.8	1 13.7	8.2	15 17.0	61.7
12	52 12.8	66.1	12 9.0	4.2	56 50.0	61.7	21 34.5	66.6	1 5.5	5.5	16 18.7	63.3
13	53 18.9	64.9	12 13.2	1.6	55 48.3	63.1	20 27.9	65.6	1 0.0	3.0	17 22.0	64.9
14	54 23.8	63.7	12 14.8	0.9	54 45.2	64.3	19 22.3	64.3	0 57.0	0.3	18 26.9	66.2
15	55 27.5	62.3	12 13.9	3.6	53 40.9	65.6	18 18.0	63.0	0 56.7	2.3	19 33.1	67.6
16	56 29.8	60.9	12 10.3	6.1	52 35.3	66.8	17 15.0	61.6	0 59.0	4.9	20 40.7	68.9
17	57 30.7	59.4	12 4.2	8.7	51 28.5	67.8	16 13.4	60.2	1 3.9	7.6	21 49.6	70.0
18	58 30.1	57.9	11 55.5	11.3	50 20.7	68.8	15 13.2	58.6	1 11.5	10.1	22 59.6	71.2
19	59 28.0	56.2	11 44.2	13.7	49 11.9	69.7	14 14.6	57.1	1 21.6	12.8	24 10.8	72.1
20	0 24.2	54.5	11 30.5	16.4	48 2.2	70.5	13 17.5	55.3	1 34.4	15.4	25 22.9	73.0
21	1 18.7	52.7	11 14.1	18.8	46 51.7	71.2	12 22.2	53.7	1 49.8	18.0	26 35.9	73.9
22	2 11.4	50.9	10 55.3	21.3	45 40.5	71.9	11 28.5	51.8	2 7.8	20.5	27 49.8	74.7
23	3 2.3	48.9	10 34.0	23.8	44 28.6	72.5	10 36.7	49.9	2 28.3	23.1	29 4.5	75.2
24	3 51.2	47.0	10 10.2	26.2	43 16.1	72.9	9 46.8	48.0	2 51.4	25.5	30 19.7	75.9
25	4 38.2	44.9	9 44.0	28.6	42 3.2	73.3	8 58.8	46.1	3 16.9	28.1	31 35.6	76.3
26	5 23.1	42.9	9 15.4	30.9	40 49.9	73.7	8 12.7	44.0	3 45.0	30.6	32 51.9	76.7
27	6 6.0	40.7	8 44.5	33.3	39 36.2	73.8	7 28.7	41.9	4 15.6	32.9	34 8.6	77.0
28	6 46.7	38.5	8 11.2	35.5	38 22.4	74.0	6 46.8	39.7	4 48.5	35.4	35 25.6	77.1
29	7 25.2	36.3	7 35.7	37.8	37 8.4	74.0	6 7.1	37.6	5 23.9	37.7	36 42.7	77.3
30	8 1.5		6 57.9		35 54.4		5 29.5		6 1.6		38 0.0	
	1°		1°		0°		0°		0°		0°	

*Variation.*

Argument. Variation corrected.

	VI <sup>s</sup>		VII <sup>s</sup>		VIII <sup>s</sup>		IX <sup>s</sup>		X <sup>s</sup>		XI <sup>s</sup>	
Deci	0°	Diff.	1°	Diff.	1°	Diff.	0°	Diff.	0°	Diff.	0°	Diff.
	' "	"	' "	"	' "	"	' "	"	' "	"	' "	"
0	38 0.0	77.3	9 58.4	37.7	10 30.5	37.6	40 5.6	74.0	9 2.1	37.8	7 58.5	36.3
1	39 17.3	77.1	10 36.1	35.4	9 52.9	39.7	38 51.6	74.0	8 24.3	35.5	8 34.8	38.5
2	40 34.4	77.0	11 11.5	32.9	9 13.2	37.6	37 37.6	73.8	7 48.8	33.3	9 13.3	40.7
3	41 51.4	76.7	11 44.4	30.6	8 31.3	41.9	36 23.8	73.7	7 15.5	30.9	9 54.0	42.9
4	43 8.1	76.3	12 15.0	28.1	7 47.3	44.0	35 10.1	73.3	6 44.6	28.6	10 36.9	44.9
5	44 24.4	75.9	12 43.1	25.5	7 1.2	46.1	33 56.8	72.9	6 16.0	26.2	11 21.8	47.0
6	45 40.3	75.2	13 8.6	23.1	6 13.2	49.9	32 43.9	72.5	5 49.8	23.8	12 8.8	48.9
7	46 55.5	74.7	13 31.7	20.5	5 23.3	51.8	31 31.4	71.9	5 26.0	21.3	12 57.7	50.9
8	48 10.2	73.9	13 52.2	18.0	4 31.5	53.7	30 19.5	71.2	5 4.7	18.8	13 48.6	52.7
9	49 24.1	73.0	14 10.2	15.4	3 37.8	55.3	29 8.3	70.5	4 45.9	16.4	14 41.3	54.5
10	50 37.1	72.1	14 25.6	12.8	2 42.5	57.1	27 57.8	69.7	4 29.5	13.7	15 35.8	56.2
11	51 49.2	71.2	14 38.4	10.1	1 45.4	58.6	26 48.1	68.8	4 15.8	11.3	16 32.0	57.9
12	53 0.4	70.0	14 48.5	7.6	0 46.8	60.2	25 39.3	67.8	4 4.5	8.7	17 29.9	59.4
13	54 10.4	68.9	14 56.1	4.9	59 46.6	61.6	24 31.5	66.8	3 55.8	6.1	18 29.3	60.9
14	55 19.3	67.6	15 1.0	2.3	58 45.0	63.0	23 24.7	65.6	3 49.7	3.6	19 30.2	62.3
15	56 26.9	66.2	15 3.3	0.3	57 42.0	64.3	22 19.1	64.3	3 46.1	0.9	20 32.5	63.7
16	57 33.1	64.9	15 3.0	3.0	56 37.7	65.6	21 14.8	63.1	3 45.2	1.6	21 36.2	64.9
17	58 38.0	63.3	15 0.0	5.5	55 32.1	66.6	20 11.7	61.7	3 46.8	4.2	22 41.1	66.1
18	59 41.3	61.7	14 54.5	8.2	54 25.5	67.8	19 10.0	60.2	3 51.0	6.8	23 47.2	67.2
19	0 43.0	60.1	14 46.3	10.8	53 17.7	68.8	18 9.8	58.8	3 57.8	9.3	24 54.4	68.2
20	1 43.1	58.3	14 35.5	13.3	52 8.9	69.6	17 11.0	57.1	4 7.1	12.0	26 2.6	69.1
21	2 41.4	56.6	14 22.2	15.9	50 59.3	70.5	16 13.9	55.5	4 19.1	14.5	27 11.7	69.9
22	3 38.0	54.7	14 6.3	18.4	49 48.8	71.2	15 18.4	53.7	4 33.6	17.0	28 21.6	70.7
23	4 32.7	52.8	13 47.9	21.0	48 37.6	71.9	14 24.7	51.9	4 50.6	19.5	29 32.3	71.3
24	5 25.5	50.8	13 26.9	23.4	47 25.7	72.4	13 32.8	50.1	5 10.1	22.0	30 43.6	71.9
25	6 16.3	48.7	13 3.5	25.8	46 13.3	72.9	12 42.7	48.2	5 32.1	24.5	31 55.5	72.3
26	7 5.0	46.6	12 37.7	28.3	45 0.4	73.4	11 54.5	46.2	5 56.6	26.9	33 7.8	72.7
27	7 51.6	44.5	12 9.4	30.7	43 47.0	73.6	11 8.3	44.2	6 23.5	29.3	34 20.5	73.0
28	8 36.1	42.3	11 38.7	32.9	42 33.4	73.8	10 24.1	42.1	6 52.8	31.7	35 33.5	73.2
29	9 18.4	40.0	11 5.8	35.3	41 19.6	74.0	9 42.0	39.9	7 24.5	34.0	36 46.7	73.3
30	9 58.4		10 30.5		40 5.6		9 2.1		7 58.5		38 0.0	
	1°		1°		0°		0°		0°		0°	

Argument. Supplement of Node+Moon's Orbit Longitude.

	Os	Vls	Diff.	Is	Vlls	Diff.	Ilss	Vlls	Diff.	Ills	IXs	Diff.	IVs	Xs	Diff.	Vs	Xls	Diff.
0	7	0.0		1	3.0		1	3.0		7	0.0		12	57.0		12	57.0	
1	6	45.6	14.4	0	56.0	7.0	1	10.4	7.4	7	14.4	14.4	13	4.0	7.0	12	49.6	7.4
2	6	31.2	14.4	0	49.5	6.5	1	18.3	7.9	7	28.8	14.4	13	10.5	6.5	12	41.7	7.9
3	6	16.9	14.3	0	43.4	6.1	1	26.5	8.2	7	43.1	14.3	13	16.6	6.1	12	33.5	8.2
4	6	2.6	14.3	0	37.8	5.6	1	35.2	8.7	7	57.4	14.3	13	22.2	5.6	12	24.8	8.7
5	5	48.4	14.2	0	32.7	5.1	1	44.2	9.0	8	11.6	14.2	13	27.3	5.1	12	15.8	9.0
			14.1			4.5			9.5			14.1			4.5			9.5
6	5	34.3	14.0	0	28.2	4.3	1	53.7	9.8	8	25.7	14.0	13	31.8	4.3	12	6.3	9.8
7	5	20.3	13.9	0	23.9	3.9	2	3.5	10.2	8	39.7	13.9	13	36.1	3.9	11	56.5	10.2
8	5	6.4	13.8	0	20.0	3.2	2	13.7	10.5	8	53.6	13.8	13	40.0	3.2	11	46.3	10.5
9	4	52.6	13.6	0	16.8	2.7	2	24.2	10.8	9	7.4	13.6	13	43.2	2.7	11	35.8	10.8
10	4	39.0	13.4	0	14.1	2.3	2	35.0	11.2	9	21.0	13.4	13	45.9	2.3	11	25.0	11.2
			13.3			1.7			11.5			13.3			1.7			11.5
11	4	25.6	13.0	0	10.1	1.3	2	57.7	11.8	9	47.7	13.0	13	49.9	1.3	11	2.3	11.8
12	4	12.3	12.8	0	8.8	0.7	3	9.5	12.1	10	0.7	12.8	13	51.2	0.7	10	50.5	12.1
13	3	59.3	12.6	0	8.1	0.3	3	21.6	12.3	10	13.5	12.6	13	51.9	0.3	10	38.4	12.3
14	3	46.5	12.3	0	7.8	0.3	3	33.9	12.6	10	26.1	12.3	13	52.2	0.3	10	26.1	12.6
15	3	33.9	12.1	0	8.1	0.7	3	46.5	12.8	10	33.4	12.1	13	51.9	0.7	10	13.5	12.8
16	3	21.6	11.8	0	8.8	1.3	3	59.3	13.0	10	50.5	11.8	13	51.2	1.3	10	0.7	13.0
17	3	9.5	11.5	0	10.1	1.7	4	12.3	13.3	11	2.3	11.5	13	49.9	1.7	9	47.7	13.3
18	2	57.7	11.2	0	11.8	2.3	4	25.6	13.4	11	13.8	11.2	13	48.2	2.3	9	34.4	13.4
19	2	46.2	10.8	0	14.1	2.7	4	39.0	13.6	11	25.0	10.8	13	45.9	2.7	9	21.0	13.6
20	2	35.0	10.5	0	16.8	3.2	4	52.6	13.8	11	35.8	10.5	13	43.2	3.2	9	7.4	13.8
21	2	21.2	10.2	0	20.0	3.9	5	6.4	13.9	11	46.3	10.2	13	40.0	3.9	8	53.6	13.9
22	2	13.7	9.8	0	23.9	4.3	5	20.3	14.0	11	56.5	9.8	13	36.1	4.3	8	39.7	14.0
23	2	3.5	9.5	0	28.2	4.5	5	34.3	14.1	12	6.3	9.5	13	31.8	4.5	8	25.7	14.1
24	1	53.7	9.0	0	32.7	5.1	5	48.4	14.2	12	15.8	9.0	13	27.3	5.1	8	11.6	14.2
25	1	44.2	8.7	0	37.8	5.6	6	2.6	14.3	12	24.8	8.7	13	22.2	5.6	7	57.4	14.3
26	1	35.2	8.2	0	43.4	6.1	6	16.9	14.3	12	33.5	8.2	13	16.6	6.1	7	43.1	14.3
27	1	26.5	7.9	0	49.5	6.5	6	31.2	14.4	12	41.7	7.9	13	10.5	6.5	7	28.8	14.4
28	1	18.3	7.4	0	56.0	7.0	6	45.6	14.4	12	49.6	7.4	13	4.0	7.0	7	14.4	14.4
29	1	10.4		1	3.0		7	0.0		12	57.0		12	57.0		7	0.0	
30	1	3.0																

TABLE LIV. *Lunar Nutation in Longitude.*

Argument. Supplement of the Node.

	Os	Is	II <sup>s</sup>	III <sup>s</sup>	IV <sup>s</sup>	V <sup>s</sup>	
	+	+	+	+	+	+	
0	0.0	8.5	14.8	17.3	15.2	8.8	30
2	0.6	9.0	15.1	17.2	14.9	8.1	28
4	1.2	9.4	15.4	17.2	14.5	7.7	26
6	1.7	10.0	15.6	17.2	14.2	7.2	24
8	2.3	10.4	15.9	17.2	13.8	6.5	22
10	2.9	10.9	16.4	17.1	13.5	6.1	20
12	3.5	11.4	16.3	17.0	13.0	5.4	18
14	4.1	11.8	16.5	16.9	12.6	4.8	16
16	4.6	12.2	16.7	16.7	12.2	4.3	14
18	5.2	12.6	16.8	16.5	11.8	3.7	12
20	5.8	13.1	16.9	16.4	11.3	3.0	10
22	6.2	13.4	17.1	16.2	10.9	2.4	8
24	6.9	13.8	17.1	15.9	10.4	1.8	6
26	7.4	14.1	17.2	15.7	9.8	1.3	4
28	7.8	14.5	17.2	15.4	9.4	0.6	2
30	8.5	14.8	17.3	15.2	8.8	0.0	0
	—	—	—	—	—	—	
	XI <sup>s</sup>	X <sup>s</sup>	IX <sup>s</sup>	VIII <sup>s</sup>	VII <sup>s</sup>	VI <sup>s</sup>	



*Moon's Distance from the North Pole of the Ecliptic.*

Argument. Supplement of Node+Moon's Orbit Longitude.

	III <sup>s</sup>		IV <sup>s</sup>		V <sup>s</sup>		VI <sup>s</sup>		VII <sup>s</sup>		VIII <sup>s</sup>	
	84°	85°	Diff. for 10	87°	Diff. for 10	89°	Diff. for 10	92°	Diff. for 10	94°		
0 0	39 16.0	20 42.7	27.2	13 46.6	46.8	48 0.0	53.8	22 13.4	46.6	15 17.3	30 0	
30	39 16.7	22 4.2	27.6	16 6.9	47.0	50 41.4	53.8	24 33.1	46.4	16 37.7	36	
1 0	39 18.8	23 27.0	28.0	18 27.8	47.2	53 22.9	53.8	26 52.2	46.0	17 56.8	29 0	
30	39 22.4	24 51.0	28.4	20 49.5	47.4	56 4.3	53.8	29 10.2	45.8	19 14.6	30	
2 0	39 27.3	26 16.2	28.8	23 11.8	47.7	58 45.7	53.8	31 27.5	45.6	20 31.3	28 0	
30	39 33.7	27 42.6	29.2	25 34.8	47.9	1 27.0	53.8	33 44.2	45.3	21 46.7	30	
3 0	39 41.5	29 10.1	29.6	27 58.5	48.1	4 8.3	53.7	36 0.2	45.0	23 0.8	27 0	
30	39 50.6	30 38.9	30.0	30 22.8	48.3	6 49.5	53.7	38 15.3	44.8	24 13.7	30	
4 0	40 1.2	32 8.8	30.4	32 47.7	48.5	9 30.6	53.7	40 29.7	44.5	25 25.3	26 0	
30	40 13.2	33 39.9	30.8	35 13.2	48.7	12 11.6	53.6	42 43.3	44.3	26 35.7	30	
5 0	40 26.7	35 12.2	31.1	37 39.3	48.9	14 52.5	53.6	44 56.2	44.0	27 44.8	25 0	
30	40 41.5	36 45.6	31.5	40 6.1	49.1	17 33.3	53.6	47 8.1	43.8	28 52.6	30	
6 0	40 57.7	38 20.1	31.9	42 33.4	49.3	20 14.0	53.5	49 19.4	43.4	29 59.0	24 0	
30	41 15.4	39 55.8	32.3	45 1.2	49.5	22 54.4	53.5	51 29.7	43.2	31 4.3	30	
7 0	41 34.4	41 32.7	32.6	47 29.6	49.7	25 34.8	53.4	53 39.3	42.9	32 8.2	23 0	
30	41 54.8	43 10.6	33.0	49 58.6	49.8	28 14.9	53.3	55 48.0	42.6	33 10.9	30	
8 0	42 16.7	44 49.7	33.4	52 28.1	50.0	30 54.9	53.3	57 55.8	42.3	34 12.2	22 0	
30	42 39.9	46 29.9	33.8	54 58.2	50.2	33 34.7	53.2	0 2.8	42.0	35 12.2	30	
9 0	43 4.6	48 11.2	34.1	57 28.7	50.4	36 14.3	53.1	2 8.9	41.7	36 10.9	21 0	
30	43 30.6	49 53.5	34.5	59 59.8	50.5	38 53.7	53.0	4 14.1	41.5	37 8.3	30	
10 0	43 58.1	51 37.0	34.9	2 31.3	50.7	41 32.8	53.0	6 18.4	41.1	38 4.4	20 0	
30	44 26.9	53 21.6	35.2	5 3.3	50.8	44 11.7	52.9	8 21.8	40.8	38 59.1	30	
11 0	44 57.1	55 7.1	35.7	7 35.8	51.0	46 50.4	52.8	10 24.3	40.5	39 52.5	19 0	
30	45 28.8	56 53.8	35.9	10 8.8	51.1	49 28.7	52.7	12 25.9	40.2	40 44.6	30	
12 0	46 1.8	58 41.6	36.2	12 42.1	51.3	52 6.8	52.6	14 26.6	39.9	41 35.3	18 0	
30	46 36.1	0 30.3	36.6	15 16.0	51.4	54 44.6	52.5	16 26.3	39.6	42 24.7	30	
13 0	47 11.9	2 20.1	37.0	17 50.2	51.6	57 22.1	52.4	18 25.0	39.3	43 12.7	17 0	
30	47 49.0	4 11.0	37.3	20 24.9	51.7	59 59.3	52.3	20 22.8	38.0	43 59.4	30	
14 0	48 27.5	6 2.9	37.6	22 59.9	51.8	2 36.2	52.2	22 19.7	38.6	44 44.7	16 0	
30	49 7.4	7 55.7	38.0	25 35.3	51.9	5 12.7	52.1	24 15.5	38.3	45 28.7	30	
15 0	49 48.7	9 49.6		28 11.1		7 48.9		26 10.4		46 11.3	15 0	
	84°	86°		88°		91°		93°		94°		
	II <sup>s</sup>	I <sup>s</sup>		Os		XI <sup>s</sup>		X <sup>s</sup>		IX <sup>s</sup>		

*Moon's Distance from the North Pole of the Ecliptic.*

Argument. Supplement of Node+Moon's Orbit Longitude.

	III <sup>s</sup>	IV <sup>s</sup>		V <sup>s</sup>		VI <sup>s</sup>		VII <sup>s</sup>		VIII <sup>s</sup>	
	84°	86°	Diff. for 10	88°	Diff. for 10	91°	Diff. for 10	93°	Diff. for 10	94°	
° ' "	° ' "	° ' "		° ' "		° ' "		° ' "		° ' "	° ' "
15 0 49 48.7	9 49.6	38.3	28 11.1	52.1	7 48.9	51.9	26 10.4	38.0	46 11.3	15 0	
30 50 31.3	11 44.5	38.6	30 47.3	52.2	10 24.7	51.8	28 4.3	37.6	46 52.6	30	
16 0 51 15.3	13 40.3	39.0	33 23.8	52.3	13 0.1	51.7	29 57.1	37.3	47 32.5	14 0	
30 52 0.6	15 37.2	39.3	36 0.7	52.4	15 35.1	51.6	31 49.0	37.0	48 11.0	30	
17 0 52 47.3	17 35.0	39.6	38 37.9	52.5	18 9.8	51.4	33 39.9	36.6	48 48.1	13 0	
30 53 35.3	19 33.7	39.9	41 15.4	52.6	20 44.0	51.3	35 29.7	36.2	49 23.9	30	
18 0 54 24.7	21 33.4	40.2	43 53.2	52.7	23 17.9	51.1	37 18.4	35.9	49 58.2	12 0	
30 55 15.4	23 34.1	40.5	46 31.3	52.8	25 51.2	51.0	39 6.2	35.6	50 31.2	30	
19 0 56 7.5	25 35.7	40.8	49 9.6	52.9	28 24.2	50.8	40 52.9	35.2	51 2.9	11 0	
30 57 0.9	27 38.2	41.1	51 48.3	53.0	30 56.7	50.7	42 38.4	34.9	51 33.1	30	
20 0 57 55.6	29 41.6	41.4	54 27.2	53.0	33 28.7	50.5	44 23.0	34.5	52 1.9	10 0	
30 58 51.7	31 45.9	41.7	57 6.3	53.1	36 0.2	50.4	46 6.5	34.1	52 29.4	30	
21 0 59 49.1	33 51.1	42.0	59 45.7	53.2	38 31.3	50.2	47 48.8	33.8	52 55.4	9 0	
30 0 47.8	35 57.2	42.3	2 25.3	53.3	41 1.8	50.0	49 30.1	33.4	53 20.1	30	
22 0 1 47.8	38 4.2	42.6	5 5.1	53.3	43 31.9	49.8	51 10.3	33.0	53 43.3	8 0	
30 2 49.1	40 12.0	42.9	7 45.1	53.4	46 1.4	49.7	52 49.4	32.6	54 5.2	30	
23 0 3 51.8	42 20.7	43.2	10 25.2	53.5	48 30.4	49.5	54 27.3	32.3	54 25.6	7 0	
30 4 55.7	44 30.3	43.4	13 5.6	53.5	50 58.8	49.3	56 4.2	31.9	54 44.6	30	
24 0 6 1.0	46 40.6	43.6	15 46.0	53.6	53 26.6	49.1	57 39.9	31.5	55 2.3	6 0	
30 7 7.4	48 51.9	44.0	18 26.7	53.6	55 53.9	48.9	59 14.4	31.1	55 18.5	30	
25 0 8 15.2	51 3.8	44.3	21 7.5	53.6	58 20.7	48.7	0 47.8	30.8	55 33.3	5 0	
30 9 24.3	53 16.7	44.5	23 48.4	53.7	0 46.8	48.5	2 20.1	30.4	55 46.8	30	
26 0 10 34.7	55 30.3	44.8	26 29.4	53.7	3 12.3	48.3	3 51.2	30.0	55 58.8	4 0	
30 11 46.3	57 44.7	45.0	29 10.5	53.7	5 37.2	48.2	5 21.1	29.6	56 9.4	30	
27 0 12 59.2	59 59.8	45.3	31 51.7	53.7	8 1.5	47.9	6 49.9	29.2	56 18.5	3 0	
30 14 13.3	2 15.8	45.6	34 33.0	53.8	10 25.2	47.7	8 17.4	28.8	56 26.3	30	
28 0 15 28.7	4 32.5	45.8	37 14.3	53.8	12 48.2	47.4	9 43.8	28.4	56 32.7	2 0	
30 16 45.4	6 49.8	46.0	39 55.7	53.8	15 10.5	47.2	11 9.0	28.0	56 37.6	30	
29 0 18 3.2	9 7.8	46.4	42 37.1	53.8	17 32.2	47.0	12 33.0	27.6	56 41.2	1 0	
30 19 22.3	11 26.9	46.6	45 18.6	53.8	19 53.1	46.7	13 55.8	27.2	56 43.3	30	
30 0 20 42.7	13 46.6		48 0.0		22 13.4		15 17.3		56 44.0	0 0	
	85°	87°		89°		92°		94°		91°	
	II <sup>s</sup>	I <sup>s</sup>		Os		XI <sup>s</sup>		X <sup>s</sup>		IX <sup>s</sup>	

TABLE LVI.

81

*Equation II of the Moon's Polar Distance.*

Argument II, corrected.

	III <sup>s</sup>	diff.	IV <sup>s</sup>	diff.	V <sup>s</sup>	diff.	VI <sup>s</sup>	diff.	VII <sup>s</sup>	diff.	VIII <sup>s</sup>	diff.	
°	"	"	"	"	"	"	"	"	"	"	"	"	°
0	0 13.8	0.1	1 24.4	4.6	4 36.9	8.0	9 0.0	9.2	13 23.1	7.9	16 35.6	4.6	30
1	0 13.9	0.2	1 29.0	4.8	4 44.9	8.1	9 9.2	9.2	13 31.0	7.8	16 40.2	4.4	29
2	0 14.1	0.4	1 33.8	4.9	4 53.0	8.1	9 18.4	9.1	13 38.8	7.8	16 44.6	4.3	28
3	0 14.5	0.6	1 38.7	5.1	5 1.1	8.2	9 27.5	9.2	13 46.6	7.6	16 48.9	4.1	27
4	0 15.1	0.7	1 43.8	5.2	5 9.3	8.3	9 36.7	9.2	13 54.2	7.6	16 53.0	3.9	26
5	0 15.8	0.9	1 49.0	5.3	5 17.6	8.4	9 45.9	9.1	14 1.8	7.5	16 56.9	3.8	25
6	0 16.7	1.0	1 54.3	5.5	5 26.0	8.4	9 55.0	9.1	14 9.3	7.4	17 0.7	3.7	24
7	0 17.7	1.2	1 59.8	5.6	5 34.4	8.5	10 4.1	9.1	14 16.7	7.3	17 4.4	3.5	23
8	0 18.9	1.4	2 5.4	5.7	5 42.9	8.5	10 13.2	9.1	14 24.0	7.2	17 7.9	3.4	22
9	0 20.3	1.5	2 11.1	5.8	5 51.4	8.6	10 22.3	9.1	14 31.2	7.0	17 11.3	3.2	21
10	0 21.8	1.7	2 16.9	6.0	6 0.0	8.7	10 31.4	9.0	14 38.2	7.0	17 14.5	3.0	20
11	0 23.5	1.8	2 22.9	6.1	6 8.7	8.7	10 40.4	9.0	14 45.2	6.9	17 17.5	2.9	19
12	0 25.3	2.0	2 29.0	6.2	6 17.4	8.8	10 49.4	9.0	14 52.1	6.8	17 20.4	2.8	18
13	0 27.3	2.1	2 35.2	6.3	6 26.2	8.8	10 58.4	8.9	14 58.9	6.6	17 23.2	2.6	17
14	0 29.4	2.3	2 41.5	6.4	6 35.0	8.8	11 7.3	8.9	15 5.5	6.6	17 25.8	2.5	16
15	0 31.7	2.5	2 47.9	6.6	6 43.8	8.9	11 16.2	8.8	15 12.1	6.4	17 28.3	2.3	15
16	0 34.2	2.6	2 54.5	6.6	6 52.7	8.9	11 25.0	8.8	15 18.5	6.3	17 30.6	2.1	14
17	0 36.8	2.8	3 1.1	6.8	7 1.6	9.0	11 33.8	8.8	15 24.8	6.2	17 32.7	2.0	13
18	0 39.6	2.9	3 7.9	6.9	7 10.6	9.0	11 42.6	8.7	15 31.0	6.1	17 34.7	1.8	12
19	0 42.5	3.0	3 14.8	7.0	7 19.6	9.0	11 51.3	8.7	15 37.1	6.0	17 36.5	1.7	11
20	0 45.5	3.2	3 21.8	7.0	7 28.6	9.1	12 0.0	8.6	15 43.1	5.8	17 38.2	1.5	10
21	0 48.7	3.4	3 28.8	7.2	7 37.7	9.1	12 8.6	8.5	15 48.9	5.7	17 39.7	1.4	9
22	0 52.1	3.5	3 36.0	7.3	7 46.8	9.1	12 17.1	8.5	15 54.6	5.6	17 41.1	1.2	8
23	0 55.6	3.7	3 43.3	7.4	7 55.9	9.1	12 25.6	8.4	16 0.2	5.5	17 42.3	1.0	7
24	0 59.3	3.8	3 50.7	7.5	8 5.0	9.1	12 34.0	8.4	16 5.7	5.3	17 43.3	0.9	6
25	1 3.1	3.9	3 58.2	7.6	8 14.1	9.2	12 42.4	8.3	16 11.0	5.2	17 44.2	0.7	5
26	1 7.0	4.1	4 5.8	7.6	8 23.3	9.2	12 50.7	8.2	16 16.2	5.1	17 44.9	0.6	4
27	1 11.1	4.3	4 13.4	7.8	8 32.5	9.1	12 58.9	8.1	16 21.3	4.9	17 45.5	0.4	3
28	1 15.4	4.4	4 21.2	7.8	8 41.6	9.2	13 7.0	8.1	16 26.2	4.8	17 45.9	0.2	2
29	1 19.8	4.6	4 29.0	7.9	8 50.8	9.2	13 15.1	8.0	16 31.0	4.6	17 46.1	0.1	1
30	1 24.4		4 36.9		9 0.0		13 23.1		16 35.6		17 46.2		0
	II <sup>s</sup>		I <sup>s</sup>		O <sup>s</sup>		XI <sup>s</sup>		X <sup>s</sup>		IX <sup>s</sup>		

TABLE LVII.

*Equation III of Moon's Polar Distance.*

Argument. Moon's True Longitude.

	III <sup>s</sup>	IV <sup>s</sup>	V <sup>s</sup>	VI <sup>s</sup>	VII <sup>s</sup>	VIII <sup>s</sup>	
°	"	"	"	"	"	"	°
0	16.0	14.9	12.0	8.0	4.0	1.1	30
6	16.0	14.5	11.3	7.2	3.3	0.7	24
12	15.8	13.9	10.5	6.3	2.6	0.4	18
18	15.6	13.4	9.7	5.5	2.1	0.2	12
24	15.3	12.7	8.8	4.7	1.5	0.0	6
30	14.9	12.0	8.0	4.0	1.1	0.0	0
	II <sup>s</sup>	I <sup>s</sup>	O <sup>s</sup>	XI <sup>s</sup>	X <sup>s</sup>	IX <sup>s</sup>	

82 TABLE LVIII.

To convert Degrees  
and Minutes into  
Decimal Parts.

Deg. &Min.	Dec parts.
0 0	000
1 5	003
1 26	4
1 48	5
2 10	6
2 31	7
2 53	8
3 14	9
3 36	10
3 58	11
4 19	12
4 41	13
5 2	14
5 24	15
5 46	16
6 7	17
6 29	18
6 50	19
7 12	20
7 34	21
7 55	22
8 17	23
8 38	24
9 0	25
9 22	26
9 43	27
10 5	28
10 26	29
10 48	30
11 10	31
11 31	32
11 53	33
12 14	34
12 36	35
12 58	36
13 19	37
13 41	38
14 2	39
14 24	40
14 46	41
15 7	42
15 29	43
15 50	44
16 12	45
16 34	46
16 55	47
17 17	48
17 38	49
18 0	50
18 22	51
18 43	52
19 5	53

TABLE LIX.

*Equations of Moon's Polar Distance.*  
Arguments, Arg. 20 of Long.; V to IX  
corrected; X not corrected; and XI  
and XII corrected.

Arg.	20	V	VI	VII	VIII	IX	X	XI	Arg.	Arg.	XII	Arg.
	"	"	"	"	"	"	"	"			"	
250	0.3	55.9	6.1	2.6	25.1	3.0	0.7	0.9	250	0	4.0	500
260	0.3	55.8	6.2	2.7	25.1	3.1	0.7	0.9	240	10	3.7	510
270	0.4	55.7	6.3	2.8	25.0	3.2	0.8	1.0	230	20	3.4	520
280	0.6	55.4	6.5	3.0	24.9	3.5	1.0	1.0	220	30	3.1	530
290	0.8	55.1	6.9	3.3	24.8	3.8	1.2	1.1	210	40	2.8	540
300	1.0	54.6	7.3	3.7	24.7	4.3	1.5	1.2	200	50	2.5	550
310	1.3	54.1	7.8	4.2	24.4	4.9	1.8	1.3	190	60	2.3	560
320	1.7	53.4	8.4	4.7	24.1	5.6	2.2	1.4	180	70	2.1	570
330	2.1	52.7	9.1	5.4	23.8	6.4	2.7	1.5	170	80	1.9	580
340	2.6	51.9	9.8	6.1	23.5	7.2	3.2	1.7	160	90	1.7	590
350	3.1	51.0	10.7	6.9	23.2	8.2	3.8	1.9	150	100	1.6	600
360	3.7	50.0	11.6	7.7	22.8	9.2	4.4	2.1	140	110	1.5	610
370	4.3	48.9	12.6	8.7	22.4	10.3	5.1	2.3	130	120	1.5	620
380	4.9	47.7	13.6	9.7	21.9	11.5	5.8	2.5	120	130	1.5	630
390	5.6	46.5	14.8	10.7	21.4	12.8	6.6	2.8	110	140	1.5	640
400	6.4	45.2	16.0	11.8	20.9	14.1	7.4	3.0	100	150	1.6	650
410	7.1	43.9	17.2	13.0	20.4	15.5	8.3	3.3	90	160	1.7	660
420	7.9	42.5	18.5	14.2	19.9	17.0	9.1	3.5	80	170	1.9	670
430	8.8	41.0	19.8	15.5	19.3	18.5	10.1	3.8	70	180	2.1	680
440	9.6	39.5	21.2	16.8	18.7	20.1	11.0	4.1	60	190	2.3	690
450	10.5	38.0	22.6	18.1	18.1	21.7	12.0	4.4	50	200	2.5	700
460	11.3	36.4	24.1	19.4	17.5	23.3	12.9	4.7	40	210	2.8	710
470	12.2	34.9	25.5	20.8	16.9	24.9	13.9	5.0	30	220	3.1	720
480	13.2	33.2	27.0	22.2	16.3	26.6	15.0	5.4	20	230	3.4	730
490	14.1	31.6	28.5	23.6	15.6	28.3	16.0	5.7	10	240	3.7	740
500	15.0	30.0	30.0	25.0	15.0	30.0	17.0	6.0	0	250	4.0	750
510	15.9	28.4	31.5	26.4	14.4	31.7	18.0	6.3	990	260	4.3	760
520	16.8	26.8	33.0	27.8	13.7	33.4	19.0	6.6	980	270	4.6	770
530	17.8	25.1	34.5	29.2	13.1	35.1	20.1	7.0	970	280	4.9	780
540	18.7	23.6	35.9	30.6	12.5	36.7	21.1	7.3	960	290	5.2	790
550	19.5	22.0	37.4	31.9	11.9	38.3	22.0	7.6	950	300	5.5	800
560	20.4	20.5	38.8	33.2	11.3	39.9	23.0	7.9	940	310	5.7	810
570	21.2	19.0	40.2	34.5	10.7	41.5	23.9	8.2	930	320	5.9	820
580	22.1	17.5	41.5	35.8	10.1	43.0	24.9	8.5	920	330	6.1	830
590	22.9	16.1	42.8	37.0	9.6	44.5	25.7	8.7	910	340	6.3	840
600	23.6	14.8	44.0	38.2	9.1	45.9	26.6	9.0	900	350	6.4	850
610	24.4	13.5	45.2	39.3	8.6	47.2	27.4	9.2	890	360	6.5	860
620	25.1	12.3	46.4	40.3	8.1	48.5	28.2	9.5	880	370	6.5	870
630	25.7	11.1	47.4	41.3	7.6	49.7	28.9	9.7	870	380	6.5	880
640	26.3	10.0	48.4	42.3	7.2	50.8	29.6	9.9	860	390	6.5	890
650	26.9	9.0	49.3	43.1	6.8	51.8	30.2	10.1	850	400	6.4	900
660	27.4	8.1	50.2	43.9	6.5	52.8	30.8	10.3	840	410	6.3	910
670	27.9	7.3	50.9	44.6	6.2	53.6	31.3	10.5	830	420	6.1	920
680	28.3	6.6	51.6	45.3	5.9	54.4	31.8	10.6	820	430	5.9	930
690	28.7	5.9	52.2	45.8	5.6	55.1	32.2	10.7	810	440	5.7	940
700	29.0	5.4	52.7	46.3	5.3	55.7	32.5	10.8	800	450	5.5	950
710	29.2	4.9	53.1	46.7	5.2	56.2	32.8	10.9	790	460	5.2	960
720	29.4	4.6	53.5	47.0	5.1	56.5	33.0	11.0	780	470	4.9	970
730	29.6	4.3	53.7	47.2	5.0	56.8	33.2	11.0	770	480	4.6	980
740	29.7	4.2	53.8	47.3	4.9	56.9	33.3	11.1	760	490	4.3	990
750	29.7	4.1	53.9	47.4	4.9	57.0	33.3	11.1	750	500	4.0	1000

Constant 8'

TABLE LX.

TABLE LXI.

*Small Equations of Moon's Parallax.**Moon's Equatorial Parallax.*

Args., 1, 2, 4, 5, 6, 8, 9, 12, 13, of Long.

Argument. Arg. of Evection.

A.	1	2	4	5	6	8	9	12	13	A.
0 0.0	1.6	0.6	1.6	1.9	0.0	3.6	1.4	2.0	100	
3 0.0	1.6	0.6	1.6	1.9	0.0	3.5	1.4	2.0	97	
6 0.0	1.5	0.6	1.5	1.8	0.0	3.1	1.4	1.9	94	
9 0.1	1.5	0.6	1.5	1.8	0.1	2.6	1.3	1.8	91	
12 0.1	1.4	0.5	1.4	1.7	0.2	1.9	1.2	1.7	88	
15 0.1	1.3	0.5	1.3	1.6	0.2	1.3	1.1	1.6	85	
18 0.2	1.1	0.4	1.1	1.4	0.3	0.7	1.0	1.4	82	
21 0.3	1.0	0.4	1.0	1.3	0.5	0.2	0.9	1.2	79	
24 0.4	0.9	0.3	0.9	1.2	0.6	0.0	0.7	1.0	76	
27 0.5	0.7	0.3	0.7	1.0	0.7	0.1	0.6	0.9	73	
30 0.5	0.6	0.2	0.6	0.9	0.8	0.4	0.5	0.7	70	
33 0.6	0.4	0.2	0.4	0.7	0.9	0.8	0.4	0.5	67	
36 0.7	0.3	0.1	0.3	0.6	1.0	1.5	0.3	0.4	64	
39 0.7	0.2	0.1	0.2	0.5	1.1	2.1	0.2	0.2	61	
42 0.8	0.1	0.0	0.1	0.4	1.1	2.8	0.1	0.1	58	
45 0.8	0.0	0.0	0.0	0.3	1.2	3.2	0.0	0.0	55	
48 0.8	0.0	0.0	0.0	0.3	1.2	3.5	0.0	0.0	52	
50 0.8	0.0	0.0	0.0	0.3	1.2	3.6	0.0	0.0	50	

Constant  $\gamma''$ 

The first two figures only of the Arguments are taken.

	O <sub>s</sub>	I <sub>s</sub>	II <sub>s</sub>	III <sub>s</sub>	IV <sub>s</sub>	V <sub>s</sub>	
0	1 20.8	1 15.6	1 1.5	42.6	24.1	10.8	30
1	1 20.8	1 15.2	1 0.9	41.9	23.6	10.5	29
2	1 20.8	1 14.9	1 0.3	41.3	23.0	10.2	28
3	1 20.7	1 14.5	59.7	40.6	22.5	9.9	27
4	1 20.7	1 14.2	59.2	40.0	21.9	9.6	26
5	1 20.6	1 13.8	58.6	39.4	21.4	9.4	25
6	1 20.6	1 13.4	57.9	38.7	20.9	9.1	24
7	1 20.5	1 13.0	57.3	38.1	20.4	8.8	23
8	1 20.4	1 12.6	56.7	37.4	19.9	8.6	22
9	1 20.3	1 12.2	56.1	36.8	19.4	8.4	21
10	1 20.2	1 11.7	55.5	36.1	18.9	8.2	20
11	1 20.1	1 11.3	54.9	35.5	18.4	8.0	19
12	1 19.9	1 10.8	54.2	34.9	17.9	7.8	18
13	1 19.8	1 10.4	53.6	34.2	17.5	7.6	17
14	1 19.6	1 9.9	53.0	33.6	17.0	7.4	16
15	1 19.5	1 9.4	52.3	33.0	16.6	7.2	15
16	1 19.3	1 9.0	51.7	32.4	16.1	7.1	14
17	1 19.1	1 8.5	51.1	31.7	15.7	6.9	13
18	1 18.9	1 8.0	50.4	31.1	15.2	6.8	12
19	1 18.7	1 7.5	49.8	30.5	14.8	6.7	11
20	1 18.4	1 7.0	49.1	29.9	14.4	6.5	10
21	1 18.2	1 6.5	48.5	29.3	14.0	6.4	9
22	1 18.0	1 5.9	47.8	28.7	13.6	6.3	8
23	1 17.7	1 5.4	47.2	28.1	13.2	6.3	7
24	1 17.4	1 4.8	46.5	27.5	12.9	6.2	6
25	1 17.1	1 4.3	45.9	26.9	12.5	6.1	5
26	1 16.9	1 3.8	45.2	26.3	12.1	6.1	4
27	1 16.6	1 3.2	44.6	25.8	11.8	6.1	3
28	1 16.2	1 2.6	43.9	25.2	11.5	6.0	2
29	1 15.9	1 2.1	43.3	24.7	11.1	6.0	1
30	1 15.6	1 1.5	42.6	24.1	10.8	6.0	0
	XI <sub>s</sub>	X <sub>s</sub>	IX <sub>s</sub>	VIII <sub>s</sub>	VII <sub>s</sub>	VI <sub>s</sub>	



*Moon's Equatorial Parallax.*

Argument.      Anomaly.

	O <sup>s</sup>	diff	I <sup>s</sup>	diff	II <sup>s</sup>	diff	III <sup>s</sup>	diff	IV <sup>s</sup>	diff	V <sup>s</sup>	diff	
°	" "	"	" "	"	" "	"	" "	"	" "	"	" "	"	°
0	58 57.7	0.0	58 27.0	2.0	57 7.9	3.1	55 29.8	3.2	54 1.9	2.5	53 3.2	1.4	30
1	58 57.7	0.1	58 25.0	2.0	57 4.8	3.2	55 26.6	3.2	53 59.4	2.5	53 1.8	1.3	29
2	58 57.6	0.2	58 23.0	2.1	57 1.6	3.2	55 23.4	3.2	53 56.9	2.4	53 0.5	1.2	28
3	58 57.4	0.3	58 20.9	2.2	56 58.4	3.2	55 20.2	3.2	53 54.5	2.4	52 59.3	1.2	27
4	58 57.1	0.3	58 18.7	2.2	56 55.2	3.2	55 17.0	3.2	53 52.1	2.4	52 58.1	1.2	26
5	58 56.8	0.4	58 16.5	2.2	56 52.0	3.2	55 13.8	3.2	53 49.7	2.3	52 57.0	1.1	25
6	58 56.4	0.4	58 14.3	2.3	56 48.8	3.3	55 10.6	3.1	53 47.4	2.3	52 55.8	1.0	24
7	58 56.0	0.6	58 12.0	2.4	56 45.5	3.2	55 7.5	3.1	53 45.1	2.2	52 54.8	1.0	23
8	58 55.4	0.6	58 9.6	2.4	56 42.3	3.3	55 4.4	3.1	53 42.9	2.3	52 53.8	1.0	22
9	58 54.8	0.6	58 7.2	2.4	56 39.0	3.3	55 1.3	3.1	53 40.6	2.1	52 52.8	0.9	21
10	58 54.2	0.8	58 4.8	2.5	56 35.7	3.3	54 58.2	3.1	53 38.5	2.2	52 51.9	0.9	20
11	58 53.4	0.8	58 2.3	2.5	56 32.4	3.3	54 55.1	3.0	53 36.3	2.1	52 51.0	0.9	19
12	58 52.6	0.8	57 59.8	2.6	56 29.1	3.3	54 52.1	3.0	53 34.2	2.1	52 50.1	0.8	18
13	58 51.8	1.0	57 57.2	2.6	56 25.8	3.3	54 49.1	3.0	53 32.1	2.0	52 49.3	0.7	17
14	58 50.8	1.0	57 54.6	2.7	56 22.5	3.3	54 46.1	3.0	53 30.1	2.0	52 48.6	0.7	16
15	58 49.8	1.1	57 51.9	2.7	56 19.2	3.3	54 43.1	2.9	53 28.1	1.9	52 47.9	0.7	15
16	58 48.7	1.1	57 49.2	2.8	56 15.9	3.3	54 40.2	2.9	53 26.2	1.9	52 47.2	0.6	14
17	58 47.6	1.2	57 46.4	2.7	56 12.6	3.3	54 37.3	2.9	53 24.3	1.9	52 46.6	0.6	13
18	58 46.4	1.3	57 43.7	2.9	56 9.3	3.3	54 34.4	2.9	53 22.4	1.8	52 46.0	0.5	12
19	58 45.1	1.3	57 40.8	2.8	56 6.0	3.3	54 31.5	2.8	53 20.6	1.8	52 45.5	0.5	11
20	58 43.8	1.4	57 38.0	2.9	56 2.7	3.4	54 28.7	2.8	53 18.8	1.8	52 45.0	0.4	10
21	58 42.4	1.5	57 35.1	2.9	55 59.3	3.3	54 25.9	2.8	53 17.0	1.7	52 44.6	0.4	9
22	58 40.9	1.5	57 32.2	2.9	55 56.0	3.3	54 23.1	2.8	53 15.3	1.6	52 44.2	0.4	8
23	58 39.4	1.6	57 29.3	3.0	55 52.7	3.3	54 20.3	2.7	53 13.7	1.7	52 43.8	0.3	7
24	58 37.8	1.6	57 26.3	3.0	55 49.4	3.3	54 17.6	2.7	53 12.0	1.6	52 43.5	0.2	6
25	58 36.2	1.8	57 23.3	3.0	55 46.1	3.3	54 14.9	2.7	53 10.4	1.5	52 43.3	0.2	5
26	58 34.4	1.7	57 20.2	3.0	55 42.8	3.2	54 12.2	2.6	53 8.9	1.5	52 43.1	0.2	4
27	58 32.7	1.8	57 17.2	3.1	55 39.6	3.2	54 9.6	2.6	53 7.4	1.5	52 42.9	0.1	3
28	58 30.9	1.9	57 14.1	3.1	55 36.4	3.3	54 7.0	2.6	53 5.9	1.4	52 42.8	0.1	2
29	58 29.0	2.0	57 11.0	3.1	55 33.1	3.3	54 4.4	2.5	53 4.5	1.3	52 42.7	0.0	1
30	58 27.0		57 7.9	3.1	55 29.8		54 1.9		53 3.2		52 42.7		0
	XI <sup>s</sup>		X <sup>s</sup>		IX <sup>s</sup>		VIII <sup>s</sup>		VII <sup>s</sup>		VI <sup>s</sup>		

*Moon's Equatorial Parallax.*

Argument. Argument of the Variation.

	O <sup>s</sup>	I <sup>s</sup>	II <sup>s</sup>	III <sup>s</sup>	IV <sup>s</sup>	V <sup>s</sup>	
°	"	"	"	"	"	"	°
0	55.6	42.3	16.0	3.7	17.6	44.0	30
1	55.6	41.5	15.3	3.8	18.5	44.8	29
2	55.5	40.7	14.5	3.8	19.3	45.6	28
3	55.5	39.8	13.8	3.9	20.1	46.3	27
4	55.3	39.0	13.1	4.1	21.0	47.0	26
5	55.2	38.1	12.4	4.3	21.9	47.7	25
6	55.0	37.2	11.7	4.5	22.7	48.4	24
7	54.8	36.3	11.1	4.7	23.6	49.1	23
8	54.6	35.5	10.4	5.0	24.5	49.7	22
9	54.3	34.6	9.8	5.3	25.4	50.3	21
10	54.0	33.7	9.2	5.6	26.3	50.9	20
11	53.7	32.7	8.7	6.0	27.2	51.5	19
12	53.3	31.8	8.2	6.3	28.2	52.1	18
13	52.9	30.9	7.7	6.8	29.1	52.6	17
14	52.5	30.0	7.2	7.2	30.0	53.1	16
15	52.0	29.1	6.7	7.7	30.9	53.5	15
16	51.5	28.2	6.3	8.2	31.8	54.0	14
17	51.0	27.2	5.9	8.7	32.8	54.4	13
18	50.5	26.3	5.6	9.3	33.7	54.8	12
19	49.9	25.4	5.3	9.8	34.6	55.1	11
20	49.4	24.5	5.0	10.5	35.5	55.4	10
21	48.8	23.6	4.7	11.1	36.4	55.7	9
22	48.1	22.7	4.5	11.7	37.3	56.0	8
23	47.4	21.9	4.3	12.4	38.2	56.2	7
24	46.8	21.0	4.1	13.1	39.0	56.4	6
25	46.1	20.1	3.9	13.8	39.9	56.6	5
26	45.4	19.3	3.8	14.5	40.8	56.8	4
27	44.6	18.5	3.7	15.3	41.6	56.9	3
28	43.9	17.6	3.7	16.1	42.4	56.9	2
29	43.1	16.8	3.7	16.8	43.2	57.0	1
30	42.3	16.0	3.7	17.6	44.0	57.0	0
	XI <sup>s</sup>	X <sup>s</sup>	IX <sup>s</sup>	VIII <sup>s</sup>	VII <sup>s</sup>	VI <sup>s</sup>	

*Reduction of the Parallax,  
and also of the Latitude.*

*Moon's Semi-diameter.*

Argument. Latitude.

Argument. Equatorial Parallax.

Lat.	Red. of par	Red. of Lat.	Eq. Par Semidia.		Eq. Par Semidia.		Eq. Par Semidia.		sec	Fro. Par.
°	"	"	"	"	"	"	"	"		
0	0.0	0 0.0	53 0	14 26.5	56 0	15 15.6	59 0	16 4.6	1	0.3
3	0.0	1 11.8	53 10	14 29.3	56 10	15 18.3	59 10	16 7.4	2	0.5
6	0.1	2 22.7	53 20	14 32.0	56 20	15 21.0	59 20	16 10.1	3	0.8
9	0.3	3 32.1	53 30	14 34.7	56 30	15 23.8	59 30	16 12.8	4	1.1
12	0.5	4 39.3	53 40	14 37.4	56 40	15 26.5	59 40	16 15.6	5	1.4
15	0.7	5 43.4	53 50	14 40.2	56 50	15 29.2	59 50	16 18.3	6	1.6
18	1.0	6 43.7	54 0	14 42.9	57 0	15 31.9	60 0	16 21.0	7	1.9
21	1.4	7 39.7	54 10	14 45.6	57 10	15 34.7	60 10	16 23.7	8	2.2
24	1.8	8 30.7	54 20	14 48.3	57 20	15 37.4	60 20	16 26.4	9	2.4
27	2.3	9 16.1	54 30	14 51.1	57 30	15 40.1	60 30	16 29.2	10	2.7
30	2.7	9 55.4	54 40	14 53.8	57 40	15 42.8	60 40	16 31.9		
33	3.3	10 28.3	54 50	14 56.5	57 50	15 45.6	60 50	16 34.6		
36	3.8	10 54.3	55 0	14 59.2	58 0	15 48.3	61 0	16 37.3		
39	4.4	11 13.2	55 10	15 2.0	58 10	15 51.0	61 10	16 40.1		
42	4.9	11 21.7	55 20	15 4.7	58 20	15 53.7	61 20	16 42.8		
45	5.5	11 28.7	55 30	15 7.4	58 30	15 56.5	61 30	16 45.5		
48	6.1	11 25.2	55 40	15 10.1	58 40	15 59.2	61 40	16 48.2		
51	6.7	11 14.1	55 50	15 12.9	58 50	16 1.9	61 50	16 51.0		
54	7.2	10 55.7	56 0	15 15.6	59 0	16 4.6	62 0	16 53.7		
57	7.8	10 30.0								
60	8.3	9 57.4								
63	8.8	9 18.3								
66	9.2	8 32.9								
69	9.7	7 42.0								
72	10.0	6 45.9								
75	10.3	5 45.4								
78	10.6	4 41.0								
81	10.8	3 33.5								
84	11.0	2 23.7								
87	11.1	1 12.3								
90	11.1	0 0.0								

TABLE LVI.

*Augmentation of Moon's Semi-diameter.*

Subsidiary Table.

Lat.	+3'	-3'
°	"	"
0	+0.0	-0.0
6	0.0	0.0
12	0.0	0.0
15	0.0	0.0
18	0.1	0.1
24	0.1	0.1
30	0.1	0.1
36	0.2	0.2
42	0.2	0.2
48	0.3	0.3
54	0.3	0.3
60	0.4	0.4
72	0.5	0.5
78	0.6	0.6
84	0.6	0.6
90	+0.6	-0.6

Alt.	Horizon. Semi-diameter.				Alt.	Horizon. Semi-diameter.			
	14° 30'	15'	16'	17'		14° 30'	15'	16'	17'
0	"	"	"	"	0	"	"	"	"
2	0.6	0.6	0.7	0.8	42	9.2	9.8	11.2	12.6
4	1.0	1.1	1.3	1.5	45	9.7	10.4	11.8	13.3
6	1.5	1.6	1.9	2.1	48	10.2	10.9	12.4	14.0
8	2.0	2.1	2.4	2.7	51	10.6	11.4	13.0	14.7
10	2.4	2.6	3.0	3.4	54	11.1	11.8	13.5	15.2
12	2.9	3.1	3.6	4.0	57	11.5	12.3	14.0	15.8
14	3.4	3.6	4.1	4.7	60	11.8	12.7	14.4	16.3
16	3.8	4.1	4.7	5.3	63	12.2	13.0	14.9	16.8
18	4.3	4.6	5.2	5.9	66	12.5	13.4	15.2	17.2
21	4.9	5.3	6.0	6.8	69	12.8	13.7	15.6	17.6
24	5.6	6.0	6.8	7.7	72	13.0	13.9	15.9	17.9
27	6.2	6.7	7.6	8.6	75	13.2	14.1	16.1	18.2
30	6.9	7.3	8.4	9.5	78	13.4	14.3	16.3	18.4
33	7.5	8.0	9.1	10.3	81	13.5	14.4	16.5	18.6
36	8.1	8.6	9.8	11.1	84	13.6	14.5	16.6	18.7
39	8.6	9.2	10.5	11.9	90	13.7	14.6	16.7	18.8

*Moon's Horary Motion in Longitude.*

Arguments. 1 to 18 of Longitude.

Arg.	2	3	4	5	6	1	7	8	9	Arg.
0	5.0	0.0	2.9	1.9	0.0	0.00	0.00	0.00	0.16	100
2	5.0	0.0	2.8	1.9	0.0	0.00	0.00	0.00	0.15	98
4	4.9	0.0	2.8	1.9	0.0	0.01	0.00	0.02	0.15	96
6	4.8	0.1	2.8	1.9	0.1	0.03	0.01	0.05	0.14	94
8	4.7	0.2	2.7	1.8	0.1	0.06	0.01	0.09	0.12	92
10	4.5	0.3	2.6	1.7	0.2	0.09	0.02	0.14	0.10	90
12	4.3	0.4	2.5	1.7	0.2	0.13	0.02	0.19	0.09	88
14	4.1	0.6	2.3	1.6	0.3	0.18	0.03	0.26	0.07	86
16	3.8	0.7	2.2	1.5	0.4	0.23	0.04	0.33	0.05	84
18	3.6	0.9	2.0	1.4	0.5	0.28	0.05	0.41	0.03	82
20	3.3	1.1	1.9	1.3	0.6	0.34	0.06	0.50	0.02	80
22	3.0	1.3	1.7	1.1	0.7	0.40	0.07	0.58	0.01	78
24	2.7	1.5	1.5	1.0	0.8	0.46	0.08	0.67	0.00	76
26	2.3	1.7	1.3	0.9	0.9	0.52	0.10	0.77	0.00	74
28	2.0	1.9	1.2	0.8	1.0	0.58	0.11	0.86	0.00	72
30	1.7	2.1	1.0	0.7	1.1	0.63	0.12	0.94	0.01	70
32	1.4	2.2	0.8	0.5	1.2	0.69	0.13	1.03	0.01	68
34	1.2	2.4	0.7	0.4	1.3	0.74	0.14	1.11	0.03	66
36	0.9	2.6	0.5	0.3	1.3	0.78	0.15	1.18	0.05	64
38	0.7	2.7	0.4	0.3	1.4	0.82	0.16	1.25	0.06	62
40	0.5	2.8	0.3	0.2	1.5	0.86	0.16	1.30	0.08	60
42	0.3	2.9	0.2	0.1	1.5	0.89	0.17	1.35	0.10	58
44	0.2	3.0	0.1	0.1	1.6	0.91	0.17	1.39	0.11	56
46	0.1	3.1	0.0	0.0	1.6	0.93	0.18	1.42	0.12	54
48	0.0	3.1	0.0	0.0	1.6	0.94	0.18	1.44	0.13	52
50	0.0	3.1	0.0	0.0	1.6	0.94	0.18	1.44	0.13	50

Arg.	10	11	12	13	14	15	16	17	18	Arg.
0	0.00	0.26	0.00	0.00	0.00	0.00	0.26	0.00	0.21	100
2	0.00	0.25	0.09	0.00	0.00	0.00	0.26	0.00	0.20	98
4	0.02	0.24	0.01	0.00	0.01	0.00	0.26	0.00	0.20	96
6	0.04	0.22	0.03	0.01	0.02	0.01	0.25	0.00	0.20	94
8	0.08	0.20	0.04	0.02	0.04	0.01	0.25	0.01	0.20	92
10	0.12	0.17	0.07	0.03	0.06	0.02	0.24	0.01	0.20	90
12	0.16	0.14	0.09	0.04	0.09	0.02	0.22	0.02	0.19	88
14	0.20	0.11	0.12	0.06	0.12	0.03	0.21	0.02	0.19	86
16	0.24	0.08	0.16	0.07	0.15	0.04	0.20	0.03	0.18	84
18	0.28	0.05	0.19	0.09	0.19	0.05	0.19	0.04	0.18	82
20	0.31	0.03	0.23	0.11	0.22	0.06	0.17	0.05	0.17	80
22	0.34	0.01	0.27	0.13	0.26	0.07	0.15	0.06	0.17	78
24	0.35	0.00	0.31	0.15	0.30	0.08	0.14	0.07	0.16	76
26	0.36	0.00	0.35	0.17	0.34	0.08	0.12	0.07	0.16	74
28	0.35	0.01	0.39	0.19	0.33	0.09	0.11	0.08	0.15	72
30	0.34	0.02	0.43	0.21	0.42	0.10	0.09	0.09	0.15	70
32	0.32	0.04	0.47	0.23	0.45	0.11	0.07	0.10	0.14	68
34	0.29	0.06	0.50	0.25	0.49	0.12	0.06	0.11	0.14	66
36	0.26	0.09	0.54	0.26	0.52	0.13	0.05	0.12	0.13	64
38	0.22	0.11	0.57	0.28	0.55	0.14	0.04	0.12	0.13	62
40	0.18	0.14	0.59	0.29	0.58	0.14	0.02	0.13	0.12	60
42	0.15	0.16	0.62	0.30	0.60	0.15	0.01	0.13	0.12	58
44	0.12	0.19	0.63	0.31	0.62	0.15	0.01	0.14	0.12	56
46	0.10	0.21	0.65	0.32	0.63	0.16	0.00	0.14	0.12	54
48	0.09	0.22	0.66	0.32	0.64	0.16	0.00	0.14	0.12	52
50	0.08	0.22	0.66	0.32	0.64	0.16	0.00	0.14	0.11	50

*Moon's Horary Motion in Longitude.*

Argument. Argument of the Evection.

	O <sup>s</sup>	I <sup>s</sup>	II <sup>s</sup>	III <sup>s</sup>	IV <sup>s</sup>	V <sup>s</sup>	
°	"	"	"	"	"	"	°
0	80.3	74.7	59.6	39.4	19.8	5.9	30
1	80.3	74.3	58.9	38.7	19.3	5.6	29
2	80.3	73.9	58.3	38.0	18.7	5.3	28
3	80.2	73.5	57.7	37.3	18.1	5.0	27
4	80.2	73.1	57.1	36.6	17.6	4.7	26
5	80.1	72.7	56.4	36.0	17.0	4.4	25
6	80.1	72.3	55.8	35.3	16.5	4.1	24
7	80.0	71.9	55.1	34.6	15.9	3.8	23
8	79.9	71.4	54.5	33.9	15.4	3.6	22
9	79.8	71.0	53.8	33.2	14.9	3.4	21
10	79.7	70.5	53.1	32.5	14.4	3.1	20
11	79.5	70.1	52.5	31.9	13.9	2.9	19
12	79.4	69.6	51.8	31.2	13.4	2.7	18
13	79.2	69.1	51.1	30.5	12.9	2.5	17
14	79.1	68.6	50.5	29.9	12.4	2.3	16
15	78.9	68.1	49.8	29.2	11.9	2.1	15
16	78.7	67.6	49.1	28.6	11.4	2.0	14
17	78.5	67.0	48.4	27.9	11.0	1.8	13
18	78.2	66.5	47.7	27.2	10.5	1.7	12
19	78.0	66.0	47.0	26.6	10.1	1.6	11
20	77.8	65.4	46.4	26.0	9.7	1.4	10
21	77.5	64.9	45.7	25.3	9.3	1.3	9
22	77.2	64.3	45.0	24.7	8.8	1.2	8
23	77.0	63.7	44.3	24.1	8.4	1.2	7
24	76.7	63.2	43.6	23.5	8.0	1.1	6
25	76.4	62.6	42.9	22.8	7.7	1.0	5
26	76.1	62.0	42.2	22.2	7.3	1.0	4
27	75.7	61.4	41.5	21.6	6.9	0.9	3
28	75.4	60.8	40.8	21.0	6.6	0.9	2
29	75.0	60.2	40.1	20.4	6.2	0.9	1
30	74.7	59.6	39.4	19.8	5.9	0.9	0
	XI <sup>s</sup>	X <sup>s</sup>	IX <sup>s</sup>	VIII <sup>s</sup>	VII <sup>s</sup>	VI <sup>s</sup>	

TABLE LXIX.

*Moon's Horary Motion in Longitude.*

Arguments. Sum of Equations, 2, 3, &amp;c., and Evection corrected.

	0''	10''	20''	
O <sup>s</sup> 0	0.0	0.2	0.5	XII <sup>s</sup> 0
I 0	0.0	0.2	0.4	XI 0
II 0	0.1	0.2	0.3	X 0
III 0	0.2	0.2	0.2	IX 0
IV 0	0.3	0.2	0.1	VIII 0
V 0	0.4	0.2	0.0	VII 0
VI 0	0.5	0.2	0.0	VI 0
	0''	10''	20''	



*Moon's Horary Motion in Longitude.*

Arguments. Sum of preceding equations, and Anomaly corrected.

	0"	10"	20"	30"	40"	50"	60"	70"	80"	90"	100"	
<sup>s</sup> °	"	"	"	"	"	"	"	"	"	"	"	<sup>s</sup> °
O 0	4.1	5.3	6.5	7.6	8.8	10.0	11.2	12.4	13.5	14.7	15.9	XII 0
5	4.1	5.3	6.5	7.7	8.8	10.0	11.2	12.3	13.5	14.7	15.9	25
10	4.2	5.4	6.5	7.7	8.8	10.0	11.2	12.3	13.5	14.6	15.8	20
15	4.3	5.5	6.6	7.7	8.9	10.0	11.1	12.3	13.4	14.5	15.7	15
20	4.5	5.6	6.7	7.8	8.9	10.0	11.1	12.2	13.3	14.4	15.5	10
25	4.8	5.8	6.9	7.9	9.0	10.0	11.0	12.1	13.1	14.2	15.2	5
I 0	5.1	6.0	7.0	8.0	9.0	10.0	11.0	12.0	13.0	14.0	14.9	XI 0
5	5.4	6.3	7.2	8.2	9.1	10.0	10.9	11.8	12.8	13.7	14.6	25
10	5.7	6.6	7.4	8.3	9.2	10.0	10.8	11.7	12.6	13.4	14.3	20
15	6.1	6.9	7.7	8.5	9.2	10.0	10.8	11.5	12.3	13.1	13.9	15
20	6.6	7.2	7.9	8.6	9.3	10.0	10.7	11.4	12.1	12.8	13.4	10
25	7.0	7.6	8.2	8.8	9.4	10.0	10.6	11.2	11.8	12.4	13.0	5
II 0	7.5	8.0	8.5	9.0	9.5	10.0	10.5	11.0	11.5	12.0	12.5	X 0
5	7.9	8.4	8.8	9.2	9.6	10.0	10.4	10.8	11.2	11.6	12.1	25
10	8.4	8.7	9.1	9.4	9.7	10.0	10.3	10.6	10.9	11.3	11.6	20
15	8.9	9.1	9.4	9.6	9.8	10.0	10.2	10.4	10.6	10.9	11.1	15
20	9.4	9.5	9.7	9.8	9.9	10.0	10.1	10.2	10.3	10.5	10.6	10
25	9.9	9.9	9.9	10.0	10.0	10.0	10.0	10.0	10.1	10.1	10.1	5
III 0	10.4	10.3	10.2	10.1	10.1	10.0	9.9	9.9	9.8	9.7	9.6	IX 0
5	10.8	10.7	10.5	10.3	10.2	10.0	9.8	9.7	9.5	9.3	9.2	25
10	11.3	11.0	10.8	10.5	10.3	10.0	9.7	9.5	9.2	9.0	8.7	20
15	11.7	11.4	11.0	10.7	10.3	10.0	9.7	9.3	9.0	8.6	8.3	15
20	12.1	11.7	11.3	10.9	10.4	10.0	9.6	9.1	8.7	8.3	7.9	10
25	12.5	12.0	11.5	11.0	10.5	10.0	9.5	9.0	8.5	8.0	7.5	5
IV 0	12.9	12.3	11.7	11.2	10.6	10.0	9.4	8.8	8.3	7.7	7.1	VIII 0
5	13.3	12.6	11.9	11.3	10.6	10.0	9.4	8.7	8.1	7.4	6.7	25
10	13.6	12.9	12.1	11.4	10.7	10.0	9.3	8.6	7.9	7.1	6.4	20
15	13.9	13.1	12.3	11.5	10.8	10.0	9.2	8.5	7.7	6.9	6.1	15
20	14.1	13.3	12.5	11.6	10.8	10.0	9.2	8.4	7.5	6.7	5.9	10
25	14.4	13.5	12.6	11.7	10.9	10.0	9.1	8.3	7.4	6.5	5.6	5
V 0	14.6	13.7	12.7	11.8	10.9	10.0	9.1	8.2	7.3	6.3	5.4	VII 0
5	14.7	13.8	12.8	11.9	10.9	10.0	9.1	8.1	7.2	6.2	5.3	25
10	14.9	13.9	12.9	12.0	11.0	10.0	9.0	8.0	7.1	6.1	5.1	20
15	15.0	14.0	13.0	12.0	11.0	10.0	9.0	8.0	7.0	6.0	5.0	15
20	15.1	14.1	13.0	12.0	11.0	10.0	9.0	8.0	7.0	5.9	4.9	10
25	15.1	14.1	13.1	12.0	11.0	10.0	9.0	8.0	6.9	5.9	4.9	5
VI 0	15.1	14.1	13.1	12.1	11.0	10.0	9.0	8.0	6.9	5.9	4.9	VI 0
	0"	10"	20"	30"	40"	50"	60"	70"	80"	90"	100"	

*Moon's Horary Motion in Longitude.*

Argument. Anomaly corrected.

	Os	diff.	Is	diff.	IIs	diff.	IIIs	diff.	IVs	diff.	Vs	diff.	
°	"		"		"		"		"		"		°
0	441.5	"	404.1	"	309.3	"	195.3	"	95.8	"	30.6	"	30
1	441.5	0.0	401.6	2.5	305.6	3.7	191.6	3.7	93.0	2.8	29.2	1.4	29
2	441.3	0.1	399.2	2.4	301.9	3.7	187.9	3.7	90.2	2.8	27.8	1.4	28
3	441.1	0.2	396.6	2.6	298.1	3.8	184.3	3.6	87.6	2.6	26.4	1.4	27
4	440.8	0.3	394.0	2.6	294.4	3.7	180.6	3.7	84.9	2.7	25.1	1.3	26
5	440.4	0.4	391.3	2.7	290.6	3.8	177.0	3.6	82.3	2.6	23.8	1.3	25
		0.5		2.7		3.8		3.6		2.6		1.2	
6	439.9	0.5	388.6	2.8	286.8	3.8	173.4	3.6	79.7	2.6	22.6	1.2	24
7	439.4	0.7	385.8	2.8	283.0	3.8	169.8	3.5	77.1	2.5	21.4	1.1	23
8	438.7	0.7	383.0	2.9	279.2	3.8	166.3	3.5	74.6	2.5	20.3	1.1	22
9	438.0	0.8	380.1	3.0	275.4	3.9	162.8	3.5	72.1	2.5	19.2	1.0	21
10	437.2	0.9	377.1	3.0	271.5	3.8	159.3	3.5	69.7	2.4	18.2	1.0	20
				3.0		3.8		3.5		2.4		1.0	
11	436.3	1.0	374.1	3.0	267.7	3.9	155.8	3.4	67.3	2.3	17.2	0.9	19
12	435.3	1.1	371.1	3.1	263.8	3.8	152.4	3.5	65.0	2.3	16.3	0.9	18
13	434.2	1.1	368.0	3.2	260.0	3.8	148.9	3.4	62.7	2.3	15.4	0.8	17
14	433.1	1.3	364.8	3.2	256.2	3.9	145.5	3.3	60.4	2.2	14.6	0.8	16
15	431.8	1.3	361.6	3.2	252.3	3.8	142.2	3.3	58.2	2.1	13.8	0.7	15
				3.2		3.8		3.3		2.1		0.7	
16	430.5	1.4	358.4	3.3	248.5	3.9	138.9	3.3	56.1	2.2	13.1	0.7	14
17	429.1	1.5	355.1	3.3	244.6	3.8	135.6	3.3	53.9	2.0	12.4	0.6	13
18	427.6	1.5	351.8	3.4	240.8	3.9	132.3	3.2	51.9	2.1	11.8	0.6	12
19	426.1	1.6	348.4	3.4	236.9	3.8	129.1	3.2	49.8	1.9	11.2	0.5	11
20	424.5	1.7	345.0	3.4	233.1	3.8	125.9	3.2	47.9	2.0	10.7	0.5	10
				3.4		3.8		3.2		2.0		0.5	
21	422.7	1.7	341.6	3.5	229.3	3.9	122.7	3.1	45.9	1.9	10.2	0.4	9
22	421.0	1.9	338.1	3.5	225.4	3.8	119.6	3.1	44.0	1.8	9.8	0.4	8
23	419.1	1.9	334.6	3.5	221.6	3.8	116.5	3.1	42.2	1.8	9.4	0.3	7
24	417.2	2.0	331.1	3.6	217.8	3.8	113.4	3.0	40.4	1.7	9.1	0.3	6
25	415.2	2.1	327.5	3.5	214.0	3.7	110.4	3.0	38.7	1.7	8.8	0.2	5
				3.5		3.7		3.0		1.7		0.2	
26	413.1	2.2	324.0	3.7	210.3	3.8	107.4	2.9	37.0	1.7	8.6	0.2	4
27	410.9	2.2	320.3	3.6	206.5	3.7	104.5	2.9	35.3	1.6	8.4	0.1	3
28	408.7	2.3	316.7	3.7	202.8	3.8	101.6	2.9	33.7	1.6	8.3	0.1	2
29	406.4	2.3	313.0	3.7	199.0	3.7	98.7	2.9	32.1	1.5	8.2	0.0	1
30	404.1		309.3		195.3		95.8		30.6		8.2		0
	XIs		Xs		IXs		VIII <sub>s</sub>		VII <sub>s</sub>		VI <sub>s</sub>		

*Moon's Horary Motion in Longitude.*

Arguments. Sum of preceding Equations, and Arg. of Variation.

	"	"	"	"	"	"	"	"	"	"	"	"	"	"	
	0	50	100	150	200	250	300	350	400	450	500	550	600		
<sup>s</sup> °	"	"	"	"	"	"	"	"	"	"	"	"	"	<sup>s</sup> °	
O 0	4.5	5.5	6.5	7.6	8.6	9.6	10.6	11.6	12.6	13.7	14.7	15.7	16.7	XII 0	
5	4.6	5.6	6.6	7.6	8.6	9.6	10.6	11.6	12.6	13.6	14.6	15.6	16.6	25	
10	4.8	5.8	6.8	7.7	8.7	9.6	10.6	11.5	12.5	13.4	14.4	15.3	16.3	20	
15	5.3	6.1	7.0	7.9	8.8	9.7	10.5	11.4	12.3	13.1	14.0	14.9	15.8	15	
20	5.8	6.6	7.4	8.2	8.9	9.7	10.5	11.2	12.0	12.8	13.5	14.3	15.1	10	
25	6.6	7.2	7.8	8.5	9.1	9.7	10.4	11.0	11.7	12.3	12.9	13.6	14.2	5	
I 0	7.4	7.8	8.3	8.8	9.3	9.8	10.3	10.8	11.3	11.8	12.3	12.7	13.2	XI 0	
5	8.3	8.6	8.9	9.2	9.5	9.9	10.2	10.5	10.8	11.2	11.5	11.8	12.1	25	
10	9.2	9.3	9.5	9.6	9.8	9.9	10.1	10.2	10.4	10.5	10.7	10.8	11.0	20	
15	10.2	10.1	10.1	10.1	10.0	10.0	10.0	10.0	9.9	9.9	9.9	9.8	9.8	15	
20	11.1	10.9	10.7	10.5	10.3	10.1	9.9	9.7	9.5	9.2	9.0	8.8	8.6	10	
25	12.1	11.7	11.3	10.9	10.5	10.2	9.8	9.4	9.0	8.6	8.3	7.9	7.5	5	
II 0	12.9	12.4	11.8	11.3	10.8	10.2	9.7	9.1	8.6	8.1	7.5	7.0	6.4	X 0	
5	13.7	13.0	12.3	11.6	11.0	10.3	9.6	8.9	8.2	7.5	6.9	6.2	5.5	25	
10	14.3	13.5	12.7	11.9	11.1	10.3	9.5	8.7	7.9	7.1	6.3	5.5	4.7	20	
15	14.9	14.0	13.1	12.2	11.3	10.4	9.5	8.6	7.7	6.8	5.8	4.9	4.0	15	
20	15.3	14.3	13.3	12.3	11.4	10.4	9.4	8.4	7.5	6.5	5.5	4.5	3.6	10	
25	15.5	14.5	13.5	12.4	11.4	10.4	9.4	8.4	7.4	6.3	5.3	4.3	3.3	5	
III 0	15.6	14.5	13.5	12.5	11.4	10.4	9.4	8.4	7.3	6.3	5.3	4.2	3.2	IX 0	
5	15.4	14.4	13.4	12.4	11.4	10.4	9.4	8.4	7.4	6.4	5.4	4.4	3.3	25	
10	15.2	14.2	13.3	12.3	11.3	10.4	9.4	8.5	7.5	6.5	5.6	4.6	3.6	20	
15	14.8	13.9	13.0	12.1	11.2	10.4	9.5	8.6	7.7	6.8	5.9	5.1	4.2	15	
20	14.2	13.4	12.6	11.9	11.1	10.3	9.5	8.8	8.0	7.2	6.4	5.6	4.9	10	
25	13.5	12.9	12.2	11.6	10.9	10.3	9.6	9.0	8.4	7.6	7.0	6.3	5.7	5	
IV 0	12.7	12.2	11.7	11.2	10.7	10.2	9.7	9.2	8.7	8.2	7.7	7.2	6.7	VIII 0	
5	11.9	11.5	11.2	10.8	10.5	10.1	9.8	9.5	9.1	8.8	8.4	8.1	7.7	25	
10	10.9	10.7	10.6	10.4	10.2	10.1	9.9	9.7	9.6	9.4	9.2	9.1	8.9	20	
15	9.9	9.9	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0	10.1	10.1	10.1	15	
20	8.9	9.1	9.3	9.5	9.7	9.9	10.1	10.3	10.5	10.7	10.9	11.1	11.3	10	
25	8.0	8.4	8.7	9.1	9.5	9.9	10.2	10.6	11.0	11.3	11.7	12.1	12.5	5	
V 0	7.1	7.6	8.2	8.7	9.2	9.8	10.3	10.9	11.4	11.9	12.5	13.0	13.6	VII 0	
5	6.3	7.0	7.6	8.3	9.0	9.7	10.4	11.1	11.8	12.5	13.2	13.9	14.6	25	
10	5.6	6.4	7.2	8.0	8.8	9.7	10.5	11.3	12.1	13.0	13.8	14.6	15.4	20	
15	5.0	5.9	6.8	7.8	8.7	9.6	10.6	11.5	12.4	13.3	14.3	15.2	16.1	15	
20	4.6	5.6	6.6	7.6	8.6	9.6	10.6	11.6	12.6	13.6	14.6	15.7	16.7	10	
25	4.3	5.4	6.4	7.5	8.5	9.6	10.6	11.7	12.7	13.8	14.9	15.9	17.0	5	
VI 0	4.2	5.3	6.4	7.4	8.5	9.6	10.6	11.7	12.8	13.9	14.9	16.0	17.1	VI 0	
	"	"	"	"	"	"	"	"	"	"	"	"	"		
	0	50	100	150	200	250	300	350	400	450	500	550	600		

*Moon's Horary Motion in Longitude.*

Argument.    Argument of the Variation.

	Os	Is	II <sup>s</sup>	III <sup>s</sup>	IV <sup>s</sup>	V <sup>s</sup>	
°	"	"	"	"	"	"	°
0	77.2	57.8	20.3	2.4	21.5	59.7	30
1	77.2	56.7	19.2	2.5	22.7	60.9	29
2	77.1	55.5	18.1	2.6	23.8	62.0	28
3	77.0	54.3	17.0	2.7	25.0	63.1	27
4	76.8	53.1	16.0	2.9	26.2	64.2	26
5	76.6	51.8	15.0	3.1	27.5	65.3	25
6	76.4	50.5	14.1	3.3	28.7	66.3	24
7	76.1	49.3	13.2	3.7	30.0	67.3	23
8	75.7	48.0	12.3	4.0	31.3	68.3	22
9	75.3	46.7	11.4	4.4	32.6	69.2	21
10	74.9	45.4	10.6	4.9	33.9	70.1	20
11	74.4	44.1	9.8	5.3	35.2	70.9	19
12	73.9	42.8	9.0	5.9	36.5	71.7	18
13	73.3	41.5	8.3	6.4	37.8	72.5	17
14	72.7	40.2	7.6	7.0	39.2	73.3	16
15	72.0	38.9	7.0	7.7	40.5	74.0	15
16	71.3	37.5	6.4	8.3	41.8	74.7	14
17	70.6	36.2	5.8	9.1	43.2	75.3	13
18	69.8	34.9	5.3	9.8	44.5	75.8	12
19	69.0	33.6	4.8	10.6	45.8	76.4	11
20	68.1	32.3	4.4	11.5	47.2	76.9	10
21	67.2	31.1	4.0	12.3	48.5	77.3	9
22	66.3	29.8	3.7	13.2	49.8	77.7	8
23	65.3	28.6	3.3	14.2	51.1	78.1	7
24	64.4	27.3	3.1	15.1	52.4	78.4	6
25	63.4	26.1	2.9	16.1	53.6	78.6	5
26	62.3	24.9	2.7	17.1	54.9	78.9	4
27	61.2	23.7	2.5	18.2	56.1	79.0	3
28	60.1	22.5	2.5	19.3	57.3	79.2	2
29	59.0	21.4	2.4	20.4	58.5	79.2	1
30	57.8	20.3	2.4	21.5	59.7	79.2	0
	XI <sup>s</sup>	X <sup>s</sup>	IX <sup>s</sup>	VIII <sup>s</sup>	VII <sup>s</sup>	VI <sup>s</sup>	

*Moon's Horary Motion in Longitude.*

Arguments. Arg. of Reduction and Sum of preceding Equations.

		"	"	"	"	"	"	"	"	"	"	"	"	"	"	"	
		0	50	100	150	200	250	300	350	400	450	500	550	600	650		
<sup>s</sup> °		"	"	"	"	"	"	"	"	"	"	"	"	"	"	<sup>s</sup> °	
O	0	3.3	3.1	2.9	2.7	2.5	2.3	2.1	1.9	1.7	1.5	1.3	1.1	0.9	0.7	XII 0	
	5	3.3	3.1	2.9	2.7	2.5	2.3	2.1	1.9	1.7	1.5	1.3	1.1	0.9	0.7	25	
	10	3.2	3.0	2.8	2.6	2.4	2.3	2.1	1.9	1.7	1.5	1.3	1.1	1.0	0.8	20	
	15	3.1	2.9	2.8	2.6	2.4	2.2	2.1	1.9	1.7	1.5	1.4	1.2	1.0	0.9	15	
	20	3.0	2.8	2.7	2.5	2.4	2.2	2.1	1.9	1.8	1.6	1.5	1.3	1.1	1.0	10	
I	25	2.8	2.7	2.6	2.4	2.3	2.2	2.1	1.9	1.8	1.7	1.5	1.4	1.3	1.2	5	
	0	2.6	2.5	2.4	2.3	2.2	2.1	2.0	1.9	1.8	1.7	1.6	1.5	1.4	1.3	XI 0	
	5	2.4	2.4	2.3	2.2	2.2	2.1	2.0	2.0	1.9	1.8	1.8	1.7	1.6	1.6	25	
	10	2.2	2.2	2.2	2.1	2.1	2.0	2.0	2.0	1.9	1.9	1.9	1.8	1.8	1.8	20	
	15	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	15	
II	20	1.8	1.8	1.8	1.9	1.9	1.9	2.0	2.0	2.1	2.1	2.1	2.2	2.2	2.2	10	
	25	1.6	1.6	1.7	1.8	1.8	1.9	2.0	2.0	2.1	2.2	2.2	2.3	2.4	2.4	5	
	0	1.4	1.5	1.6	1.7	1.8	1.9	2.0	2.1	2.2	2.3	2.4	2.5	2.6	2.7	X 0	
	5	1.2	1.3	1.4	1.6	1.7	1.8	1.9	2.1	2.2	2.3	2.5	2.6	2.7	2.8	25	
	10	1.0	1.2	1.3	1.5	1.6	1.8	1.9	2.1	2.2	2.4	2.5	2.7	2.9	3.0	20	
III	15	0.9	1.1	1.2	1.4	1.6	1.8	1.9	2.1	2.3	2.5	2.6	2.8	3.0	3.1	15	
	20	0.8	1.0	1.2	1.4	1.6	1.7	1.9	2.1	2.3	2.5	2.7	2.9	3.0	3.2	10	
	25	0.7	0.9	1.1	1.3	1.5	1.7	1.9	2.1	2.3	2.5	2.7	2.9	3.1	3.3	5	
	0	0.7	0.9	1.1	1.3	1.5	1.7	1.9	2.1	2.3	2.5	2.7	2.9	3.1	3.3	IX 0	
	5	0.7	0.9	1.1	1.3	1.5	1.7	1.9	2.1	2.3	2.5	2.7	2.9	3.1	3.3	25	
IV	10	0.8	1.0	1.2	1.4	1.6	1.7	1.9	2.1	2.3	2.5	2.7	2.9	3.0	3.2	20	
	15	0.9	1.1	1.2	1.4	1.6	1.8	1.9	2.1	2.3	2.5	2.6	2.8	3.0	3.1	15	
	20	1.0	1.2	1.3	1.5	1.6	1.8	1.9	2.1	2.2	2.4	2.5	2.7	2.9	3.0	10	
	25	1.2	1.3	1.4	1.6	1.7	1.8	1.9	2.1	2.2	2.3	2.5	2.6	2.7	2.8	5	
	0	1.4	1.5	1.6	1.7	1.8	1.9	2.0	2.1	2.2	2.3	2.4	2.5	2.6	2.7	VIII 0	
V	5	1.6	1.6	1.7	1.8	1.8	1.9	2.0	2.0	2.1	2.2	2.2	2.3	2.4	2.4	25	
	10	1.8	1.8	1.8	1.9	1.9	1.9	2.0	2.0	2.1	2.1	2.1	2.2	2.2	2.2	20	
	15	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	15	
	20	2.2	2.2	2.2	2.1	2.1	2.0	2.0	2.0	1.9	1.9	1.9	1.8	1.8	1.8	10	
	25	2.4	2.4	2.3	2.2	2.2	2.1	2.0	2.0	1.9	1.8	1.8	1.7	1.6	1.6	5	
VI	0	2.6	2.5	2.4	2.3	2.2	2.1	2.0	1.9	1.8	1.7	1.6	1.5	1.4	1.3	VII 0	
	5	2.8	2.7	2.6	2.4	2.3	2.2	2.1	1.9	1.8	1.7	1.5	1.4	1.3	1.2	25	
	10	3.0	2.8	2.7	2.5	2.4	2.2	2.1	1.9	1.8	1.6	1.5	1.3	1.1	1.0	20	
	15	3.1	2.9	2.8	2.6	2.4	2.2	2.1	1.9	1.7	1.5	1.4	1.2	1.0	0.9	15	
	20	3.2	3.0	2.8	2.6	2.4	2.3	2.1	1.9	1.7	1.5	1.3	1.1	1.0	0.8	10	
VII	25	3.3	3.1	2.9	2.7	2.5	2.3	2.1	1.9	1.7	1.5	1.3	1.1	0.9	0.7	5	
	0	3.3	3.1	2.9	2.7	2.5	2.3	2.1	1.9	1.7	1.5	1.3	1.1	0.9	0.7	VI 0	
		"	"	"	"	"	"	"	"	"	"	"	"	"	"		
		0	50	100	150	200	250	300	350	400	450	500	550	600	650		



*Moon's Horary Motion in Long.*

Arg.    Arg. of Reduction.

	Os VIs	Is VI s	I's VIII s	
°	"	"	"	°
0	2.1	6.0	14.0	30
1	2.1	6.3	14.2	29
2	2.1	6.5	14.4	28
3	2.1	6.8	14.7	27
4	2.2	7.0	14.9	26
5	2.2	7.3	15.1	25
6	2.2	7.5	15.3	24
7	2.3	7.8	15.5	23
8	2.4	8.1	15.7	22
9	2.5	8.4	15.9	21
10	2.5	8.6	16.1	20
11	2.6	8.9	16.2	19
12	2.7	9.2	16.4	18
13	2.9	9.4	16.6	17
14	3.0	9.7	16.7	16
15	3.1	10.0	16.9	15
16	3.3	10.3	17.0	14
17	3.4	10.6	17.1	13
18	3.6	10.8	17.3	12
19	3.8	11.1	17.4	11
20	3.9	11.4	17.5	10
21	4.1	11.6	17.5	9
22	4.3	11.9	17.6	8
23	4.5	12.2	17.7	7
24	4.7	12.5	17.8	6
25	4.9	12.7	17.8	5
26	5.1	13.0	17.8	4
27	5.3	13.2	17.9	3
28	5.6	13.5	17.9	2
29	5.8	13.7	17.9	1
30	6.0	14.0	17.9	0
	XIs Vs	Xs IVs	IXs IIIs	

Constant to be added 27°24'.0.

*Moon's Horary Motion in Long.*  
(Equation of the second order.)

Arguments.    Arg's of Table LXX.

Arg.	"	"	"
	0	50	100
<sup>s</sup> °	"	"	"
O    0	0.05	0.05	0.05
I    0	0.08	0.05	0.02
II   0	0.10	0.05	0.00
III   0	0.10	0.05	0.00
IV   0	0.09	0.05	0.01
V    0	0.07	0.05	0.03
VI   0	0.05	0.05	0.05
VII   0	0.03	0.05	0.07
VIII   0	0.01	0.05	0.09
IX   0	0.00	0.05	0.10
X    0	0.00	0.05	0.10
XI   0	0.02	0.05	0.08
XII   0	0.05	0.05	0.05
	"	"	"
	0	50	100

TABLE LXXVII.

*Moon's Horary Motion in Longitude.*

(Equations of the second order.)

Arguments.    Arguments of Tables LXXII and LXXIV.

		Variation.								Reduction.	
		"	"	"	"	"	"	"	"	"	"
		0	100	200	300	400	500	600	0	600	
<sup>s</sup>	<sup>s</sup> °	"	"	"	"	"	"	"	"	"	"
O.   VI.   0	0.14	0.14	0.14	0.14	0.14	0.14	0.14	0.14	0.03	0.03	
I.   VII.   0	0.22	0.19	0.16	0.13	0.10	0.06	0.02		0.01	0.05	
I.   VII. 15	0.23	0.20	0.17	0.13	0.10	0.05	0.01		0.01	0.06	
II.   VIII.   0	0.22	0.19	0.16	0.13	0.10	0.07	0.03		0.01	0.05	
III.   IX.   0	0.14	0.14	0.14	0.14	0.14	0.14	0.14		0.03	0.03	
IV.   X.    0	0.06	0.09	0.12	0.15	0.18	0.21	0.26		0.05	0.01	
IV.   X.   15	0.05	0.08	0.11	0.15	0.18	0.23	0.28		0.05	0.00	
V.   XI.    0	0.06	0.09	0.12	0.15	0.18	0.22	0.26		0.05	0.01	
VI.   XII.   0	0.14	0.14	0.14	0.14	0.14	0.14	0.14		0.03	0.03	

*Moon's Horary Motion in Longitude.*

(Equations of the second order.)

Arguments. Args. of Evection, Anomaly, Variation, Reduction.

		Evec.	Anom.	Var.	Red.	Evec.	Anom.	Var.	Red.		
	<sup>s</sup> °	"	"	"	"	"	"	"	"	<sup>s</sup> °	
O	0	0.16	1.05	0.34	0.08	0.16	1.05	0.34	0.08	XII	0
	5	0.15	0.93	0.28	0.09	0.18	1.17	0.40	0.06		25
	10	0.13	0.81	0.22	0.10	0.19	1.28	0.46	0.05		20
	15	0.12	0.70	0.17	0.11	0.21	1.40	0.51	0.04		15
	20	0.10	0.59	0.12	0.12	0.22	1.50	0.56	0.03		10
	25	0.09	0.49	0.08	0.13	0.24	1.60	0.60	0.02		5
I	0	0.08	0.40	0.05	0.14	0.25	1.70	0.63	0.01	XI	0
	5	0.07	0.31	0.02	0.15	0.26	1.78	0.66	0.01		25
	10	0.05	0.24	0.01	0.15	0.27	1.86	0.67	0.00		20
	15	0.04	0.17	0.01	0.15	0.28	1.92	0.67	0.00		15
	20	0.03	0.12	0.01	0.15	0.29	1.98	0.67	0.00		10
	25	0.03	0.07	0.03	0.15	0.30	2.02	0.65	0.01		5
II	0	0.02	0.04	0.06	0.14	0.31	2.05	0.62	0.01	X	0
	5	0.01	0.02	0.09	0.13	0.32	2.08	0.59	0.02		25
	10	0.01	0.00	0.13	0.12	0.32	2.09	0.54	0.03		20
	15	0.00	0.00	0.18	0.11	0.32	2.10	0.50	0.04		15
	20	0.00	0.00	0.24	0.10	0.33	2.09	0.44	0.05		10
	25	0.00	0.02	0.29	0.09	0.33	2.08	0.39	0.06		5
III	0	0.00	0.04	0.35	0.08	0.33	2.06	0.33	0.08	IX	0
	5	0.00	0.07	0.40	0.06	0.33	2.03	0.27	0.09		25
	10	0.01	0.10	0.46	0.05	0.32	2.00	0.22	0.10		20
	15	0.01	0.14	0.51	0.04	0.32	1.96	0.17	0.11		15
	20	0.01	0.18	0.56	0.03	0.31	1.91	0.12	0.12		10
	25	0.02	0.23	0.60	0.02	0.31	1.87	0.08	0.13		5
IV	0	0.03	0.28	0.63	0.01	0.30	1.82	0.05	0.14	VIII	0
	5	0.03	0.34	0.66	0.01	0.29	1.76	0.02	0.15		25
	10	0.04	0.39	0.67	0.00	0.28	1.70	0.01	0.15		20
	15	0.05	0.45	0.68	0.00	0.27	1.64	0.00	0.15		15
	20	0.06	0.52	0.67	0.00	0.26	1.58	0.00	0.15		10
	25	0.08	0.58	0.66	0.01	0.25	1.52	0.02	0.15		5
V	0	0.09	0.64	0.64	0.01	0.24	1.45	0.04	0.14	VII	0
	5	0.10	0.71	0.60	0.02	0.23	1.39	0.08	0.13		25
	10	0.11	0.78	0.56	0.03	0.22	1.32	0.12	0.12		20
	15	0.12	0.84	0.51	0.04	0.20	1.25	0.16	0.11		15
	20	0.14	0.91	0.46	0.05	0.19	1.18	0.22	0.10		10
	25	0.15	0.98	0.40	0.06	0.18	1.12	0.28	0.09		5
VI	0	0.16	1.05	0.34	0.08	0.16	1.05	0.34	0.08	VI	0

## TABLE LXXIX.

*Moon's Horary Motion in Latitude.*

Argument. Arg. I of Latitude.

	O <sup>s</sup>	I <sup>s</sup>	II <sup>s</sup>	III <sup>s</sup>	IV <sup>s</sup>	V <sup>s</sup>	
°	"	"	"	"	"	"	°
0	378.0	351.3	289.2	200.0	110.8	45.7	30
1	378.0	352.7	286.5	196.9	108.1	44.2	29
2	377.9	351.1	283.8	193.8	105.4	42.7	28
3	377.8	349.4	281.0	190.7	102.8	41.3	27
4	377.6	347.7	278.3	187.5	100.2	39.9	26
5	377.3	346.0	275.5	184.4	97.7	38.6	25
6	377.0	344.2	272.6	181.3	95.1	37.3	24
7	376.7	342.3	269.8	178.2	92.6	36.1	23
8	376.3	340.5	266.9	175.1	90.2	34.9	22
9	375.8	338.5	264.0	172.1	87.7	33.8	21
10	375.3	336.6	261.1	169.0	85.3	32.7	20
11	374.7	334.5	258.1	165.9	83.0	31.6	19
12	374.1	332.5	255.2	162.9	80.7	30.7	18
13	373.5	330.4	252.2	159.8	78.1	29.7	17
14	372.7	328.3	249.2	156.8	76.1	28.9	16
15	372.0	326.1	246.2	153.8	73.9	28.0	15
16	371.1	323.9	243.2	150.8	71.7	27.3	14
17	370.3	321.9	240.2	147.8	69.6	26.5	13
18	369.3	319.3	237.1	144.8	67.5	25.9	12
19	368.4	317.0	234.1	141.9	65.5	25.3	11
20	367.3	314.7	231.0	138.9	63.4	24.7	10
21	366.2	312.3	227.9	136.0	61.5	24.2	9
22	365.1	309.8	224.9	133.1	59.5	23.7	8
23	363.9	307.4	221.8	130.2	57.7	23.3	7
24	362.7	304.9	218.7	127.4	55.8	23.0	6
25	361.4	302.3	215.6	124.5	54.0	22.7	5
26	360.1	299.8	212.5	121.7	52.3	22.4	4
27	358.7	297.2	209.3	119.0	50.6	22.2	3
28	357.3	294.6	206.2	116.2	48.9	22.1	2
29	355.8	291.9	203.1	113.5	47.3	22.0	1
30	354.3	289.2	200.0	110.8	45.7	22.0	0
	XI <sup>s</sup>	X <sup>s</sup>	IX <sup>s</sup>	VIII <sup>s</sup>	VII <sup>s</sup>	VI <sup>s</sup>	

## TABLE LXXX.

*Moon's Horary Motion in Latitude.*

Arguments. Args. V, VI, VII, VIII, IX, X, XI, and XII, of Latitude.

Arg.	V	VI	VII	VIII	IX	X	XI	XII	Arg.
	"	"	"	"	"	"	"	"	
0	0.00	0.50	0.34	0.00	0.50	0.04	0.12	0.08	1000
50	0.01	0.49	0.33	0.00	0.49	0.04	0.12	0.07	950
100	0.04	0.45	0.30	0.02	0.45	0.04	0.11	0.05	900
150	0.09	0.40	0.27	0.04	0.40	0.03	0.10	0.03	850
200	0.16	0.33	0.22	0.06	0.33	0.03	0.08	0.01	800
250	0.23	0.25	0.17	0.09	0.25	0.02	0.06	0.00	750
300	0.30	0.17	0.12	0.12	0.17	0.01	0.04	0.01	700
350	0.37	0.10	0.07	0.14	0.10	0.01	0.02	0.03	650
400	0.42	0.05	0.04	0.16	0.05	0.00	0.01	0.05	600
450	0.45	0.01	0.01	0.18	0.01	0.00	0.00	0.07	550
500	0.46	0.00	0.00	0.18	0.00	0.00	0.00	0.08	500

TABLE LXXXI. *Moon's Horary Motion in Latitude.* 97  
 Arguments. Preceding equation, and Sum of equations of Horary  
 Motion in Longitude, except the last two.

Pr. eq.	0''	50''	100''	150''	200''	250''	300''	350''	400''	450''	500''	550''	600''	650''	Diff.
20	59.0	54.5	50.0	45.4	40.9	36.4	31.8	27.3	22.8	18.2	13.7	9.1	4.6	0.1	4.5
30	57.4	53.1	48.9	44.6	40.3	36.0	31.7	27.4	23.2	18.9	14.6	10.3	6.0	1.7	4.3
40	55.8	51.8	47.7	43.7	39.7	35.6	31.6	27.6	23.6	19.5	15.5	11.5	7.4	3.4	4.0
50	54.2	50.4	46.6	42.9	39.1	35.3	31.5	27.7	24.0	20.2	16.4	12.6	8.8	5.1	3.8
60	52.6	49.1	45.5	42.0	38.5	34.9	31.4	27.9	24.4	20.8	17.3	13.8	10.2	6.7	3.5
70	51.0	47.7	44.4	41.1	37.9	34.6	31.3	28.0	24.8	21.5	18.2	14.9	11.7	8.4	3.3
80	49.3	46.3	43.3	40.3	37.3	34.2	31.2	28.2	25.2	22.1	19.1	16.1	13.1	10.0	3.0
90	47.7	45.0	42.2	39.4	36.7	33.9	31.1	28.3	25.6	22.8	20.0	17.3	14.5	11.7	2.8
100	46.1	43.6	41.1	38.6	36.0	33.5	31.0	28.5	26.0	23.4	20.9	18.4	15.9	13.4	2.5
110	44.5	42.2	40.0	37.7	35.4	33.2	30.9	28.6	26.4	24.1	21.8	19.6	17.3	15.0	2.3
120	42.9	40.9	38.9	36.9	34.8	32.8	30.8	28.8	26.8	24.8	22.7	20.7	18.7	16.7	2.0
130	41.3	39.5	37.8	36.0	34.2	32.5	30.7	28.9	27.2	25.4	23.7	21.9	20.1	18.4	1.8
140	39.7	38.2	36.7	35.1	33.6	32.1	30.6	29.1	27.6	26.1	24.6	23.0	21.5	20.0	1.5
150	38.1	36.8	35.5	34.3	33.0	31.8	30.5	29.2	28.0	26.7	25.5	24.2	23.0	21.7	1.3
160	36.5	35.4	34.4	33.4	32.4	31.4	30.4	29.4	28.4	27.4	26.4	25.4	24.4	23.3	1.0
170	34.8	34.1	33.3	32.6	31.8	31.1	30.3	29.5	28.8	28.0	27.3	26.5	25.8	25.0	0.8
180	33.2	32.7	32.2	31.7	31.2	30.7	30.2	29.7	29.2	28.7	28.2	27.7	27.2	26.7	0.5
190	31.6	31.4	31.1	30.9	30.6	30.4	30.1	29.8	29.6	29.3	29.1	28.8	28.6	28.3	0.3
200	30.0	30.0	30.0	30.0	30.0	30.0	30.0	30.0	30.0	30.0	30.0	30.0	30.0	30.0	0.0
210	28.4	28.6	28.9	29.1	29.4	29.6	29.9	30.2	30.4	30.7	30.9	31.2	31.4	31.7	0.3
220	26.8	27.3	27.8	28.3	28.8	29.3	29.8	30.3	30.8	31.3	31.8	32.3	32.8	33.3	0.5
230	25.2	25.9	26.7	27.4	28.2	28.9	29.7	30.5	31.2	32.0	32.7	33.5	34.2	35.0	0.8
240	23.5	24.6	25.6	26.6	27.6	28.6	29.6	30.6	31.6	32.6	33.6	34.6	35.6	36.7	1.0
250	21.9	23.2	24.5	25.7	27.0	28.2	29.5	30.8	32.0	33.3	34.5	35.8	37.1	38.3	1.3
260	20.3	21.8	23.3	24.9	26.4	27.9	29.4	30.9	32.4	33.9	35.4	37.0	38.5	40.0	1.5
270	18.7	20.5	22.2	24.0	25.8	27.5	29.3	31.1	32.8	34.6	36.3	38.1	39.9	41.6	1.8
280	17.1	19.1	21.1	23.1	25.2	27.2	29.2	31.2	33.2	35.2	37.3	39.3	41.3	43.3	2.0
290	15.5	17.8	20.0	22.3	24.6	26.8	29.1	31.4	33.6	35.9	38.2	40.4	42.7	45.0	2.3
300	13.9	16.4	18.9	21.4	24.0	26.5	29.0	31.5	34.0	36.6	39.1	41.6	44.1	46.6	2.5
310	12.3	15.0	17.8	20.6	23.3	26.1	28.9	31.7	34.4	37.2	40.0	42.7	45.5	48.3	2.8
320	10.7	13.7	16.7	19.7	22.7	25.8	28.8	31.8	34.8	37.9	40.9	43.9	46.9	50.0	3.0
330	9.0	12.3	15.6	18.9	22.1	25.4	28.7	32.0	35.2	38.5	41.8	45.1	48.3	51.6	3.3
340	7.4	10.9	14.5	18.0	21.5	25.1	28.6	32.1	35.6	39.2	42.7	46.2	49.8	53.3	3.5
350	5.8	9.6	13.4	17.1	20.9	24.7	28.5	32.3	36.0	39.8	43.6	47.4	51.2	54.9	3.8
360	4.2	8.2	12.3	16.3	20.3	24.4	28.4	32.4	36.4	40.5	44.5	48.5	52.6	56.6	4.0
370	2.6	6.9	11.1	15.4	19.7	24.0	28.3	32.6	36.8	41.1	45.4	49.7	54.0	58.3	4.3
380	1.0	5.5	10.0	14.6	19.1	23.6	28.2	32.7	37.2	41.8	46.3	50.9	55.4	59.9	4.5
	0''	50''	100''	150''	200''	250''	300''	350''	400''	450''	500''	550''	600''	650''	

TABLE LXXXII. *Moon's Horary Motion in Latitude.*  
 Argument. Arg. II. of Latitude.

°	Os	Is	IIs	IIIs	IVs	Vs	°
0	9.3	8.7	7.1	5.0	2.9	1.3	30
3	9.3	8.6	6.9	4.8	2.7	1.2	27
6	9.2	8.5	6.7	4.6	2.5	1.1	24
9	9.2	8.3	6.5	4.3	2.3	1.0	21
12	9.2	8.2	6.3	4.1	2.1	0.9	18
15	9.1	8.0	6.1	3.9	2.0	0.9	15
18	9.1	7.9	5.9	3.7	1.8	0.8	12
21	9.0	7.7	5.7	3.5	1.7	0.8	9
24	8.9	7.5	5.4	3.3	1.5	0.8	6
27	8.8	7.3	5.2	3.1	1.4	0.7	3
30	8.7	7.1	5.0	2.9	1.3	0.7	0
	XIs	Xs	IXs	VIIIs	VIIs	VIs	

*Moon's Horary Motion in Latitude.*

Arguments. Preceding equation, and Sum of equations of Horary Motion in Longitude, except the last two.

Prec. equ.	" 0	" 100	" 200	" 300	" 400	" 500	" 600	" 700
"	"	"	"	"	"	"	"	"
0	2.1	1.8	1.5	1.2	0.9	0.6	0.3	0.0
1	1.9	1.6	1.4	1.1	0.9	0.7	0.4	0.2
2	1.7	1.5	1.3	1.1	1.0	0.8	0.6	0.3
3	1.5	1.4	1.2	1.1	1.0	0.9	0.8	0.6
4	1.3	1.2	1.2	1.1	1.1	1.0	0.9	0.9
5	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1
6	0.9	1.0	1.0	1.1	1.1	1.2	1.3	1.3
7	0.7	0.8	1.0	1.1	1.2	1.3	1.4	1.6
8	0.5	0.7	0.9	1.1	1.2	1.4	1.6	1.9
9	0.3	0.6	0.8	1.1	1.3	1.5	1.8	2.0
10	0.1	0.4	0.7	1.0	1.3	1.6	1.9	2.2
"	"	"	"	"	"	"	"	"
0	100	200	300	400	500	600	700	

Constant to be subtracted 237".2.

TABLE LXXXV.

*Moon's Horary Motion in Latitude.*

(Equations of second order.)

Arguments. Preceding equation, and Sum of equations of Horary Motion in Longitude, except the last two.

Prec. equ.	" 0	" 100	" 200	" 300	" 400	" 500	" 600	" 700
"	"	"	"	"	"	"	"	"
0.00	0.65	0.57	0.48	0.39	0.31	0.21	0.12	0.00
0.10	0.62	0.55	0.47	0.39	0.31	0.23	0.15	0.04
0.20	0.69	0.53	0.46	0.39	0.32	0.25	0.18	0.09
0.30	0.66	0.51	0.45	0.39	0.33	0.27	0.21	0.13
0.40	0.63	0.48	0.44	0.39	0.34	0.29	0.24	0.17
0.50	0.50	0.46	0.43	0.38	0.35	0.30	0.27	0.21
0.60	0.47	0.44	0.42	0.38	0.36	0.32	0.29	0.25
0.70	0.44	0.42	0.40	0.38	0.36	0.34	0.32	0.30
0.80	0.41	0.40	0.39	0.38	0.37	0.36	0.35	0.34
0.90	0.38	0.38	0.38	0.38	0.38	0.38	0.38	0.38
1.00	0.35	0.36	0.37	0.38	0.39	0.40	0.41	0.42
1.10	0.32	0.34	0.36	0.38	0.40	0.42	0.44	0.46
1.20	0.29	0.32	0.34	0.38	0.40	0.44	0.47	0.51
1.30	0.26	0.30	0.33	0.38	0.41	0.46	0.49	0.55
1.40	0.23	0.28	0.32	0.37	0.42	0.47	0.52	0.59
1.50	0.20	0.25	0.31	0.37	0.43	0.49	0.55	0.63
1.60	0.17	0.23	0.30	0.37	0.44	0.51	0.58	0.67
1.70	0.14	0.21	0.29	0.37	0.45	0.53	0.61	0.72
1.80	0.11	0.19	0.28	0.37	0.45	0.55	0.64	0.76
"	"	"	"	"	"	"	"	"
0	100	200	300	400	500	600	700	

*Moon's Hor. Motion in Lat*

(Equa. of second order.)

Argument. Arg. I of Lat.

	I	I	
"	"	"	"
O	0	0.90	0.90
5	0.83	0.97	XII
10	0.75	1.05	25
15	0.68	1.12	20
20	0.61	1.19	15
25	0.54	1.26	10
I	0	0.47	5
5	0.41	1.33	XI
10	0.35	1.39	0
15	0.29	1.45	25
20	0.24	1.51	20
25	0.20	1.56	15
II	0	0.16	10
5	0.12	1.60	5
10	0.09	1.64	X
15	0.07	1.68	0
20	0.05	1.71	25
25	0.04	1.73	20
III	0	0.04	15
5	0.04	1.75	10
10	0.05	1.76	5
15	0.07	1.76	IX
20	0.09	1.76	0
25	0.12	1.76	25
IV	0	0.16	20
5	0.20	1.60	15
10	0.24	1.56	10
15	0.29	1.51	5
20	0.35	1.45	VIII
25	0.41	1.39	0
V	0	0.47	25
5	0.54	1.26	20
10	0.61	1.19	15
15	0.68	1.12	10
20	0.75	1.05	5
25	0.83	0.97	VI
VI	0	0.90	0



*Mean New Moons and Arguments, in January.*

Years.	Mean New Moon in. January.	I.	II.	III.	IV.	N.
	<i>d. h. m.</i>					
1821	2 17 59	0092	7859	80	78	823
1822	21 15 32	0602	7182	78	66	930
1823	11 0 20	0304	5787	61	55	953
1824 B	29 21 53	0814	5110	59	43	060
1825	18 6 41	0516	3716	42	32	083
1826	7 15 30	0218	2321	25	21	105
1827	26 13 3	0728	1644	24	09	213
1828 B	15 21 51	0430	0250	07	98	235
1829	4 6 40	0131	8855	90	87	257
1830	23 4 12	0642	8178	88	75	365
1831	12 13 1	0343	6784	71	64	387
1832 B	1 21 50	0045	5389	54	53	409
1833	19 19 22	0555	4712	53	42	517
1834	9 4 11	0257	3318	36	31	539
1835	28 1 43	0768	2641	34	19	647
1836 B	17 10 32	0469	1246	17	08	669
1837	5 19 20	0171	9852	00	97	692
1838	24 16 53	0681	9175	99	85	799
1839	14 1 42	0383	7780	82	74	822
1840 B	3 10 30	0085	6386	65	63	844
1841	21 8 3	0595	5709	63	51	951
1842	10 16 51	0297	4314	46	40	974
1843	29 14 24	0807	3637	44	28	081
1844 B	18 23 13	0509	2243	28	17	104
1845	7 8 1	0211	0848	11	06	126
1846	26 5 34	0721	0171	09	94	234
1847	15 14 22	0423	8777	92	84	256
1848 B	4 23 11	0125	7382	75	73	278
1849	22 20 43	0635	6705	73	61	386
1850	12 5 32	0337	5311	56	50	408
1851	1 14 21	0038	3916	40	39	431
1852 B	20 11 53	0549	3239	38	27	538
1853	8 20 42	0251	1845	21	16	560
1854	27 18 14	0761	1168	19	04	668
1855	17 3 3	0463	9773	02	93	690
1856 B	6 11 51	0164	8379	85	82	713
1857	24 9 24	0675	7702	84	70	820
1858	13 18 13	0376	6307	67	59	843
1859	3 3 1	0078	4913	50	48	865
1860 B	22 0 34	0588	4236	48	36	972

*Mean Lunations and Changes of the Arguments.*

Num.	Lunations.			I.	II.	III.	IV.	N.
	d.	h.	m.					
$\frac{1}{2}$	14	18	22	404	5359	58	50	43
1	29	12	44	808	717	15	99	85
2	59	1	28	1617	1434	31	98	170
3	88	14	12	2425	2151	46	97	256
4	118	2	56	3234	2869	61	96	341
5	147	15	40	4042	3586	76	95	426
6	177	4	24	4851	4303	92	95	511
7	206	17	8	5659	5020	7	94	596
8	236	5	52	6468	5737	22	93	682
9	265	18	36	7276	6454	37	92	767
10	295	7	20	8085	7171	53	91	852
11	324	20	5	8893	7889	68	90	937
12	354	8	49	9702	8606	83	89	22
13	383	21	33	510	9323	98	88	108

TABLE LXXXVIII.

*Number of Days from the commencement of the year to the first of each month.*

Months.	Com.	Bis.
	Days.	Days.
January . . .	0	0
February . . .	31	31
March . . . .	59	60
April . . . .	90	91
May . . . .	120	121
June . . . .	151	152
July . . . .	181	182
August . . . .	212	213
September . . .	243	244
October . . . .	273	274
November . . .	304	305
December . . .	334	335

*Equations for New and Full Moon.*

Arg.	I		II		Arg.	I		II		Arg.	III	IV	Arg
	<i>h</i>	<i>m</i>	<i>h</i>	<i>m</i>		<i>h</i>	<i>m</i>	<i>h</i>	<i>m</i>		<i>m</i>	<i>m</i>	
0	4	20	10	10	5000	4	20	10	10	25	3	31	25
100	4	36	9	36	5100	4	5	10	50	26	3	31	24
200	4	52	9	2	5200	3	49	11	30	27	3	30	23
300	5	8	8	28	5300	3	34	12	9	28	3	30	22
400	5	24	7	55	5400	3	19	12	48	29	3	30	21
500	5	40	7	22	5500	3	4	13	26	30	3	30	20
600	5	55	6	49	5600	2	49	14	3	31	3	30	19
700	6	10	6	17	5700	2	35	14	39	32	4	30	18
800	6	24	5	46	5800	2	21	15	13	33	4	29	17
900	6	38	5	15	5900	2	8	15	46	34	4	29	16
1000	6	51	4	46	6000	1	55	16	18	35	4	29	15
1100	7	4	4	17	6100	1	42	16	48	36	5	28	14
1200	7	15	3	50	6200	1	31	17	16	37	5	28	13
1300	7	27	3	24	6300	1	19	17	42	38	5	27	12
1400	7	37	2	59	6400	1	9	18	6	39	5	27	11
1500	7	47	2	35	6500	0	59	18	28	40	6	26	10
1600	7	55	2	14	6600	0	50	18	48	41	6	26	9
1700	8	3	1	53	6700	0	42	19	6	42	7	25	8
1800	8	10	1	35	6800	0	34	19	21	43	7	25	7
1900	8	16	1	18	6900	0	28	19	33	44	7	24	6
2000	8	21	1	3	7000	0	22	19	44	45	8	23	5
2100	8	25	0	51	7100	0	17	19	52	46	8	23	4
2200	8	29	0	40	7200	0	14	19	57	47	9	22	3
2300	8	31	0	32	7300	0	11	20	0	48	9	21	2
2400	8	32	0	25	7400	0	9	20	1	49	10	21	1
2500	8	32	0	21	7500	0	8	19	59	50	10	20	0
2600	8	31	0	19	7600	0	8	19	55	51	10	19	99
2700	8	29	0	20	7700	0	9	19	48	52	11	19	98
2800	8	26	0	23	7800	0	11	19	40	53	11	18	97
2900	8	23	0	28	7900	0	15	19	29	54	12	17	96
3000	8	18	0	36	8000	0	19	19	17	55	12	17	95
3100	8	12	0	47	8100	0	24	19	2	56	13	16	94
3200	8	6	0	59	8200	0	30	18	45	57	13	15	93
3300	7	58	1	14	8300	0	37	18	27	58	13	15	92
3400	7	50	1	32	8400	0	45	18	6	59	14	14	91
3500	7	41	1	52	8500	0	53	17	45	60	14	14	90
3600	7	31	2	14	8600	1	3	17	21	61	15	13	89
3700	7	21	2	38	8700	1	13	16	56	62	15	13	88
3800	7	9	3	4	8800	1	25	16	30	63	15	12	87
3900	6	58	3	32	8900	1	36	16	3	64	15	12	86
4000	6	45	4	2	9000	1	49	15	34	65	16	11	85
4100	6	32	4	34	9100	2	2	15	5	66	16	11	84
4200	6	19	5	7	9200	2	16	14	34	67	16	11	83
4300	6	5	5	41	9300	2	30	14	3	68	16	10	82
4400	5	51	6	17	9400	2	45	13	31	69	17	10	81
4500	5	36	6	54	9500	3	0	12	58	70	17	10	80
4600	5	21	7	32	9600	3	16	12	25	71	17	10	79
4700	5	6	8	11	9700	3	32	11	52	72	17	10	78
4800	4	51	8	50	9800	3	48	11	18	73	17	10	77
4900	4	35	9	30	9900	4	4	10	44	74	17	9	76
5000	4	20	10	10	10000	4	20	10	10	75	17	9	75

*Mean Right Ascensions and Declinations of 50 principal Fixed Stars, for the beginning of 1840.*

Stars' Name.	Mag	Right Ascen.			Annual Var.	Declination.			Ann. Var.
		<i>h</i>	<i>m</i>	<i>s</i>		<i>°</i>	<i>'</i>	<i>"</i>	
1 <i>Algenib</i>	2.3	0	5	0.31	+ 3.0775	14	17	38.82 N	+ 20.051
2 <i>β Andromedae</i>	2	1	0	46.7	3.309	34	46	17.2 N	19.35
3 <i>Polaris</i>	2.3	1	2	10.38	16.1962	88	27	21.96 N	19.339
4 <i>Achernar</i>	1	1	31	44.88	2.2351	58	3	5.13 S	— 18.473
5 <i>α Arietis</i>	3	1	58	9.94	3.3457	22	42	11.81 N	+ 17.456
6 <i>α Ceti</i>	2.3	2	53	55.34	+ 3.1257	3	27	30.09 N	+ 14.561
7 <i>α Persei</i>	2.3	3	12	55.97	4.2280	49	17	8.74 N	13.371
8 <i>Aldebaran</i>	1	4	26	44.77	3.4264	16	10	56.82 N	7.949
9 <i>Capella</i>	1	5	4	52.67	4.4066	45	49	42.81 N	4.793
10 <i>Rigel</i>	1	5	6	51.09	2.8783	8	23	29.29 S	— 4.620
11 <i>β Tauri</i>	2	5	16	10.96	+ 3.7820	28	27	58.20 N	+ 3.825
12 <i>γ Orionis</i>	2	5	16	33.1	3.210	6	11	55.3 N	+ 3.82
13 <i>α Columbae</i>	2	5	33	51.52	2.1688	34	9	47.41 S	— 2.291
14 <i>α Orionis</i>	1	5	46	30.71	3.2430	7	22	17.14 N	+ 1.191
15 <i>Canopus</i>	1	6	20	24.18	1.3278	52	36	38.42 S	1.778
16 <i>Sirius</i>	1	6	38	5.76	+ 2.6458	16	30	4.79 S	+ 4.449
17 <i>Castor</i>	3	7	24	23.06	3.8572	32	13	58.89 N	— 7.206
18 <i>Procyon</i>	1.2	7	30	55.53	3.1448	5	37	48.92 N	8.720
19 <i>Pollux</i>	2	7	35	31.07	3.6840	28	24	25.57 N	8.107
20 <i>α Hydrae</i>	2	9	19	43.57	2.9500	7	58	4.83 S	+ 15.341
21 <i>Regulus</i>	1	9	59	50.93	+ 3.2220	12	44	49.70 N	— 17.356
22 <i>α Ursae Majoris</i>	1.2	10	53	47.98	3.8077	62	36	48.93 N	19.221
23 <i>β Leonis</i>	2.3	11	40	53.69	3.0660	15	28	1.16 N	19.985
24 <i>β Virginis</i>	3.4	11	42	21.4	3.124	2	40	2.6 N	19.98
25 <i>γ Ursae Majoris</i>	2	11	45	22.93	3.1914	54	35	4.67 N	20.014
26 <i>α 2 Crucis</i>	2	12	17	43.7	+ 3.258	62	12	47.9 S	+ 19.99
27 <i>Spica</i>	1	13	16	46.36	3.1502	10	19	24.39 S	18.945
28 <i>θ Centauri</i>	2	13	57	18.0	3.491	35	34	41.9 S	17.499
29 <i>α Draconis</i>	3.4	14	0	2.8	1.625	65	8	32.1 N	— 17.37
30 <i>Arcturus</i>	1	14	8	21.96	2.7335	20	1	7.67 N	18.956
31 <i>α 2 Centauri</i>	1	14	28	47.84	+ 4.0086	60	10	6.24 S	+ 15.152
32 <i>α 2 Librae</i>	3	14	42	2.44	3.3088	15	22	18.25 S	15.256
33 <i>β Ursae Minoris</i>	3	14	51	14.66	— 0.2787	74	48	34.18 N	— 14.712
34 <i>γ 2 Ursae Minoris</i>	3.4	15	21	1.3	— 0.179	72	24	14.1 N	12.81
35 <i>α Coronae Borealis</i>	2	15	27	54.87	+ 2.5277	27	15	27.71 N	12.361
36 <i>α Serpentis</i>	2.3	15	36	23.43	+ 2.9386	6	56	2.80 N	— 11.770
37 <i>β Scorpii</i>	2	15	56	8.68	3.4729	19	21	38.82 S	+ 10.330
38 <i>Antares</i>	1	16	19	36.49	3.6625	26	4	13.13 S	8.519
39 <i>α Herculis</i>	3.4	17	7	21.30	2.7317	14	34	41.43 N	— 4.576
40 <i>α Ophiuchi</i>	2	17	27	30.56	2.7724	12	40	58.65 N	2.844
41 <i>δ Ursae Minoris</i>	3	18	23	56.48	— 19.2072	86	35	28.89 N	+ 2.161
42 <i>Vega</i>	1	18	31	31.19	+ 2.0116	38	38	16.85 N	2.742
43 <i>Altair</i>	1	19	42	58.61	2.9255	8	27	0.21 N	8.701
44 <i>α 2 Capricorni</i>	3	20	9	10.34	3.3323	13	2	5.57 S	— 10.705
45 <i>α Cygni</i>	1	20	35	58.80	2.0416	44	42	41.38 N	+ 12.614
46 <i>α Aquarii</i>	3	21	57	33.93	+ 3.0835	1	5	38.00 S	— 17.256
47 <i>Fomalhaut</i>	1	22	48	47.67	3.3114	30	28	4.91 S	19.092
48 <i>β Pegasi</i>	2	22	56	1.1	2.878	27	13	1.7 N	+ 19.255
49 <i>Markab</i>	2	22	56	47.75	2.9771	14	20	46.92 N	19.295
50 <i>α Andromedae</i>	1	24	0	7.72	3.0704	28	12	27.06 N	20.056

*Constants for the Aberration and Nutation in Right Ascension and Declination of the Stars in the preceding Catalogue*

	Aberration.				Nutation.			
	$\phi$	M	$\theta$	N	$\phi'$	M'	$\theta'$	N'
1	<sup>s</sup> 8 28 47	0.1087	<sup>s</sup> 7 27 12	0.9657	<sup>s</sup> 6 8 24	0.0300	<sup>s</sup> 5 28 30	0.8381
2	8 13 39	0.1830	6 19 12	1.0740	6 19 53	0.0838	5 10 8	0.8496
3	8 13 51	1.6526	5 16 57	1.3052	8 16 7	1.3427	5 10 22	0.8493
4	8 5 20	0.3801	10 26 46	1.2798	4 10 12	0.0775	5 0 31	0.8629
5	7 28 26	0.1397	7 0 2	0.8972	6 11 1	0.0695	4 22 53	0.8765
6	7 14 11	0.1149	8 23 8	0.8678	6 1 26	0.0322	4 8 16	0.9078
7	7 9 30	0.3020	5 3 5	1.0630	6 18 13	0.1849	4 3 47	0.9179
8	6 21 43	0.1447	7 23 12	0.5760	6 3 27	0.0726	3 17 54	0.9502
9	6 12 51	0.2875	3 25 37	0.9112	6 5 46	0.1830	3 10 29	0.9605
10	6 12 20	0.1355	9 3 42	1.0300	5 28 47	1.9966	3 10 4	0.9608
11	6 10 13	0.1873	4 19 21	0.3917	6 2 52	0.1008	3 8 19	0.9626
12	6 10 6	0.1340	8 26 4	0.7851	6 0 40	0.0441	3 8 14	0.9626
13	6 6 5	0.2145	9 4 24	1.2348	5 26 18	1.8750	3 4 57	0.9648
14	6 3 13	0.1361	8 28 23	0.7521	6 0 15	0.0481	3 2 37	0.9657
15	5 25 22	0.3491	8 25 53	1.2960	6 8 46	1.6679	2 26 15	0.9657
16	5 21 21	0.1501	8 25 51	1.1152	6 1 51	1.9658	2 22 58	0.9636
17	5 10 40	0.2010	1 2 17	0.6620	5 24 2	0.1257	2 14 6	0.9535
18	5 9 6	0.1297	9 6 54	0.8071	5 28 47	0.0414	2 12 47	0.9513
19	5 8 2	0.1829	0 14 32	0.6052	5 24 2	0.1114	2 11 53	0.9499
20	4 12 39	0.1158	8 17 31	0.9967	6 3 41	0.0081	1 18 37	0.9007
21	4 2 22	0.1162	10 3 47	0.8457	5 23 47	0.0480	1 7 59	0.8782
22	3 18 7	0.4366	0 3 28	1.2394	4 18 58	0.2407	0 21 57	0.8520
23	3 5 21	0.1117	10 6 20	0.9621	5 20 56	0.0344	0 6 35	0.8393
24	3 4 57	0.0958	9 6 51	0.9075	5 28 25	0.0253	0 6 5	0.8390
25	3 4 8	0.3229	11 17 28	1.2298	4 21 46	0.1465	0 5 5	0.8388
26	2 25 19	0.4261	6 8 5	1.2585	7 16 2	0.2089	11 24 14	0.8390
27	2 9 22	0.1066	8 3 31	0.8862	6 5 51	0.0154	11 5 6	0.8559
28	1 28 40	0.1942	6 7 12	1.0176	6 17 31	0.1062	10 23 8	0.8760
29	1 27 53	0.4824	10 23 28	1.2995	3 25 50	0.1090	10 22 16	0.8777
30	1 25 46	0.1336	9 28 18	1.0974	5 18 49	1.9937	10 20 1	0.8822
31	1 20 32	0.4123	5 7 54	1.1820	6 29 6	0.2460	10 14 36	0.8937
32	1 17 26	0.1273	7 18 24	0.6923	6 6 29	0.0593	10 11 28	0.9006
33	1 14 42	0.6961	10 15 5	1.3087	2 26 45	0.2235	10 8 47	0.9066
34	1 7 20	0.6386	10 7 33	1.3087	2 27 7	0.0960	10 1 45	0.9225
35	1 5 45	0.1704	9 22 28	1.1785	5 17 18	1.9510	10 0 18	0.9257
36	1 3 43	0.1237	9 8 22	0.9994	5 27 30	0.0058	9 28 26	0.9298
37	0 28 58	0.1485	7 4 4	0.6237	6 5 20	0.0795	9 24 12	0.9386
38	0 23 24	0.1728	5 27 59	0.5816	6 5 49	0.1029	9 19 21	0.9478
39	0 12 13	0.1451	9 5 25	1.0962	5 27 45	1.9742	9 9 58	0.9610
40	0 7 34	0.1427	9 3 4	1.0786	5 28 48	1.9803	9 6 9	0.9642
41	11 23 47	1.3571	8 22 49	1.2321	11 19 31	0.8257	8 24 57	0.9650
42	11 22 50	0.2393	8 24 29	1.2545	6 5 31	1.8436	8 24 10	0.9644
43	11 6 15	0.1309	8 22 59	1.0237	6 2 16	1.9988	8 10 21	0.9472
44	11 0 2	0.1341	9 29 33	0.6961	5 26 12	0.0609	8 4 55	0.9368
45	10 23 29	0.2668	8 0 39	1.2634	6 28 32	1.9042	7 29 0	0.9242
46	10 2 57	0.1057	9 2 31	0.8988	5 29 26	0.0264	7 8 37	0.8794
47	9 19 26	0.1638	11 7 34	1.0271	5 13 8	0.0765	6 23 30	0.8540
48	9 17 29	0.1491	7 17 0	1.1171	6 17 2	0.0162	6 21 13	0.8511
49	9 17 17	0.1120	8 2 5	1.0138	6 8 23	0.0157	6 20 58	0.8508
50	9 0 6	0.1495	7 6 42	1.0785	6 17 20	0.0444	6 0 8	0.8380



*Mean Longitudes and Latitudes of some of the principal Fixed Stars for the beginning of 1840, with their Annual Variations.*

Stars' Name.	Mag	Longitude.				Annual Var.	Latitude.				Annual Var.
		s	°	'	"	"	°	'	"	"	"
$\alpha$ Arietis	3	1	5	25	27.6	50.277	9	57	40.9	N	+ 0.161
Aldebaran	1	2	7	33	5.9	50.210	5	28	38.0	S	— 0.335
Capella	1	2	19	37	17.8	50.302	22	51	44.4	N	— 0.052
Polaris	2.3	2	26	19	20.1	47.959	66	4	59.5	N	+ 0.552
Sirius	1	3	11	52	32.9	49.488	39	34	4.3	S	+ 0.319
Canopus	1	3	12	44	59.6	49.366	75	50	57.6	S	+ 0.459
Pollux	2	3	21	0	22.0	49.502	6	40	20.2	N	+ 0.255
Regulus	1	4	27	36	13.2	49.946	0	27	38.3	N	+ 0.220
Spica	1	6	21	36	29.2	50.085	2	2	29.7	S	+ 0.171
Arcturus	1	6	22	0	4.7	50.711	30	51	17.5	N	+ 0.214
Antares	1	8	7	31	45.2	50.120	4	32	51.6	S	+ 0.424
Altair	1.2	9	29	31	5.9	50.795	29	18	37.3	N	+ 0.080
Fomalhaut	1	11	1	36	22.0	50.595	21	6	49.7	S	+ 0.213
Achernar	1	11	13	2	5.3	50.346	17	6	17.3	S	— 0.083
$\alpha$ Pegasi	2	11	21	15	24.7	50.112	19	24	40.9	N	+ 0.098

TABLE added to TABLE XC.

*Mean Right Ascensions and Declinations of Polaris and  $\delta$  Ursae Minoris for 1830, 1840, 1850, and 1860.*

Stars.	Years	Right Asc.			Ann. Var.	Declination.			Ann. Var.
		°	'	"	"	°	'	"	"
Polaris	1830	0	59	30.76	+ 15.478	88	24	8.82	+ 19.371
	1840	1	2	10.32	16.470	88	27	22.43	19.309
	1850	1	5	0.29	17.567	88	30	35.40	19.240
	1860	1	8	1.79	18.784	88	33	47.64	19.163
$\delta$ Ursae Minoris	1830	18	27	5.13	— 19.167	86	35	5.70	+ 2.363
	1840	18	23	53.03	19.241	86	35	27.93	2.085
	1850	18	20	40.21	19.305	86	35	47.36	1.805
	1860	18	17	26.77	19.360	86	36	3.97	1.523

*Second Differences.*

Hours & Minutes.				1'	2'	3'	4'	5'	6'	7'	8'	9'	10'	11'
<i>h</i>	<i>m</i>	<i>h</i>	<i>m</i>	"	"	"	"	"	"	"	"	"	"	"
0	0	12	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
0	10	11	50	0.4	0.8	1.2	1.6	2.0	2.4	2.9	3.3	3.7	4.1	4.5
0	20	11	40	0.8	1.6	2.4	3.2	4.1	4.9	5.7	6.5	7.3	8.1	8.9
0	30	11	30	1.2	2.4	3.6	4.8	6.0	7.2	8.4	9.6	10.8	12.0	13.2
0	40	11	20	1.6	3.1	4.7	6.3	7.9	9.4	11.0	12.6	14.2	15.7	17.3
0	50	11	10	1.9	3.9	5.8	7.8	9.7	11.6	13.6	15.5	17.4	19.4	21.4
1	0	11	0	2.3	4.6	6.9	9.2	11.5	13.8	16.0	18.3	20.6	22.9	25.2
1	10	10	50	2.6	5.3	7.9	10.5	13.2	15.8	18.4	21.1	23.7	26.3	29.0
1	20	10	40	3.0	5.9	8.9	11.9	14.8	17.8	20.7	23.7	26.7	29.6	32.6
1	30	10	30	3.3	6.6	9.8	13.1	16.4	19.7	23.0	26.3	29.5	32.8	36.1
1	40	10	20	3.6	7.2	10.8	14.4	17.9	21.5	25.1	28.7	32.3	35.9	39.5
1	50	10	10	3.9	7.8	11.6	15.5	19.4	23.3	27.2	31.0	34.9	38.8	42.7
2	0	10	0	4.2	8.3	12.5	16.7	20.8	25.0	29.2	33.3	37.5	41.7	45.8
2	10	9	50	4.4	8.9	13.3	17.8	22.2	26.6	31.1	35.5	40.0	44.4	48.8
2	20	9	40	4.7	9.4	14.1	18.8	23.5	28.2	32.9	37.6	42.3	47.0	51.7
2	30	9	30	4.9	9.9	14.8	19.8	24.7	29.7	34.6	39.6	44.5	49.5	54.4
2	40	9	20	5.2	10.4	15.6	20.7	25.9	31.1	36.3	41.5	46.7	51.9	57.0
2	50	9	10	5.4	10.8	16.2	21.6	27.1	32.5	37.9	43.3	48.7	54.1	59.5
3	0	9	0	5.6	11.3	16.9	22.5	28.1	33.8	39.4	45.0	50.6	56.3	61.9
3	10	8	50	5.8	11.7	17.5	23.3	29.1	35.0	40.8	46.6	52.4	58.3	64.1
3	20	8	40	6.0	12.0	18.1	24.1	30.1	36.1	42.1	48.1	54.2	60.2	66.2
3	30	8	30	6.2	12.4	18.6	24.8	31.0	37.2	43.4	49.6	55.8	62.0	68.2
3	40	8	20	6.4	12.7	19.1	25.5	31.8	38.2	44.6	50.9	57.3	63.7	70.0
3	50	8	10	6.5	13.0	19.6	26.1	32.6	39.1	45.7	52.2	58.7	65.2	71.7
4	0	8	0	6.7	13.3	20.0	26.7	33.3	40.0	46.7	53.3	60.0	66.7	73.3
4	10	7	50	6.8	13.6	20.4	27.2	34.0	40.8	47.6	54.4	61.2	68.0	74.8
4	20	7	40	6.9	13.8	20.8	27.7	34.6	41.5	48.4	55.4	62.3	69.2	76.1
4	30	7	30	7.0	14.1	21.1	28.1	35.2	42.2	49.2	56.2	63.3	70.3	77.3
4	40	7	20	7.1	14.3	21.4	28.5	35.6	42.8	49.9	57.0	64.2	71.3	78.4
4	50	7	10	7.2	14.4	21.6	28.9	36.1	43.3	50.5	57.7	64.9	72.2	79.4
5	0	7	0	7.3	14.6	21.9	29.2	36.5	43.8	51.0	58.3	65.6	72.9	80.2
5	10	6	50	7.4	14.7	22.1	29.4	36.8	44.1	51.5	58.8	66.2	73.6	80.9
5	20	6	40	7.4	14.8	22.2	29.6	37.0	44.4	51.9	59.3	66.7	74.1	81.5
5	30	6	30	7.4	14.9	22.3	29.8	37.2	44.7	52.1	59.6	67.0	74.5	81.9
5	40	6	20	7.5	15.0	22.4	29.9	37.4	44.9	52.3	59.8	67.3	74.8	82.2
5	50	6	10	7.5	15.0	22.5	30.0	37.5	45.0	52.5	60.0	67.4	74.9	82.4
6	0	6	0	7.5	15.0	22.5	30.0	37.5	45.0	52.5	60.0	67.5	75.0	82.5

*Second Differences.*

Hours & Min.		10"	20"	30"	40"	50"	1"	2"	3"	4"	5"	6"	7"	8"	9"
<i>h</i>	<i>m</i>	<i>h</i>	<i>m</i>	"	"	"	"	"	"	"	"	"	"	"	"
0	0	12	0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
0	10	11	50	0.1	0.1	0.2	0.3	0.3	0.0	0.0	0.0	0.0	0.0	0.1	0.1
0	20	11	40	0.1	0.3	0.4	0.5	0.7	0.0	0.0	0.0	0.1	0.1	0.1	0.1
0	30	11	30	0.2	0.4	0.6	0.8	1.0	0.0	0.0	0.1	0.1	0.1	0.1	0.2
0	40	11	20	0.3	0.5	0.8	1.0	1.3	0.0	0.1	0.1	0.1	0.1	0.2	0.2
0	50	11	10	0.3	0.6	1.0	1.3	1.6	0.0	0.1	0.1	0.1	0.2	0.2	0.3
1	0	11	0	0.4	0.8	1.1	1.5	1.9	0.0	0.1	0.1	0.2	0.2	0.3	0.3
1	10	10	50	0.4	0.9	1.3	1.8	2.2	0.0	0.1	0.1	0.2	0.2	0.3	0.4
1	20	10	40	0.5	1.0	1.5	2.0	2.5	0.0	0.1	0.1	0.2	0.2	0.3	0.4
1	30	10	30	0.5	1.1	1.6	2.2	2.7	0.1	0.1	0.2	0.2	0.3	0.4	0.5
1	40	10	20	0.6	1.2	1.8	2.4	3.0	0.1	0.1	0.2	0.2	0.3	0.4	0.5
1	50	10	10	0.6	1.3	1.9	2.6	3.2	0.1	0.1	0.2	0.3	0.4	0.5	0.6
2	0	10	0	0.7	1.4	2.1	2.8	3.5	0.1	0.1	0.2	0.3	0.4	0.5	0.6
2	10	9	50	0.7	1.5	2.2	3.0	3.7	0.1	0.1	0.2	0.3	0.4	0.5	0.7
2	20	9	40	0.8	1.6	2.3	3.1	3.9	0.1	0.2	0.2	0.3	0.4	0.5	0.7
2	30	9	30	0.8	1.6	2.5	3.3	4.1	0.1	0.2	0.2	0.3	0.4	0.5	0.7
2	40	9	20	0.9	1.7	2.6	3.5	4.3	0.1	0.2	0.3	0.3	0.4	0.5	0.8
2	50	9	10	0.9	1.8	2.7	3.6	4.5	0.1	0.2	0.3	0.4	0.5	0.6	0.8
3	0	9	0	0.9	1.9	2.8	3.8	4.7	0.1	0.2	0.3	0.4	0.5	0.6	0.8
3	10	8	50	1.0	1.9	2.9	3.9	4.9	0.1	0.2	0.3	0.4	0.5	0.6	0.9
3	20	8	40	1.0	2.0	3.0	4.0	5.0	0.1	0.2	0.3	0.4	0.5	0.6	0.9
3	30	8	30	1.0	2.1	3.1	4.1	5.2	0.1	0.2	0.3	0.4	0.5	0.6	0.9
3	40	8	20	1.1	2.1	3.2	4.2	5.3	0.1	0.2	0.3	0.4	0.5	0.6	1.0
3	50	8	10	1.1	2.2	3.3	4.3	5.4	0.1	0.2	0.3	0.4	0.5	0.7	1.0
4	0	8	0	1.1	2.2	3.3	4.4	5.6	0.1	0.2	0.3	0.4	0.6	0.7	1.0
4	10	7	50	1.1	2.3	3.4	4.5	5.7	0.1	0.2	0.3	0.5	0.6	0.7	1.0
4	20	7	40	1.2	2.3	3.5	4.6	5.8	0.1	0.2	0.3	0.5	0.6	0.7	1.0
4	30	7	30	1.2	2.3	3.5	4.7	5.9	0.1	0.2	0.4	0.5	0.6	0.7	1.1
4	40	7	20	1.2	2.4	3.6	4.8	5.9	0.1	0.2	0.4	0.5	0.6	0.7	1.1
4	50	7	10	1.2	2.4	3.6	4.8	6.0	0.1	0.2	0.4	0.5	0.6	0.7	1.1
5	0	7	0	1.2	2.4	3.6	4.9	6.1	0.1	0.2	0.4	0.5	0.6	0.7	1.1
5	10	6	50	1.2	2.5	3.7	4.9	6.1	0.1	0.2	0.4	0.5	0.6	0.7	1.1
5	20	6	40	1.2	2.5	3.7	4.9	6.1	0.1	0.2	0.4	0.5	0.6	0.7	1.1
5	30	6	30	1.2	2.5	3.7	5.0	6.2	0.1	0.2	0.4	0.5	0.6	0.7	1.1
5	40	6	20	1.2	2.5	3.7	5.0	6.2	0.1	0.2	0.4	0.5	0.6	0.7	1.1
5	50	6	10	1.2	2.5	3.7	5.0	6.2	0.1	0.2	0.4	0.5	0.6	0.7	1.1
6	0	6	0	1.3	2.6	3.8	5.0	6.3	0.1	0.2	0.4	0.5	0.6	0.7	1.1

*Third Differences.*

Time after noon or midnight.	10"	20"	30"	40"	50"	1'	2'	3'	4'	5'	Time after noon or midnight.
+	"	"	"	"	"	"	"	"	"	"	—
0h. 0m.	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	12h. 0m.
0 30	0.0	0.1	0.1	0.1	0.2	0.2	0.4	0.5	0.7	0.9	11 30
1 0	0.1	0.1	0.2	0.2	0.3	0.3	0.6	1.0	1.3	1.5	11 0
1 30	0.1	0.1	0.2	0.3	0.3	0.4	0.8	1.2	1.6	2.1	10 30
2 0	0.1	0.2	0.2	0.3	0.4	0.5	0.9	1.4	1.9	2.3	10 0
2 30	0.1	0.2	0.2	0.3	0.4	0.5	1.0	1.4	1.9	2.4	9 30
3 0	0.1	0.2	0.2	0.3	0.4	0.5	0.9	1.4	1.9	2.3	9 0
3 30	0.1	0.1	0.2	0.3	0.4	0.4	0.9	1.3	1.7	2.2	8 30
4 0	0.1	0.1	0.2	0.2	0.3	0.4	0.7	1.1	1.5	1.9	8 0
4 30	0.0	0.1	0.1	0.2	0.2	0.3	0.6	0.9	1.2	1.5	7 30
5 0	0.0	0.1	0.1	0.1	0.2	0.2	0.4	0.6	0.8	1.0	7 0
5 30	0.0	0.0	0.1	0.1	0.1	0.1	0.2	0.3	0.4	0.5	6 30
6 0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	6 0
+											—

TABLE XCV.

*Fourth Differences.*

Time after noon or midnight.	10"	20"	30"	40"	50"	1'	2'	3'	Time after noon or midnight.
h. m.	"	"	"	"	"	"	"	"	h. m.
0 0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	12 0
0 30	0.0	0.1	0.1	0.1	0.2	0.2	0.4	0.6	11 30
1 0	0.1	0.1	0.2	0.3	0.3	0.4	0.8	1.2	11 0
1 30	0.1	0.2	0.3	0.4	0.5	0.6	1.2	1.7	10 30
2 0	0.1	0.2	0.4	0.5	0.6	0.7	1.5	2.2	10 0
2 30	0.1	0.3	0.4	0.6	0.7	0.9	1.8	2.7	9 30
3 0	0.2	0.3	0.5	0.7	0.9	1.0	2.1	3.1	9 0
3 30	0.2	0.4	0.6	0.8	0.9	1.1	2.3	3.4	8 30
4 0	0.2	0.4	0.6	0.8	1.0	1.2	2.5	3.7	8 0
4 30	0.2	0.4	0.7	0.9	1.1	1.3	2.6	3.9	7 30
5 0	0.2	0.5	0.7	0.9	1.1	1.4	2.7	4.1	7 0
5 30	0.2	0.5	0.7	0.9	1.2	1.4	2.8	4.2	6 30
6 0	0.2	0.5	0.7	0.9	1.2	1.4	2.8	4.2	6 0

	0	1	2	3	4	5	6	7	8	9
"	0	60	120	180	240	300	360	420	480	540
0		1.7782	1.4771	1.3010	1.1761	1.0792	1.0000	9331	8751	8239
1	3.5563	1.7710	1.4735	1.2986	1.1743	1.0777	9988	9320	8742	8231
2	3.2553	1.7639	1.4699	1.2962	1.1725	1.0763	9976	9310	8733	8223
3	3.0792	1.7570	1.4664	1.2939	1.1707	1.0749	9964	9300	8724	8215
4	2.9542	1.7501	1.4629	1.2915	1.1689	1.0734	9952	9289	8715	8207
5	2.8573	1.7434	1.4594	1.2891	1.1671	1.0720	9940	9279	8706	8199
6	2.7782	1.7368	1.4559	1.2868	1.1654	1.0706	9928	9269	8697	8191
7	2.7112	1.7302	1.4525	1.2845	1.1636	1.0692	9916	9259	8688	8183
8	2.6532	1.7238	1.4491	1.2821	1.1619	1.0678	9905	9249	8679	8175
9	2.6021	1.7175	1.4457	1.2798	1.1601	1.0663	9893	9238	8670	8167
10	2.5563	1.7112	1.4424	1.2775	1.1584	1.0649	9881	9228	8661	8159
11	2.5149	1.7050	1.4390	1.2753	1.1566	1.0635	9869	9218	8652	8152
12	2.4771	1.6990	1.4357	1.2730	1.1549	1.0621	9858	9208	8643	8144
13	2.4424	1.6930	1.4325	1.2707	1.1532	1.0608	9846	9198	8635	8136
14	2.4102	1.6871	1.4292	1.2685	1.1515	1.0594	9834	9188	8626	8128
15	2.3802	1.6812	1.4260	1.2663	1.1498	1.0580	9823	9178	8617	8120
16	2.3522	1.6755	1.4228	1.2640	1.1481	1.0566	9811	9168	8608	8112
17	2.3259	1.6698	1.4196	1.2618	1.1464	1.0552	9800	9158	8599	8104
18	2.3010	1.6642	1.4165	1.2596	1.1447	1.0539	9788	9148	8591	8097
19	2.2775	1.6587	1.4133	1.2574	1.1430	1.0525	9777	9138	8582	8089
20	2.2553	1.6532	1.4102	1.2553	1.1413	1.0512	9765	9128	8573	8081
21	2.2341	1.6478	1.4071	1.2531	1.1397	1.0498	9754	9119	8565	8073
22	2.2139	1.6425	1.4040	1.2510	1.1380	1.0484	9742	9109	8556	8066
23	2.1946	1.6372	1.4010	1.2488	1.1363	1.0471	9731	9099	8547	8058
24	2.1761	1.6320	1.3979	1.2467	1.1347	1.0458	9720	9089	8539	8050
25	2.1584	1.6269	1.3949	1.2445	1.1331	1.0444	9708	9079	8530	8043
26	2.1413	1.6218	1.3919	1.2424	1.1314	1.0431	9697	9070	8522	8035
27	2.1249	1.6168	1.3890	1.2403	1.1298	1.0418	9686	9060	8513	8027
28	2.1091	1.6118	1.3860	1.2382	1.1282	1.0404	9675	9050	8504	8020
29	2.0939	1.6069	1.3831	1.2362	1.1266	1.0391	9664	9041	8496	8012
30	2.0792	1.6021	1.3802	1.2341	1.1249	1.0378	9652	9031	8487	8004
31	2.0649	1.5973	1.3773	1.2320	1.1233	1.0365	9641	9021	8479	7997
32	2.0512	1.5925	1.3745	1.2300	1.1217	1.0352	9630	9012	8470	7989
33	2.0378	1.5878	1.3716	1.2279	1.1201	1.0339	9619	9002	8462	7981
34	2.0248	1.5832	1.3688	1.2259	1.1186	1.0326	9608	8992	8453	7974
35	2.0122	1.5786	1.3660	1.2239	1.1170	1.0313	9597	8983	8445	7966
36	2.0000	1.5740	1.3632	1.2218	1.1154	1.0300	9586	8973	8437	7959
37	1.9881	1.5695	1.3604	1.2198	1.1138	1.0287	9575	8964	8428	7951
38	1.9765	1.5651	1.3576	1.2178	1.1123	1.0274	9564	8954	8420	7944
39	1.9652	1.5607	1.3549	1.2159	1.1107	1.0261	9553	8945	8411	7936
40	1.9542	1.5563	1.3522	1.2139	1.1091	1.0248	9542	8935	8403	7929
41	1.9435	1.5520	1.3495	1.2119	1.1076	1.0235	9532	8926	8395	7921
42	1.9331	1.5477	1.3468	1.2099	1.1061	1.0223	9521	8917	8386	7914
43	1.9228	1.5435	1.3441	1.2080	1.1045	1.0210	9510	8907	8378	7906
44	1.9128	1.5393	1.3415	1.2061	1.1030	1.0197	9499	8898	8370	7899
45	1.9031	1.5351	1.3388	1.2041	1.1015	1.0185	9488	8888	8361	7891
46	1.8935	1.5310	1.3362	1.2022	1.0999	1.0172	9478	8879	8353	7884
47	1.8842	1.5269	1.3336	1.2003	1.0984	1.0160	9467	8870	8345	7877
48	1.8751	1.5229	1.3310	1.1984	1.0969	1.0147	9456	8861	8337	7869
49	1.8661	1.5189	1.3284	1.1965	1.0954	1.0135	9446	8851	8328	7862
50	1.8573	1.5149	1.3259	1.1946	1.0939	1.0122	9435	8842	8320	7855
51	1.8487	1.5110	1.3233	1.1927	1.0924	1.0110	9425	8833	8312	7847
52	1.8403	1.5071	1.3208	1.1908	1.0909	1.0098	9414	8824	8304	7840
53	1.8320	1.5032	1.3183	1.1889	1.0894	1.0085	9404	8814	8296	7832
54	1.8239	1.4994	1.3158	1.1871	1.0880	1.0073	9393	8805	8288	7825
55	1.8159	1.4956	1.3133	1.1852	1.0865	1.0061	9383	8796	8279	7818
56	1.8081	1.4918	1.3108	1.1834	1.0850	1.0049	9372	8787	8271	7811
57	1.8004	1.4881	1.3083	1.1816	1.0835	1.0036	9362	8778	8263	7803
58	1.7929	1.4844	1.3059	1.1797	1.0821	1.0024	9351	8769	8255	7796
59	1.7855	1.4808	1.3034	1.1779	1.0806	1.0012	9341	8760	8247	7789
60	1.7782	1.4771	1.3010	1.1761	1.0792	1.0000	9331	8751	8239	7782




	10	11	12	13	14	15	16	17	18	19	20	21
"	600	660	720	780	840	900	960	1020	1080	1140	1200	1260
0	7782	7368	6990	6642	6320	6021	5740	5477	5229	4994	4771	4559
1	7774	7361	6984	6637	6315	6016	5736	5473	5225	4990	4768	4556
2	7767	7354	6978	6631	6310	6011	5731	5469	5221	4986	4764	4552
3	7760	7348	6972	6625	6305	6006	5727	5464	5217	4983	4760	4549
4	7753	7341	6966	6620	6300	6001	5722	5460	5213	4979	4757	4546
5	7745	7335	6960	6614	6294	5997	5718	5456	5209	4975	4753	4542
6	7738	7328	6954	6609	6289	5992	5713	5452	5205	4971	4750	4539
7	7731	7322	6948	6603	6284	5987	5709	5447	5201	4967	4746	4535
8	7724	7315	6942	6598	6279	5982	5704	5443	5197	4964	4742	4532
9	7717	7309	6936	6592	6274	5977	5700	5439	5193	4960	4739	4528
10	7710	7302	6930	6587	6269	5973	5695	5435	5189	4956	4735	4525
11	7703	7296	6924	6581	6264	5968	5691	5430	5185	4952	4732	4522
12	7696	7289	6918	6576	6259	5963	5686	5426	5181	4949	4728	4518
13	7688	7283	6912	6570	6254	5958	5682	5422	5177	4945	4724	4515
14	7681	7276	6906	6565	6248	5954	5677	5418	5173	4941	4721	4511
15	7674	7270	6900	6559	6243	5949	5673	5414	5169	4937	4717	4508
16	7667	7264	6894	6554	6238	5944	5669	5409	5165	4933	4714	4505
17	7660	7257	6888	6548	6233	5939	5664	5405	5161	4930	4710	4501
18	7653	7251	6882	6543	6228	5935	5660	5401	5157	4926	4707	4498
19	7646	7244	6877	6538	6223	5930	5655	5397	5153	4922	4703	4494
20	7639	7238	6871	6532	6218	5925	5651	5393	5149	4918	4699	4491
21	7632	7232	6865	6527	6213	5920	5646	5389	5145	4915	4696	4488
22	7625	7225	6859	6521	6208	5916	5642	5384	5141	4911	4692	4484
23	7618	7219	6853	6516	6203	5911	5637	5380	5137	4907	4689	4481
24	7611	7212	6847	6510	6198	5906	5633	5376	5133	4903	4685	4477
25	7604	7206	6841	6505	6193	5902	5629	5372	5129	4900	4682	4474
26	7597	7200	6836	6500	6188	5897	5624	5368	5125	4896	4678	4471
27	7590	7193	6830	6494	6183	5892	5620	5364	5122	4892	4675	4467
28	7583	7187	6824	6489	6178	5888	5615	5359	5118	4889	4671	4464
29	7577	7181	6818	6484	6173	5883	5611	5355	5114	4885	4668	4460
30	7570	7175	6812	6478	6168	5878	5607	5351	5110	4881	4664	4457
31	7563	7168	6807	6473	6163	5874	5602	5347	5106	4877	4660	4454
32	7556	7162	6801	6467	6158	5869	5598	5343	5102	4874	4657	4450
33	7549	7156	6795	6462	6153	5864	5594	5339	5098	4870	4653	4447
34	7542	7149	6789	6457	6148	5860	5589	5335	5094	4866	4650	4444
35	7535	7143	6784	6451	6143	5855	5585	5331	5090	4863	4646	4440
36	7528	7137	6778	6446	6138	5850	5580	5326	5086	4859	4643	4437
37	7522	7131	6772	6441	6133	5846	5576	5322	5082	4855	4639	4434
38	7515	7124	6766	6435	6128	5841	5572	5318	5079	4852	4636	4430
39	7508	7118	6761	6430	6123	5836	5567	5314	5075	4848	4632	4427
40	7501	7112	6755	6425	6118	5832	5563	5310	5071	4844	4629	4424
41	7494	7106	6749	6420	6113	5827	5559	5306	5067	4841	4625	4420
42	7488	7100	6743	6414	6108	5823	5554	5302	5063	4837	4622	4417
43	7481	7093	6738	6409	6103	5818	5550	5298	5059	4833	4618	4414
44	7474	7087	6732	6404	6099	5813	5546	5294	5055	4830	4615	4410
45	7467	7081	6726	6398	6094	5809	5541	5290	5051	4826	4611	4407
46	7461	7075	6721	6393	6089	5804	5537	5285	5048	4822	4608	4404
47	7454	7069	6715	6388	6084	5800	5533	5281	5044	4819	4604	4400
48	7447	7063	6709	6383	6079	5795	5528	5277	5040	4815	4601	4397
49	7441	7057	6704	6377	6074	5790	5524	5273	5036	4811	4597	4394
50	7434	7050	6698	6372	6069	5786	5520	5269	5032	4808	4594	4390
51	7427	7044	6692	6367	6064	5781	5516	5265	5028	4804	4590	4387
52	7421	7038	6687	6362	6059	5777	5511	5261	5025	4800	4587	4384
53	7414	7032	6681	6357	6055	5772	5507	5257	5021	4797	4584	4380
54	7407	7026	6676	6351	6050	5768	5503	5253	5017	4793	4580	4377
55	7401	7020	6670	6346	6045	5763	5498	5249	5013	4789	4577	4374
56	7394	7014	6664	6341	6040	5758	5494	5245	5009	4786	4573	4370
57	7387	7008	6659	6336	6035	5754	5490	5241	5005	4782	4570	4367
58	7381	7002	6653	6331	6030	5749	5486	5237	5002	4778	4566	4364
59	7374	6996	6648	6325	6025	5745	5481	5233	4998	4775	4563	4361
60	7368	6990	6642	6320	6021	5740	5477	5229	4994	4771	4559	4357

	22	23	24	25	26	27	28	29	30	31	32	33
	1320	1380	1440	1500	1560	1620	1680	1740	1800	1860	1920	1980
0	4357	4164	3979	3802	3632	3468	3310	3158	3010	2868	2730	2596
1	4354	4161	3976	3799	3629	3465	3307	3155	3008	2866	2728	2594
2	4351	4158	3973	3796	3626	3463	3305	3153	3005	2863	2725	2592
3	4347	4155	3970	3793	3623	3460	3302	3150	3003	2861	2723	2590
4	4344	4152	3967	3791	3621	3457	3300	3148	3001	2859	2721	2588
5	4341	4149	3964	3788	3618	3454	3297	3145	2998	2856	2719	2585
6	4338	4145	3961	3785	3615	3452	3294	3143	2996	2854	2716	2583
7	4334	4142	3958	3782	3612	3449	3292	3140	2993	2852	2714	2581
8	4331	4139	3955	3779	3610	3446	3289	3138	2991	2849	2712	2579
9	4328	4136	3952	3776	3607	3444	3287	3135	2989	2847	2710	2577
10	4325	4133	3949	3773	3604	3441	3284	3133	2986	2845	2707	2574
11	4321	4130	3946	3770	3601	3438	3282	3130	2984	2842	2705	2572
12	4318	4127	3943	3768	3598	3436	3279	3128	2981	2840	2703	2570
13	4315	4124	3940	3765	3596	3433	3276	3125	2979	2838	2701	2568
14	4311	4120	3937	3762	3593	3431	3274	3123	2977	2835	2698	2566
15	4308	4117	3934	3759	3590	3428	3271	3120	2974	2833	2696	2564
16	4305	4114	3931	3756	3587	3425	3269	3118	2972	2831	2694	2561
17	4302	4111	3928	3753	3585	3423	3266	3115	2969	2828	2692	2559
18	4298	4108	3925	3750	3582	3420	3264	3113	2967	2826	2689	2557
19	4295	4105	3922	3747	3579	3417	3261	3110	2965	2824	2687	2555
20	4292	4102	3919	3745	3576	3415	3259	3108	2962	2821	2685	2553
21	4289	4099	3917	3742	3574	3412	3256	3105	2960	2819	2683	2551
22	4285	4096	3914	3739	3571	3409	3253	3103	2958	2817	2681	2548
23	4282	4092	3911	3736	3568	3407	3251	3101	2955	2815	2678	2546
24	4279	4089	3908	3733	3565	3404	3248	3098	2953	2812	2676	2544
25	4276	4086	3905	3730	3563	3401	3246	3096	2950	2810	2674	2542
26	4273	4083	3902	3727	3560	3399	3243	3093	2948	2808	2672	2540
27	4269	4080	3899	3725	3557	3396	3241	3091	2946	2805	2669	2538
28	4266	4077	3896	3722	3555	3393	3238	3088	2943	2803	2667	2535
29	4263	4074	3893	3719	3552	3391	3236	3086	2941	2801	2665	2533
30	4260	4071	3890	3716	3549	3388	3233	3083	2939	2798	2663	2531
31	4256	4068	3887	3713	3546	3386	3231	3081	2936	2796	2660	2529
32	4253	4065	3884	3710	3544	3383	3228	3078	2934	2794	2658	2527
33	4250	4062	3881	3708	3541	3380	3225	3076	2931	2792	2656	2525
34	4247	4059	3878	3705	3538	3378	3223	3073	2929	2789	2654	2522
35	4244	4055	3875	3702	3535	3375	3220	3071	2927	2787	2652	2520
36	4240	4052	3872	3699	3533	3372	3218	3069	2924	2785	2649	2518
37	4237	4049	3869	3696	3530	3370	3215	3066	2922	2782	2647	2516
38	4234	4046	3866	3693	3527	3367	3213	3064	2920	2780	2645	2514
39	4231	4043	3863	3691	3525	3365	3210	3061	2917	2778	2643	2512
40	4228	4040	3860	3688	3522	3362	3208	3059	2915	2775	2640	2510
41	4224	4037	3857	3685	3519	3359	3205	3056	2912	2773	2638	2507
42	4221	4034	3855	3682	3516	3357	3203	3054	2910	2771	2636	2505
43	4218	4031	3852	3679	3514	3354	3200	3052	2908	2769	2634	2503
44	4215	4028	3849	3677	3511	3351	3198	3049	2905	2766	2632	2501
45	4212	4025	3846	3674	3508	3349	3195	3047	2903	2764	2629	2499
46	4209	4022	3843	3671	3506	3346	3193	3044	2901	2762	2627	2497
47	4205	4019	3840	3668	3503	3344	3190	3042	2898	2760	2625	2494
48	4202	4016	3837	3665	3500	3341	3188	3039	2896	2757	2623	2492
49	4199	4013	3834	3663	3497	3338	3185	3037	2894	2755	2621	2490
50	4196	4010	3831	3660	3495	3336	3183	3034	2891	2753	2618	2488
51	4193	4007	3828	3657	3492	3333	3180	3032	2889	2750	2616	2486
52	4189	4004	3825	3654	3489	3331	3178	3030	2887	2748	2614	2484
53	4186	4001	3822	3651	3487	3328	3175	3027	2884	2746	2612	2482
54	4183	3998	3820	3649	3484	3325	3173	3025	2882	2744	2610	2480
55	4180	3995	3817	3646	3481	3323	3170	3022	2880	2741	2607	2477
56	4177	3991	3814	3643	3479	3320	3168	3020	2877	2739	2605	2475
57	4174	3988	3811	3640	3476	3318	3165	3018	2875	2737	2603	2473
58	4171	3985	3808	3637	3473	3315	3163	3015	2873	2735	2601	2471
59	4167	3982	3805	3635	3471	3313	3160	3013	2870	2732	2599	2469
60	4164	3979	3802	3632	3468	3310	3158	3010	2868	2730	2596	2467

	34	35	36	37	38	39	40	41	42	43	44	45
"	2040	2100	2160	2220	2280	2340	2400	2460	2520	2580	2640	2700
0	2467	2341	2218	2099	1984	1871	1761	1654	1549	1447	1347	1249
1	2465	2339	2216	2098	1982	1869	1759	1652	1547	1445	1345	1248
2	2462	2337	2214	2096	1980	1867	1757	1650	1546	1443	1344	1246
3	2460	2335	2212	2094	1978	1865	1755	1648	1544	1442	1342	1245
4	2458	2333	2210	2092	1976	1863	1754	1647	1542	1440	1340	1243
5	2456	2331	2208	2090	1974	1862	1752	1645	1540	1438	1339	1241
6	2454	2328	2206	2088	1972	1860	1750	1643	1539	1437	1337	1240
7	2452	2326	2204	2086	1970	1858	1748	1641	1537	1435	1335	1238
8	2450	2324	2202	2084	1968	1856	1746	1640	1535	1433	1334	1237
9	2448	2322	2200	2082	1967	1854	1745	1638	1534	1432	1332	1235
10	2445	2320	2198	2080	1965	1852	1743	1636	1532	1430	1331	1233
11	2443	2318	2196	2078	1963	1850	1741	1634	1530	1428	1329	1232
12	2441	2316	2194	2076	1961	1849	1739	1633	1528	1427	1327	1230
13	2439	2314	2192	2074	1959	1847	1737	1631	1527	1425	1326	1229
14	2437	2312	2190	2072	1957	1845	1736	1629	1525	1423	1324	1227
15	2435	2310	2188	2070	1955	1843	1734	1627	1523	1422	1322	1225
16	2433	2308	2186	2068	1953	1841	1732	1626	1522	1420	1321	1224
17	2431	2306	2184	2066	1951	1839	1730	1624	1520	1418	1319	1222
18	2429	2304	2182	2064	1950	1838	1728	1622	1518	1417	1317	1221
19	2426	2302	2180	2062	1948	1836	1727	1620	1516	1415	1316	1219
20	2424	2300	2178	2061	1946	1834	1725	1619	1515	1413	1314	1217
21	2422	2298	2176	2059	1944	1832	1723	1617	1513	1412	1313	1216
22	2420	2296	2174	2057	1942	1830	1721	1615	1511	1410	1311	1214
23	2418	2294	2172	2055	1940	1828	1719	1613	1510	1408	1309	1213
24	2416	2291	2170	2053	1938	1827	1718	1612	1508	1407	1308	1211
25	2414	2289	2169	2051	1936	1825	1716	1610	1506	1405	1306	1209
26	2412	2287	2167	2049	1934	1823	1714	1608	1504	1403	1304	1208
27	2410	2285	2165	2047	1933	1821	1712	1606	1503	1402	1303	1206
28	2408	2283	2163	2045	1931	1819	1711	1605	1501	1400	1301	1205
29	2405	2281	2161	2043	1929	1817	1709	1603	1499	1398	1300	1203
30	2403	2279	2159	2041	1927	1816	1707	1601	1498	1397	1298	1201
31	2401	2277	2157	2039	1925	1814	1705	1599	1496	1395	1296	1200
32	2399	2275	2155	2037	1923	1812	1703	1598	1494	1393	1295	1198
33	2397	2273	2153	2035	1921	1810	1702	1596	1493	1392	1293	1197
34	2395	2271	2151	2033	1919	1808	1700	1594	1491	1390	1291	1195
35	2393	2269	2149	2032	1918	1806	1698	1592	1489	1388	1290	1193
36	2391	2267	2147	2030	1916	1805	1696	1591	1487	1387	1288	1192
37	2389	2265	2145	2028	1914	1803	1694	1589	1486	1385	1287	1190
38	2387	2263	2143	2026	1912	1801	1693	1587	1484	1383	1285	1189
39	2384	2261	2141	2024	1910	1799	1691	1585	1482	1382	1283	1187
40	2382	2259	2139	2022	1908	1797	1689	1584	1481	1380	1282	1186
41	2380	2257	2137	2020	1906	1795	1687	1582	1479	1378	1280	1184
42	2378	2255	2135	2018	1904	1794	1686	1580	1477	1377	1278	1182
43	2376	2253	2133	2016	1903	1792	1684	1578	1476	1375	1277	1181
44	2374	2251	2131	2014	1901	1790	1682	1577	1474	1373	1275	1179
45	2372	2249	2129	2012	1899	1788	1680	1575	1472	1372	1274	1178
46	2370	2247	2127	2010	1897	1786	1678	1573	1470	1370	1272	1176
47	2368	2245	2125	2009	1895	1785	1677	1571	1469	1368	1270	1174
48	2366	2243	2123	2007	1893	1783	1675	1570	1467	1367	1269	1173
49	2364	2241	2121	2005	1891	1781	1673	1568	1465	1365	1267	1171
50	2362	2239	2119	2003	1889	1779	1671	1566	1464	1363	1266	1170
51	2359	2237	2117	2001	1888	1777	1670	1565	1462	1362	1264	1168
52	2357	2235	2115	1999	1886	1775	1668	1563	1460	1360	1262	1167
53	2355	2233	2113	1997	1884	1774	1666	1561	1459	1359	1261	1165
54	2353	2231	2111	1995	1882	1772	1664	1559	1457	1357	1259	1163
55	2351	2229	2109	1993	1880	1770	1663	1558	1455	1355	1257	1162
56	2349	2227	2107	1991	1878	1768	1661	1556	1454	1354	1256	1160
57	2347	2225	2105	1989	1876	1766	1659	1554	1452	1352	1254	1159
58	2345	2223	2103	1987	1875	1765	1657	1552	1450	1350	1253	1157
59	2343	2220	2101	1986	1873	1763	1655	1551	1449	1349	1251	1156
60	2341	2218	2099	1984	1871	1761	1654	1549	1447	1347	1249	1154



'	46	47	48	49	50	51	52	53	54	55	56	57	58	59
"	2760	2820	2880	2940	3000	3060	3120	3180	3240	3300	3360	3420	3480	3540
0	1154	1061	0969	0880	0792	0706	0621	0539	0458	0378	0300	0223	0147	0073
1	1152	1059	0968	0878	0790	0704	0620	0537	0456	0377	0298	0221	0145	0072
2	1151	1057	0966	0877	0789	0703	0619	0536	0455	0375	0297	0220	0145	0071
3	1149	1056	0965	0875	0787	0702	0617	0535	0454	0374	0296	0219	0142	0069
4	1148	1054	0963	0874	0786	0700	0616	0533	0452	0373	0294	0218	0142	0068
5	1146	1053	0962	0872	0785	0699	0615	0532	0451	0371	0293	0216	0141	0067
6	1145	1051	0960	0871	0783	0697	0613	0531	0450	0370	0292	0215	0140	0066
7	1143	1050	0959	0869	0782	0696	0612	0529	0448	0369	0291	0214	0139	0064
8	1141	1048	0957	0868	0780	0694	0610	0528	0447	0367	0289	0213	0137	0063
9	1140	1047	0956	0866	0779	0693	0609	0526	0446	0366	0288	0211	0136	0062
10	1138	1045	0954	0865	0777	0692	0608	0525	0444	0365	0287	0210	0135	0061
11	1137	1044	0953	0863	0776	0690	0606	0524	0443	0363	0285	0209	0134	0060
12	1135	1042	0951	0862	0774	0689	0605	0522	0442	0362	0284	0208	0132	0058
13	1134	1041	0950	0860	0773	0687	0603	0521	0440	0361	0283	0206	0131	0057
14	1132	1039	0948	0859	0772	0686	0602	0520	0439	0359	0282	0205	0130	0056
15	1130	1037	0947	0857	0770	0685	0601	0518	0438	0358	0280	0204	0129	0055
16	1129	1036	0945	0856	0769	0683	0599	0517	0436	0357	0279	0202	0127	0053
17	1127	1034	0944	0855	0767	0682	0598	0516	0435	0356	0278	0201	0126	0052
18	1126	1033	0942	0853	0766	0680	0596	0514	0434	0354	0276	0200	0125	0051
19	1124	1031	0941	0852	0764	0679	0595	0513	0432	0353	0275	0199	0124	0050
20	1123	1030	0939	0850	0763	0678	0594	0512	0431	0352	0274	0197	0122	0049
21	1121	1028	0938	0849	0762	0676	0592	0510	0430	0350	0273	0196	0121	0047
22	1119	1027	0936	0847	0760	0675	0591	0509	0428	0349	0271	0195	0120	0046
23	1118	1025	0935	0846	0759	0673	0590	0507	0427	0348	0270	0194	0119	0045
24	1116	1024	0933	0844	0757	0672	0588	0506	0426	0346	0269	0192	0117	0044
25	1115	1022	0932	0843	0756	0670	0587	0505	0424	0345	0267	0191	0116	0042
26	1113	1021	0930	0841	0754	0669	0585	0503	0423	0344	0266	0190	0115	0041
27	1112	1019	0929	0840	0753	0668	0584	0502	0422	0342	0265	0189	0114	0040
28	1110	1018	0927	0838	0751	0666	0583	0501	0420	0341	0264	0187	0112	0039
29	1109	1016	0926	0837	0750	0665	0581	0499	0419	0340	0262	0186	0111	0038
30	1107	1015	0924	0835	0749	0663	0580	0498	0418	0339	0261	0185	0110	0036
31	1105	1013	0923	0834	0747	0662	0579	0497	0416	0337	0260	0184	0109	0035
32	1104	1012	0921	0833	0746	0661	0577	0495	0415	0336	0258	0182	0107	0034
33	1102	1010	0920	0831	0744	0659	0576	0494	0414	0335	0257	0181	0106	0033
34	1101	1008	0918	0830	0743	0658	0574	0493	0412	0333	0256	0180	0105	0031
35	1099	1007	0917	0828	0741	0656	0573	0491	0411	0332	0255	0179	0104	0030
36	1098	1005	0915	0827	0740	0655	0572	0490	0410	0331	0253	0177	0103	0029
37	1096	1004	0914	0825	0739	0654	0570	0489	0408	0329	0252	0176	0101	0028
38	1095	1002	0912	0824	0737	0652	0569	0487	0407	0328	0251	0175	0100	0027
39	1093	1001	0911	0822	0736	0651	0568	0486	0406	0327	0250	0174	0099	0025
40	1091	0999	0909	0821	0734	0649	0566	0484	0404	0326	0248	0172	0098	0024
41	1090	0998	0908	0819	0733	0648	0565	0483	0403	0324	0247	0171	0096	0023
42	1088	0996	0906	0818	0731	0647	0563	0482	0402	0323	0246	0170	0095	0022
43	1087	0995	0905	0816	0730	0645	0562	0480	0400	0322	0244	0169	0094	0021
44	1085	0993	0903	0815	0729	0644	0561	0479	0399	0320	0243	0167	0093	0019
45	1084	0992	0902	0814	0727	0642	0559	0478	0398	0319	0242	0166	0091	0018
46	1082	0990	0900	0812	0726	0641	0558	0476	0396	0318	0241	0165	0090	0017
47	1081	0989	0899	0811	0724	0640	0557	0475	0395	0316	0239	0163	0089	0016
48	1079	0987	0897	0809	0723	0638	0555	0474	0394	0315	0238	0162	0088	0015
49	1078	0986	0896	0808	0721	0637	0554	0472	0392	0314	0237	0161	0087	0013
50	1076	0984	0894	0806	0720	0635	0552	0471	0391	0313	0235	0160	0085	0012
51	1074	0983	0893	0805	0719	0634	0551	0470	0390	0311	0234	0158	0084	0011
52	1073	0981	0891	0803	0717	0633	0550	0468	0388	0310	0233	0157	0083	0010
53	1071	0980	0890	0802	0716	0631	0548	0467	0387	0309	0232	0156	0082	0008
54	1070	0978	0888	0801	0714	0630	0547	0466	0386	0307	0230	0155	0080	0007
55	1068	0977	0887	0799	0713	0628	0546	0464	0384	0306	0229	0153	0079	0006
56	1067	0975	0885	0798	0711	0627	0544	0463	0383	0305	0228	0152	0078	0005
57	1065	0974	0884	0796	0710	0626	0543	0462	0382	0304	0227	0151	0077	0004
58	1064	0972	0883	0795	0709	0624	0541	0460	0381	0302	0225	0150	0075	0002
59	1062	0971	0881	0793	0707	0623	0540	0459	0379	0301	0224	0148	0074	0001
60	1061	0969	0880	0792	0706	0621	0539	0458	0378	0300	0223	0147	0073	0000






THE SOLAR SYSTEM

*Relative Magnitudes of the Planets.*

Jupiter .  
 Saturn .  
 Uranus .  
 Neptune .  
 Mars .  
 Venus .  
 Earth .  
 Mercury .



*Relative Magnitudes of the sun seen from each Planet*

from Mars  
 from Saturn  
 from Jupiter  
 from Neptune  
 from Uranus  
 from Venus  
 from Earth  
 from Mercury




























Fig 11















37



































A 000 210 521 1

N82e

AUXILIARY  
STACK

JUL 72

QB  
43  
N82e



